is densely covered with green columnar emergences, whose cortical cells contain chlorophyll, starch, tannin, and a substance giving the reactions of a fat. Irregular masses of a similarly reacting material are frequently found covering the cuticle.

A *single* stoma occurs on the free surface of each emergence, and in the embryo of this species stomata are confined to the emergences.

The cuticle covering the general epidermis is continuous over the guard-cells of each stoma, except for a small oval slit which allows of communication between the intercellular space and the air. The stomata thus suggest either a xerophytic habit for the plant or an abnormal function for themselves.

The emergences flourish during the germinating (epiphytic) stage, and later, when semi-parasitism is achieved, cease to be functional.

II. Mode of penetration into the host of L. neelgherensis.—Unlike many species, *L. neelgherensis* develops no well-marked organ of attachment (suctorial disc) at the free end of its hypocotyl.

Where much resistance to the entry of the sucker is offered by the host, there are formed at the edges of the attached surface of the hypocotyl a series of acropetally arising, hair-bearing cortical ridges. The later-formed ridges, wedging themselves in between the older ones and the bark, force these older ridges away. The firmly-attached hairs of each ridge so forced away tear off masses of the bark, and thus the softer tissues, through which the sucker readily and cleanly bores, are exposed by instalments. Where the sucker comes in contact with lignified structures, dissolution is more gradual, and stages of disintegration (erosion figures) are to be observed.

In *L. loniceroides*, where a well-marked suctorial disc is formed, attachment occurs once for all. This attachment is maintained (1) by the growth of the edge of the disc hard against the bark; (2) by the outgrowing hairs forming a matted sclerotic mass firmly fixed into the outer layers of the host.

F. W. KEEBLE.

PRELIMINARY NOTES ON FLORAL DEVIATIONS IN SOME SPECIES OF POLYGONUM¹.—The genus has long been known to show considerable departures from the arrangement and number of parts accepted as most typical (Per. 5. St. 5 + 3, C. 3), such as is found in *P. Convolvulus*. Eichler's 'Blüthendiagramme,'

¹ Abstract of a paper read before Section K at the Liverpool meeting of the British Association, 1896.

for example, shows diagrams of several species as if characterized by constant differences of structures. Observation shows that in some species (e.g. *P. Convolvulus*) variations are comparatively infrequent and slight, but that in most (e.g. *P. Persicaria* and *aviculare*) they are extremely frequent, and lead to very great changes in floral structure. Often it is scarcely possible in such species to find two flowers alike on the same branch, or even on the same plant. Within a species individual plants show wide differences in the frequency and extent of variations.

A comparison of different species shows that while each varies, so as in the more variable species to cover almost the whole range observed in the genus, each shows a tendency to certain lines of variation. These tendencies are more alike usually in the more nearly allied species, so as to correspond in the main with the groups based on habit, and they lead from group to group.

The modes of variation commonly observed include almost all the recognized modes of departure from floral symmetry. They affect all the whorls. The *perianth* in some species is very constant. In others it habitually shows cohesion of two or more segments, or abortion in different degrees, or suppression of one or two (usually the inner) segments. Choris of a segment is less frequent. Enations from one or more segments are frequent in certain species, rare or absent in others. The *outer stamens* often show cohesion of the two in each pair, varying from the slightest union of the bases of the filaments to absolute union of even the anthers. Abortion (in all degrees to complete suppression) of one or more stamens is not rare, frequently reducing this whorl to $\bar{3}$ (less often to 2) in *aviculare*. Choris is not rare, especially of the unpaired stamen. The *inner stamens* seldom show cohesion (except in *aviculare* and its allies) with stamens of the outer whorl. Abortion (in all degrees to complete suppression) is very frequent, and in certain species (*amphibium*) this whorl has completely disappeared. In *aviculare* and allied species the inner whorl shows abortion less than the outer. Choris in the inner whorl most frequently shows itself in the posterior stamen. Adhesions of stamens to perianth segments and petalody of stamens are not frequent.

(In *P. amphibium*, the land form near Aberdeen very generally has the anthers very small or abortive, and the stamens hidden within the perianth, while the form growing in water has the anthers well

developed, and some or all exserted; neither form appears to seed habitually.)

The *pistil* in some species is very constant, while in others it shows all stages of cohesion and reduction to two carpels, this being the almost invariable number in certain species. Abortion is less frequent, and complete suppression cannot be distinguished from complete cohesion. Chorisis is very frequent in *aviculare* and some other species, in all degrees from a mere enlargement of one or more stigmas to an increase in number (up to seven), with corresponding modifications in structure in the ovary. Only one ovule has been observed in each ovary.

Markedly teratological forms have been met with, but are not included in this summary.

No very definite relation has been traced between the position of a flower on the axis and deviations in structure, though pressure tends to abortion or suppression of parts, especially of the sexual organs. (The flowers examined have chiefly been those sufficiently open to allow the natural arrangement to be noted without manipulation, to avoid displacement of parts, hence cleistogamous flowers are scarcely included.) The variability appears rather to express the result of an innate tendency to vary where not subject to the check of loss of fertility, the variations in *Polygonum* not leading to this loss.

The same number of parts in a whorl may be due to very different causes, and still more may the same number of stamens express very different arrangements in the flower; hence such a statement in a specific description as 'stamens usually six' is insufficient.

J. W. H. TRAIL.

Trail, J. W. H. 1896. "Preliminary notes on floral deviations in some species of *Polygonum*." *Annals of botany* 10, 627–629.

<https://doi.org/10.1093/oxfordjournals.aob.a088635>.

View This Item Online: <https://www.biodiversitylibrary.org/item/235772>

DOI: <https://doi.org/10.1093/oxfordjournals.aob.a088635>

Permalink: <https://www.biodiversitylibrary.org/partpdf/318447>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.