material for the early stages of spore-formation, I pull up a plant on which one ear or more is smutted, and dissect out the ears of the eight or ten or more shoots that so far have no ears visible externally. Once I found a plant with two ears smutted and six or eight sound, but this was the only time that all the ears on a plant were not equally affected; and once I found an ear with the three lowest kernels alone smutted while all those above were normal. To keep up my stock of spores for germination in class-work, I gather the smutted ears before the spores are shed, and keep them in small tin tobacco-boxes. On one occasion a small beetle got shut up too, and was imprisoned for a week or two before I found and released it. I noticed many spores adhering in short threads, really the excreta of the said insect. I tried the germination of the excretal and normal spores in a culture-cell (made by fastening a glass ring to a slide and growing the spores in a hanging drop of water on the cover-slip inverted on the ring), and found the former to germinate in about half the time required for the latter. For this summer's material I smutted some Barley with spores taken from Barley (Row 5 of my plot at the Botanic Garden), some with spores from Oats (Row 7), and some with spores from Wheat (which, however, gave no result). Omitting details, I may say that I found 70-80 % only of the grain grew, and I gathered eighty smutted ears from Row 5 and forty-six from Row 7, amounting to 37 % and 21 % respectively of the total number of ears produced.

Phytophthora infestans.—As Potatoes are not to be absolutely depended upon, and as growing and infecting successive sets of potato-tops under bell-jars involves much space and time, I have for the past three years used *Solanum laciniatum* as class-material. I can rely on finding *Phytophthora infestans* upon it and in good condition, and its petioles and midribs furnish material much superior to potato-leaves. I prefer to gather all my *Phytophthora* material before the dew is off, and I keep it in a vasculum until my class at 11.0, when I use it fresh.

W. G. P. ELLIS.

BOTANICAL LABORATORY, CAMBRIDGE.

THE FUNCTIONS OF LATEX.—Although several attempts have been made to ascertain the functions of latex in plants, they have not led to any very conclusive information respecting it. Accordingly, at the suggestion of Prof. Marshall Ward, I undertook an

examination of the subject from several points of view. My apology for bringing forward the results in so incomplete a state is that, at present, I have not the time at my disposal to complete it, owing to my departure for the tropics, and that it may possibly prove suggestive enough to excite interest in the problem and so help in its solution.

The work of Schullerus¹, Treub², and Faivre³ is too well known to need description, as is also Schimper's⁴ criticism of it.

Haberlandt's work⁵ is disputed, but as he still maintains his original views in the last edition of his *Physiologische Pflanzenanatomie*, I have thought it worth while to repeat it.

In leaves in which the palisade-tissue is loosely packed, as in *Euphorbia mellifera* and *Euphorbia punicea*, frequent examples of the palisade-cells converging to the blind apex of a laticiferous cell, or a cell in direct connexion with it, can be met with.

In *Euphorbia spinosa*, where the palisade-layer is double, the branches of the laticiferous tubes run between the two layers and send out terminal branches into the upper layer of palisade-cells, thus forming a kind of brush-work. The blind endings are usually on the upper surface of the palisade-cells.

These terminations may be traced by mounting the sections in phenyl-hydrazine, the high refractive power of which renders it a very useful clearing agent. At the same time a marked staining action occurs in the palisade-cells and in the cells in which Fry⁶ has described 'proteid aggregates,' which may be worth further investigation.

In laticiferous plants whose assimilation is carried on by the stem owing to partial or complete reduction of the leaf-surface, the assimilating cells form a compact layer several cells in thickness, e. g. *Euphorbia arborescens* and *coerulescens*. The laticiferous cells pass directly through this layer and end, for the most part, on the upper surface of the outermost layer of assimilating cells.

Taking all observed cases into consideration, I think we may safely

¹ Schullerus, Abhandl. d. bot. Vereins d. Prov. Brandenb., 1882.

² Treub, Anns. du Jardin Bot. de Buitenzerg, vol. iii, p. 37.

³ Faivre, Ann. des Sci. Nat., sér. 5, t. x. p. 33. C. R. t. lxxxviii. p. 269, etc.

⁴ Schimper, Bot. Zeit., 1885, p. 771 et seq.

⁵ Haberlandt, Sitzb. d. k. Akad. Wien, 1883, p. 51.

⁶ Fry, Annals of Botany, Vol. v, p. 413.

say that the blind endings of the laticiferous system are generally connected with the palisade-cells.

If it is improbable that a plant would build up so elaborate a system for the storage of waste products, it is possible that this method of terminating among cells in which assimilation is carried on may point to a storage or means of transit of reserve products.

This is further borne out by the extremely early appearance of the system in the leaf. A longitudinal section through a stem-apex of *Euphorbia pulcherrima*, for example, shows, in leaves just beginning to unfold, a well-developed system with its earliest branches extending to the palisade-layer.

Again, if latex is simply waste material, old leaves should contain large quantities of it, but this is by no means the case.

If young leaves and old leaves are lightly punctured, the young leaves exude far more latex than the old ones, and the same is true for young and old shoots.

Micro-chemical tests give the same results. If starch is present in the latex, a section of a young leaf shows the tubes well stored with it; while in an old or a fallen leaf it is difficult to find a trace, even if the iodine-method of staining is used. The starch has evidently been handed on from the laticiferous system to other parts of the plant.

Testing for proteids with Millons' reagent, or by the xanthoproteic reaction, gives completely parallel results. *Ficus Carica* is a possible exception, for here one occasionally finds granular masses in the tubes which give proteid reactions.

Though caoutchouc may be met with in fallen leaves, yet, from what is known of its chemistry, it is hardly likely that this substance has any nutrient value in the plant; but there is a possibility that, as do many other terpenes, it may absorb oxygen as ozone. When the endings of the tubes, and Schunck and Brebner's¹ demonstration of the presence of ozone in the palisade-layer, are taken into consideration, this view becomes more tenable. Direct proof of this has not been obtained, though testing latex with Wurster's papers and with mercury points to the presence of minute quantities of free ozone. I hope to obtain evidence upon this point, among others, while examining the india-rubber industry in East Africa.

To determine whether any physiological connexion exists between

¹ Schunck and Brebner, *Annals of Botany*, Vol. vi, p. 167.

the laticiferous system and the assimilating tissues, chemical methods had to be applied.

So far the chemical side of the question appears to have been badly neglected, although it was attacked so long ago by v. Mohl¹. We may pass over the few analyses made of latex yielding economic products, and briefly notice that Weiss and Wiesner² have shown the presence of sugars in latex, and that Green³ has examined its proteids.

As the sugars are so intimately connected with assimilation, it was decided to estimate them in latex drawn from plants under varying conditions with regard to their capability for assimilation.

The method employed was to take a known quantity of latex and remove the proteids by precipitation with excess of alcohol. After standing for a day, the solution was filtered off and evaporated to dryness over a water-bath. The sugars were then extracted from the resinous mass with warm water and estimated with Fehling's solution. The precipitate was weighed as CuO. With ordinary care this method gives very constant results, which is a point of considerable value when working with small quantities of latex. Where tannins are present they should be extracted with hide powder or ethyl acetate; but as the process leads to a certain amount of loss, it is as well to use latex free from them when possible.

We will first examine the results obtained when assimilation is checked by darkening the plants. Four plants of *Euphorbia pulcherrima* were placed in a dark room, and another set were kept in a greenhouse at a subtropical temperature.

The undarkened plants showed the presence of sugars equivalent to .021 grms. CuO per c.c. After being darkened for a day the amount of sugar fell to .002 grms. CuO per c.c., and after another day no trace could be found. *Euphorbia arborescens* gave the same results.

Judging solely from the bulk of the proteid precipitate, the proteids had also undergone a considerable diminution in quantity.

If one may dismiss pathological changes due to abnormal conditions in this case, we may say that there is some definite connexion

¹ v. Mohl, Bot. Zeit., 1843, p. 553 et seq.

² Weiss and Wiesner, Bot. Zeit., 1861, p. 41; and 1862, p. 125.

³ Green, Proc. Roy. Soc., 1886, No. 242, p. 28.

between the laticiferous system and the assimilating tissues. Further experiments confirm this conclusion; for if the plants are grown under normal conditions and the latex examined before and after the day's assimilation, an increase is found at the end of the period, as is shown by the following experiments made with *Euphorbia Peplus* in the early part of November, 1896, e. g.:—

Nov. 3, 1896, a bright sunny day—

Latex collected at 10 A.M.	gave	·0302	grms. CuO per c.c.
„ „ „ 4 P.M.	„	·036	„ „ „

A sharp frost put an end to this series of experiments. The result of exposure to this cold was that the plants gave no exudation of latex on wounding even when brought into a warm room; on applying pressure a thin opalescent liquid was squeezed out.

Further experiments were accordingly carried out with *Euphorbia pulcherrima*, with the following results:—

Collected at 10 A.M.	Collected at 4 P.M.	
·032	·049	grs. CuO per c.c.
·030	·046	„ „
·033	·046	„ „
·030	·047	„ „

Here an increase of sugars after the day's assimilation is evident. With *Euphorbia arborescens* very discordant results were obtained:—

Collected at 10 A.M.	Collected at 4 P.M.	
·032	·032	grs. CuO per c.c.
·034	·0345	„ „
·0185	·013	„ „
·012	·018	„ „
·021	·0195	„ „

The discrepancies in the foregoing results were found to be due to the fact that the latex had been drawn from different parts of the plant, the younger portions being richer in sugar than the older.

When we consider that some of the chief constituents of latex are starch, proteids, and sugar, all substances of undoubted nutrient value, and that the proteids are typical circulatory forms, and that the quantities of sugar vary with the plant's assimilation, it is evident that one function of latex is to carry reserve-materials in the plant.

Whether caoutchouc, resins, and tannins have any value, nutrient or otherwise, in the plant still remains to be determined. If they prove to be simply waste products, we shall be confronted with a peculiar method of dealing with waste matter which does not show much analogy to methods known at present.

R. H. BIFFEN.

FRANK SMART STUDENT,
GONVILLE AND CAIUS COLLEGE, CAMBRIDGE.

ON PEZIZA AURANTIA.—In March, 1896, the large pond in the Botanic Gardens, Cambridge, was emptied and cleaned, and large quantities of mud, chiefly the peculiar blue clay—gault—so common here, were removed from the bottom, together with plant-remains, such as the rhizomes and stems of water-lilies, *Epilobium*, *Scirpus*, *Polygonum*, &c. This mud was put into heaps in the garden under the shade of trees. At the same time a heap of similar blue clay, obtained during the digging for sewerage operations in the streets outside the gardens, was made near these heaps of pond-mud.

In September we noticed the apparently sudden outburst of enormous numbers of brilliant orange-scarlet *Peziza* cups on the heaps from the pond, and these increased during October and November, forming one of the most magnificent crops of these Fungi I have ever seen.

Examination showed the *Peziza* to be *P. aurantia* (Oed.), the cups, asci, and spores exhibiting the characters given in Phillips' British Discomycetes, p. 56, as those of the typical species, and reference to Masee (Brit. Fungus-Flora, iv, p. 448), Lindau (Engler, Pflanzenfam., 1 Th. 1 Abth. Lfg. 130, p. 187), and Winter (Rabenh., Die Pilze, B. III, p. 970) confirmed this, though these authors differ in the authorities cited for the name and synonyms.

Brefeld has not succeeded in germinating the spores of this species, and so I made attempts to cultivate it, and hoped to get out its life-history, but in vain.

It was quite easy to obtain the spores, though the majority of them appeared hardly ripe; nevertheless, numerous attempts to germinate them failed. In hanging-drops of water, gelatine culture-media, agar, &c., they did not stir under any of the given conditions, nor could I get them to grow in test-tubes on what experience suggested were probably suitable media.

Biffen, Roland H. 1897. "The function of latex." *Annals of botany* 11, 334–339.
<https://doi.org/10.1093/oxfordjournals.aob.a088654>.

View This Item Online: <https://www.biodiversitylibrary.org/item/233542>

DOI: <https://doi.org/10.1093/oxfordjournals.aob.a088654>

Permalink: <https://www.biodiversitylibrary.org/partpdf/318466>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.