Lastly, of the several theories thus far proposed to account for the occurrence of the abnormalities, none is capable of wide application.

Sometimes the abnormalities reappear year after year; sometimes they prove inconstant. Had we a fuller knowledge, some explanation, partial or complete, might be forthcoming; for frequently, both in their distribution in the catkin and on the branch, the changes in sex show a tendency to arrangement. At times the male is above the female; at times the reverse is the case. Rarely there are three or four belts of flowers on one catkin, male succeeding female, and female male, in definite order.

ROYAL GARDENS, KEW.

I. H. BURKILL.

THE ANATOMY OF THE STEM OF SPECIES OF LYCOPodium.—Ten species of *Lycopodium* have been examined; among these two types may be distinguished.

1. Type of *L. clavatum* (L.). The oval stelic arrangement is marked by a considerable amount of xylem, broken up into patches by bands of phloem. Centrally these bands are strap-shaped, but at the ends of the long axes the areas of phloem are external, and occur as curved and flattened wedges. Large cells without contents, sieve-tubes, appear in the centre of the strap-shaped bands. Protophloems and protoxylems are external, forming a continuous ring, as figured by Hofmeister; so that, using De Bary's terminology, the arrangement of the bundles is radial. Pericyclic and the so-called endodermal cells occur in concentric zones, 1-3 cells broad. The former swell up, especially in glycerine or glycerine-jelly; the latter are generally considerably lignified. The cells of the cortex lying just external to the endodermal cells are thickened and lignified, forming a third concentric zone several cells deep. To this type conform *L. alpinum* (L.), *L. Phlegmaria* (L.), *L. dendroides* (?), and *L. cernuum* (L.).

2. Type of *L. squarrosum*. The type which contrasts most markedly with the former is found in *L. squarrosum* (Forst.), *L. dichotomum* (Jacq.), and *L. nummularifolium* (Blume). The phloems occur as islands in the sea of xylem, or as inserted peninsulas. The phloems are centrally built up, with the apparent sieve-tubes in the centre. Protoxylems are well marked, and lie externally, but protophloems are not to be distinguished. Endodermal cells and pericycle are found as in the previous type. The sclerenchymatous sheath is wanting, or very slightly developed.

The two remaining species, *L. Dalhousieanum* (Spring) and *L. Selago* (L.), are, to some degree, intermediate types. The phloem in *L. Dalhousieanum* shows both types, strap-shaped and centric. In the branches the structure becomes simpler. There are two narrow strips of xylem, with an intermediate strip of phloem, so that a prominent row of sieve-tubes occupies the very centre of the stelic cylinder. *L. Selago* in its structure is modified on that of *L. clavatum*. An interesting feature of *L. Selago* and *L. squarrosus* is the occurrence of root-structures running through the stem. These consist of steles containing a crescent-shaped mass of xylem, with protoxylems towards each tip, while the concave portion is filled up with phloem. A characteristic sclerenchymatous sheath surrounds the stele. In *L. Selago* these root-structures are found even above the point where the stem branches, but in *L. squarrosus* they have fused with the central cylinder before branching occurs.

C. E. JONES.

UNIVERSITY COLLEGE, LIVERPOOL.

REPRODUCTION IN DICTYOTA DICHOTOMA.—1. *Dictyota* is an annual. In this country it germinates during the summer, remains small during the winter, grows very rapidly in June, and begins to form its reproductive cells in July.

2. The tetraspores are produced throughout the season, and all stages may be found together on the same plant. The sexual cells, however, show a remarkable periodicity. The formation, maturation, and liberation of each crop occupies a fortnight, the interval between two spring-tides. The sori are formed during neap-tides, and the cells are liberated during or immediately after the highest spring-tides.

3. When liberated the oospheres are not invested with walls. In this condition they strongly attract the antherozoids, become fertilized, and at once start germinating. The plantlets are similar to those figured by Thuret as resulting from the germination of the tetraspores.

4. If not fertilized the eggs lose the power of attracting antherozoids, they form walls, and, as already described by Thuret and Bornet, they germinate parthenogenetically. After one or a few divisions, sometimes accompanied by formation of a rhizoid-rudiment, the process stops and the plantlets die.

5. Towards the close of the season some sori fail to mature within

Jones, C E. 1898. "The anatomy of the stem of species of Lycopodium." *Annals of botany* 12, 558–559. <https://doi.org/10.1093/oxfordjournals.aob.a088709>.

View This Item Online: <https://www.biodiversitylibrary.org/item/233102>

DOI: <https://doi.org/10.1093/oxfordjournals.aob.a088709>

Permalink: <https://www.biodiversitylibrary.org/partpdf/318521>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.