GREAT-FOOTED HAWK, Wils. Amer. Orn., vol. ix. p. 120.

FALCO PEREGRINUS, Bonap. Syn., p. 27.

COMMON OR WANDERING FALCON, *Falco peregrinus*, Nutt. Man., vol. i. p. 53.

GREAT-FOOTED HAWK, *Falco peregrinus*, Aud. Amer. Orn., vol. i. p. 85; vol. v. p. 365.

FALCO PEREGRINUS, *Peregrine Falcon*, Swains. & Rieh., F. Bor. Amer., vol. ii. p. 23.

Bill blackish-blue at the tip, pale green at the base, cere oil-green; bare orbital space orange. Iris hazel. Feet lemon-yellow; claws brownish-black. Head and hind neck greyish-black, tinged with blue; the rest of the upper parts dark bluish-grey, indistinctly barred with deep brown. Quills blackish-brown, the inner webs marked with transverse elliptical spots of reddish-white. Tail greyish-brown, marked with about twelve bars, the last of which is broad, the rest diminishing in size and intensity of tint. Throat and fore-neck white; a broad band of blackish-blue from the angle of the mouth downwards; cheeks whitish-grey; sides, breast and thighs reddish-white, transversely marked with dark brown spots in longitudinal series. Under wing feathers whitish, transversely barred.

Length $16\frac{1}{2}$ inches; extent of wings 30; bill $1\frac{1}{8}$ along the ridge; tarsus $1\frac{7}{8}$, middle toe $2\frac{1}{2}$.

As the bird gets old, the colours of the upper parts acquire a lighter tint in the male, and sometimes the back is ash-grey; but in the female, they gradually assume a deeper hue.

PIGEON HAWK.

+ FALCO COLUMBARIUS, Linn.

PLATE XXI.—MALE AND FEMALE.

The Pigeon Hawk ranges very extensively over the United States, and extends its migrations far beyond their limits on either side. Mr. TOWNSEND found it on the Rocky Mountains, as well as along the shores of the Columbia River. Dr. RICHARDSON mentions it as not uncommon about York Factory, in latitude 57° , and it is not improbable that it wanders farther, as he speaks of having seen a small hawk on the north shore of Great Bear Lake, in latitude 66° , which may have been a male as small as the one represented in my plate. I found it very abundant in the Texas early in May, when I shot as many as five on a small island in a short time.

Mr. HUTCHINS's description of the eggs of this bird, which he says are

Pigeon Falcon.

Drawn from Nature by J.J. Audubon. F.R.S.F.L.S.

Lith.^d Printed & Col^d by J.T. Bowen. Philad^a

white, and from two to four in number, as well as the situation of its nest, as given in his Notes on the Hudson's Bay Birds, is greatly at variance with my own observations. The eggs in three instances, which occurred at Labrador, were five; they measured an inch and three-quarters in length, an inch and a quarter in breadth, and were rather elongated; their ground colour a dull yellowish-brown, thickly clouded with irregular blotches of dull dark reddish-brown. In that country they are laid about the first of June. In the beginning of July I found five in a nest that were ready to be hatched. The nests were placed on the top branches of the low firs peculiar to that country, about ten or twelve feet from the ground, and were composed of sticks, slightly lined with moss and a few feathers. At this season the old birds evinced great concern respecting their eggs or young, remaining about them, and shewing all the tokens of anger and vexation which other courageous species exhibit on similar occasions. The young are at first covered with yellowish down; but I had no opportunity of watching their progress, as all that were taken on board the Ripley died in a few days. This species also breeds in Nova Scotia and New Brunswick.

A male from the Texas. Length to end of tail $13\frac{1}{4}$ inches, to end of wings $11\frac{5}{12}$, to end of claws $11\frac{6}{12}$; extent of wings 26.

The mouth resembles that of the other Falcons; its breadth $\frac{9\frac{1}{2}}{12}$ ths. The tongue is short, $\frac{6}{12}$ ths long, fleshy, deeply emarginate and papillate at the base, broadly grooved above, the tip rounded and slightly emarginate. The œsophagus is $4\frac{1}{2}$ inches long, its width at the upper part half an inch. The stomach is very large, round, $1\frac{1}{2}$ inches in diameter, with a very thin muscular coat; its central tendons $\frac{4}{12}$ ths in diameter. The proventriculus is $\frac{9}{12}$ ths long; its glands very numerous, and cylindrical. The intestine is $26\frac{3}{4}$ inches long, $\frac{2\frac{1}{2}}{12}$ ths in its greatest diameter. There are merely two slight indications of cœca; and the cloaca is globular, with a diameter of 1 inch.

The trachea is $2\frac{3}{4}$ inches long, a little flattened; the rings 58, well ossified; its breadth at the upper part $\frac{3}{12}$ ths, at the lower $\frac{2}{12}$ ths. The contractor muscles cover the anterior surface entirely in the upper third, and are of moderate strength, as are the sterno-tracheales; a pair of inferior laryngeal muscles going to the membrane between the last tracheal and first bronchial half ring. The bronchial half rings are 15 and 18.

PIGEON HAWK, *Falco columbarius*, Wils. Amer. Orn., vol. ii. p. 107.

FALCO COLUMBARIUS, Bonap. Syn., p. 38.

PIGEON HAWK, *Falco columbarius*, Nutt. Man., vol. i. p. 60.

LITTLE CORPORAL HAWK, *Falco temerarius*, Nutt. Man., vol. i. p. 61. Adult Male.

FALCO COLUMBARIUS, *Pigeon Hawk*, Swains. and Rich. F. Bor. Amer., vol. ii. p. 35.

FALCO ÆSALON, *Merlin*, Swains. and Rich. F. Bor. Amer., vol. ii. p. 37.

PIGEON HAWK, *Falco columbarius*, Aud. Orn. Biog., vol. i. p. 466; Young, vol. i. p. 381, Male; vol. v. p. 368.

Wings from two to three inches shorter than the tail, on the middle feathers of which are five, on the lateral six, broad whitish bands. Adult male with the cere greenish-yellow, the feet pale orange, the upper parts light bluish-grey, each feather with a black central line; lower parts reddish or yellowish-white, the breast and sides with large oblong brown spots; tibial feathers light red, streaked with blackish-brown. Female with the cere and legs greenish-yellow, the upper parts dark greyish-brown, the lower pale red, spotted as in the male. Young with the head light reddish-brown, streaked with dusky, the upper parts brownish-grey, the feathers margined and spotted with pale red, throat white, lower parts pale red, streaked with brown. The tail-bands vary from pale red to white.

THE AMERICAN SPARROW-HAWK.

+ *FALCO SPARVERIUS*, Linn.

PLATE XXII.—MALE AND FEMALE.

We have few more beautiful Hawks in the United States than this active little species, and I am sure, none half so abundant. It is found in every district from Louisiana to Maine, as well as from the Atlantic shores to the western regions. Every one knows the Sparrow-Hawk, the very mention of its name never fails to bring to mind some anecdote connected with its habits, and, as it commits no depredations on poultry, few disturb it, so that the natural increase of the species experiences no check from man. During the winter months especially it may be seen in the Southern States about every old field, orchard, barn-yard, or kitchen-garden, but seldom indeed in the interior of the forest.

Beautifully erect, it stands on the highest fence-stake, the broken top of a tree, the summit of a grain stack, or the corner of the barn, patiently and silently waiting until it spy a mole, a field-mouse, a cricket, or a grasshopper, on which to pounce. If disappointed in its expectation, it leaves its stand and removes to another, flying low and swiftly until within a few yards of the spot on which it wishes to alight, when all of a sudden, and in the most graceful manner, it rises towards it and settles with incomparable firmness of manner, merely suffering its beautiful tail to vibrate gently for awhile, its wings being closed with the swiftness of thought. Its keen eye perceives something beneath, when down it darts, secures the object in its talons, returns to its stand, and devours its prey piece by piece. This done, the

Pigeon Falcon.

Drawn from Nature by J.J. Audubon. F.R.S.F.L.S.

Lith.^d Printed & Col^d by J.T. Bowen. Philad^a

Audubon, John James. 1840. "Pigeon Hawk, *Falco columbarius*, Linn. [Pl. 21]."
The birds of America : from drawings made in the United States and their territories 1, 88–90. <https://doi.org/10.5962/p.319134>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124833>

DOI: <https://doi.org/10.5962/p.319134>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319134>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.