feathers lighter, the outer web pale brownish-grey; fore part of neck and a portion of the breast and sides ash-grey, the rest of the lower parts pale yellow.

Male, $5\frac{2}{12}$, wing, $2\frac{5}{12}$.

Columbia River. Fur countries. Labrador. Newfoundland. Rare in the Atlantic States.

THE WHITE OAK.

QUERCUS PRINUS, *Willd.* Sp. Pl., vol. iv. p. 439. *Pursh*, Fl. Amer., vol. ii. p. 633.—QUERCUS PRINUS PALUSTRIS, *Mich.* Arbr. Forest. de L'Amér. Sept. vol. ii. p. 51. Pl. 7.—MONOCIA POLYANDRIA, *Linn.*—AMENTACEÆ, *Juss.*

Leaves oblongo-oval, acute, largely toothed, the teeth nearly equal, dilated, and callous at the tip; cupule craterate, attenuated at the base; acorn ovate. This species grows in low shady woods, and along the margins of rivers, from Pennsylvania to Florida. The wood is porous, and of inferior quality.

SMALL HEADED FLYCATCHER.

+ MUSCICAPA MINUTA, *Wilson.*

PLATE LXVII.—MALE.

The sight of the figure of this species brings to my recollection a curious incident of long-past days, when I drew it at Louisville in Kentucky. It was in the early part of the spring of 1808, thirty-two years ago, that I procured a specimen of it while searching the margins of a pond.

In those happy days, kind reader, I thought not of the minute differences by which one species may be distinguished from another in words, or of the necessity of comparing tarsi, toes, claws, and quills, although I have, as you are aware, troubled you with tedious details of this sort. When ALEXANDER WILSON visited me at Louisville, he found in my already large collection of drawings, a figure of the present species, which, being at that time unknown to him, he copied and afterwards published in his great work, but without acknowledging the privilege that had thus been granted to him. I have more than once regretted this, not by any means so much on my own ac-

Small-headed Flycatcher
Virginian Spider-wort. Tradescantia virginica?
 Male

 

count, as for the sake of one to whom we are so deeply indebted for his elucidation of our ornithology.

I consider this Flycatcher as among the scarcest of those that visit our middle districts; for, although it seems that WILSON procured one that "was shot on the 24th of April, in an orchard," and afterwards "several individuals of this species in various quarters of New Jersey, particularly in swamps," all my endeavours to trace it in that section of the country have failed, as have those of my friend EDWARD HARRIS, Esq., who is a native of that State, resides there, and is well acquainted with all the birds found in the district. I have never seen it out of Kentucky, and even there it is a very uncommon bird. In Philadelphia, Baltimore, New York, or farther eastward or southward, in our Atlantic districts, I never saw a single individual, not even in museums, private collections, or for sale in bird-stuffers' shops.

In its habits this species is closely allied to the Hooded and Green Blackcap Flycatchers, being fond of low thick coverts, whether in the interior of swamps, or by the margins of sluggish pools, from which it only removes to higher situations after a continuation of wet weather, when I have found it on rolling grounds, and amid woods comparatively clear of under-growth.

Differing from the true Flycatchers, this species has several rather pleasing notes, which it enunciates at pretty regular intervals, and which may be heard at the distance of forty or fifty yards in calm weather. I have more than once seen it attracted by an imitation of these notes. While chasing insects on wing, although it clicks its bill on catching them, the sound thus emitted is comparatively weak, as is the case with the species above mentioned, it being stronger however in the Green Blackcap than in this or the Hooded species. Like these birds, it follows its prey to some distance at times, whilst at others, it searches keenly among the leaves for its prey, but, I believe, never alights on the ground, not even for the purpose of drinking, which act it performs by passing lightly over the water and sipping, as it were, the quantity it needs.

All my efforts to discover its nest in the lower parts of Kentucky, where I am confident that it breeds, have proved fruitless; and I have not heard that any other person has been more successful.

SMALL-HEADED FLYCATCHER, *Muscicapa minuta*, Wils. Amer. Orn., vol. vi. p. 62.

SYLVIA MINUTA, Bonap. Syn., p. 86.

SMALL-HEADED SYLVAN FLYCATCHER, Nutt. Man., vol. i. p. 296.

SMALL-HEADED FLYCATCHER, *Muscicapa minuta*, Aud. Orn. Biog., vol. v. p. 291.

Wings short, the second quill longest; tail of moderate length, even; general colour of upper parts light greenish-brown; wings and tail dark olive-brown, the outer feathers of the latter with a terminal white spot on the inner

web; a narrow white ring surrounding the eye; two bands of dull white on the wing; sides of the head and neck greenish-yellow, the rest of the lower parts pale yellow, gradually fading into white behind.

Male, 5, $8\frac{1}{4}$.

Kentucky, Pennsylvania, and New Jersey. Exceedingly rare. Migratory.

THE VIRGINIAN SPIDER-WORT.

TRADESCANTIA VIRGINICA, Willd. Sp. Pl., vol. ii. p. 16. Pursh, Fl. Amer., vol. i. p. 218.

—HEXANDRIA MONOGYNIA, Linn.—JUNCI, Juss.

This species is distinguished by its erect, succulent stem; elongated, lanceolate, smooth leaves; and umbellate, subsessile flowers, which are of a deep purple colour, with yellow anthers.

THE AMERICAN REDSTART.

†MUSCICAPA RUTICILLA, Linn.

PLATE LXVIII.—MALE AND FEMALE.

This is one of the most lively, as well as one of the handsomest, of our Flycatchers, and ornaments our woods during spring and summer, when it cannot fail to attract the attention of any person who may visit the interior of the shady forests. It is to be met with over the whole of the United States, where it arrives, according to the different localities, between the beginning of March and the 1st of May. It takes its departure, on its way southward, late in September, and in the beginning of October.

It keeps in perpetual motion, hunting along the branches sidewise, jumping to either side in search of insects and larvæ, opening its beautiful tail at every movement which it makes, then closing it, and flirting it from side to side, just allowing the transparent beauty of the feathers to be seen for a moment. The wings are observed gently drooping during these motions, and its pleasing notes, which resemble the sounds of *Tetee-weet*, *Tetee-weet*, are then emitted. Should it observe an insect on the wing, it immediately flies in pursuit of it, either mounts into the air in its wake, or comes towards the ground spirally and in many zig-zags. The insect secured, the lovely Redstart reascends, perches, and sings a different note, equally clear, and

Small-headed Flycatcher
Virginian Spider-wort. Tradescantia virginica?
 Male

Audubon, John James. 1840. "Small-Headed Flycatcher, *Muscicapa minuta*, Wilson. [Pl. 67]." *The birds of America : from drawings made in the United States and their territories* 1, 238–240. <https://doi.org/10.5962/p.319180>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124833>

DOI: <https://doi.org/10.5962/p.319180>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319180>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.