web; a narrow white ring surrounding the eye; two bands of dull white on the wing; sides of the head and neck greenish-yellow, the rest of the lower parts pale yellow, gradually fading into white behind.

Male, 5, $8\frac{1}{4}$.

Kentucky, Pennsylvania, and New Jersey. Exceedingly rare. Migratory.

THE VIRGINIAN SPIDER-WORT.

TRADESCANTIA VIRGINICA, Willd. Sp. Pl., vol. ii. p. 16. Pursh, Fl. Amer., vol. i. p. 218.
—HEXANDRIA MONOGYNIA, Linn.—JUNCI, Juss.

This species is distinguished by its erect, succulent stem; elongated, lanceolate, smooth leaves; and umbellate, subsessile flowers, which are of a deep purple colour, with yellow anthers.

THE AMERICAN REDSTART.

†MUSCICAPA RUTICILLA, Linn.

PLATE LXVIII.—MALE AND FEMALE.

This is one of the most lively, as well as one of the handsomest, of our Flycatchers, and ornaments our woods during spring and summer, when it cannot fail to attract the attention of any person who may visit the interior of the shady forests. It is to be met with over the whole of the United States, where it arrives, according to the different localities, between the beginning of March and the 1st of May. It takes its departure, on its way southward, late in September, and in the beginning of October.

It keeps in perpetual motion, hunting along the branches sidewise, jumping to either side in search of insects and larvæ, opening its beautiful tail at every movement which it makes, then closing it, and flirting it from side to side, just allowing the transparent beauty of the feathers to be seen for a moment. The wings are observed gently drooping during these motions, and its pleasing notes, which resemble the sounds of *Tetee-weet*, *Tetee-weet*, are then emitted. Should it observe an insect on the wing, it immediately flies in pursuit of it, either mounts into the air in its wake, or comes towards the ground spirally and in many zig-zags. The insect secured, the lovely Redstart reascends, perches, and sings a different note, equally clear, and

American Redstart
Virginian Hornbeam or Ironwood Tree.
 1. Male 2 Female

which may be expressed by the syllables *wizz, wizz, wizz*. While following insects on the wing, it keeps its bill constantly open, snapping as if it procured several of them on the same excursion. It is frequently observed balancing itself in the air, opposite the extremity of a bunch of leaves, and darting into the midst of them after the insects there concealed.

When one approaches the nest of this species, the male exhibits the greatest anxiety respecting its safety, passes and repasses, fluttering and snapping its bill within a few feet, as if determined to repel the intruder. They now and then alight on the ground, to secure an insect, but this only for a moment. They are more frequently seen climbing along the trunks and large branches of trees for an instant, and then shifting to a branch, being, as I have said, in perpetual motion. It is also fond of giving chase to various birds, snapping at them without any effect, as if solely for the purpose of keeping up the natural liveliness of its disposition.

The young males of this species do not possess the brilliancy and richness of plumage which the old birds display, until the second year, the first being spent in the garb worn by the females; but, towards the second autumn, appear mottled with pure black and vermilion on their sides. Notwithstanding their want of full plumage, they breed and sing the first spring like the old males.

I have looked for several minutes at a time on the ineffectual attacks which this bird makes on wasps while busily occupied about their own nests. The bird approaches and snaps at them, but in vain; for the wasp elevating its abdomen, protrudes its sting, which prevents its being seized. The male bird is represented in the plate in this posture.

Its nest is generally made on a low bush or sapling, and has the appearance of hanging to the twigs. It is slight, and is composed of lichens and dried fibres of rank weeds or grape vines, nicely lined with soft cottony materials. The female lays from four to six white eggs, sprinkled with ash-grey and blackish dots. It rears only a single brood in a season. The old birds, I am inclined to think, leave the United States a month or three weeks before the young, some of which linger in the deep swamps of the States of Mississippi and Louisiana until the beginning of November.

This bird differs in no essential respect from the Flycatchers above mentioned. Its mouth has the same structure, being only a little more concave in front. The tongue is of the same form, but proportionally narrower, its tip slit. The œsophagus is 1 inch 8 twelfths long, its average width 1 twelfth. The stomach $4\frac{1}{2}$ twelfths by $3\frac{3}{4}$ twelfths. Intestine 3 inches 10 twelfths long, its greatest width barely 1 twelfth; cœca little more than $\frac{1}{2}$ twelfth long, and $7\frac{1}{2}$ twelfths distant from the extremity. Trachea $1\frac{1}{4}$ inch long, of 55 rings,

with 2 dimidiate; its muscles as in the other species, but the inferior laryngeal proportionally a little larger; bronchi of about 12 half rings.

AMERICAN REDSTART, *Muscicapa ruticilla*, Wils. Amer. Orn., vol. i. p. 103.

MUSCICAPA RUTICILLA, Bonap. Syn., p. 68.

AMERICAN REDSTART, *Muscicapa ruticilla*, Aud. Amer. Orn., vol. i. p. 202; vol. v. p. 428.

AMERICAN REDSTART, *Muscicapa ruticilla*, Nutt. Man., vol. i. p. 291.

Second and third quills equal and longest, fourth longer than first; tail rounded. Male with the head, neck all round, fore part of breast, and back, glossy bluish-black; sides of the breast, lower wing-coverts, a patch on the wings formed by the margins of the primaries and the basal half of most of the secondaries, together with three-fourths of both webs of the outer four tail-feathers on each side, and the outer web of the next, bright orange-red; abdomen and lower tail-coverts white. Female with the upper parts yellowish-brown; the head grey; the quills greyish-brown; the tail darker; the parts yellow which in the male are bright orange; the rest of the lower parts white, tinged with yellow. Young similar to the female, more grey above, and with less yellow beneath.

Male, 5, 6½. Female.

Throughout the United States. Abundant. Migratory.

THE VIRGINIAN HORNBEAM, OR IRON-WOOD TREE.

OSTRYA VIRGINICA, Willd., Sp. Pl., vol. iv. p. 469. Pursh, Flor. Amer., vol. ii. p. 623.—

MONÆCIA POLYANDRIA, Linn.—AMENTACEÆ, Juss.

This species is distinguished by its ovato-oblong leaves, which are somewhat cordate at the base, unequally serrated and acuminate, and its twin, ovate, acute cones. It is a small tree, attaining a height of from twenty to thirty feet, and a diameter of about one foot. The wood is white, and close grained. The common name in America is *iron-wood*, which it receives on account of the great hardness of the wood.

American Redstart
Virginian Hornbeam or Ironwood Tree.
 1. Male 2 Female

Audubon, John James. 1840. "The American Redstart, *Muscicapa ruticilla*, Linn. [Pl. 68]." *The birds of America : from drawings made in the United States and their territories* 1, 240–242. <https://doi.org/10.5962/p.319181>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124833>

DOI: <https://doi.org/10.5962/p.319181>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319181>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.