narrow feathers. Bill brownish-black. Iris hazel. Feet and claws greyish-blue. Head all round, neck and under parts generally, of a bright rich pure yellow, paler on the abdomen, and passing into white on the under tail-coverts. Fore part of the back and lesser wing-coverts yellowish-green. Lower back and wings light greyish-blue. Inner webs of the quills blackish. Inner webs of the tail-feathers bluish-grey at the base, then white to near the tip, which is black, as well as the outer webs. The two middle feathers blackish, tinged with greyish-blue.

Length $5\frac{1}{2}$ inches, extent of wings $8\frac{1}{2}$; beak along the ridge $\frac{7}{12}$, along the gap $\frac{3}{4}$; tarsus $1\frac{1}{2}$.

Adult Female.

The differences which the female exhibits are so slight as scarcely to be describable, the tints being merely a little duller.

GOLDEN-WINGED SWAMP-WARBLER.

+*HELINAIA CHRYSOPTERA*, Linn.

PLATE CVII.—MALE AND FEMALE.

Although I have met with this species entering the United States from the Texas in the month of April, and have procured several specimens in Kentucky and Louisiana, as well as a single one in New Jersey, I never had the good fortune to find its nest. When it first makes its appearance in Louisiana or Kentucky, it usually resorts to the higher branches of trees, where, amid the opening leaflets and blossoms, it actively searches for its insect food, occasionally following its prey on wing to some distance, and moving by short leaps among the twigs, in the manner of *Helinaia carbonata*, which, in its elongated and slender shape, it in some measure resembles. The flight of this species is, unlike that of the Cape May Warbler, *Sylvicola maritima*, elevated, swift, and irregularly undulated, until it is about to alight, when it dives toward the spot selected by it, as most Warblers are wont to do. I never saw a bird of this species in autumn, and therefore infer that its southward journey must be accomplished in a very secret and careful manner, or by night. A male and a female are figured in their perfect spring plumage.

GOLDEN-WINGED WARBLER, *Sylvia crysoptera*, Wils. Amer. Orn., vol. ii. p. 113.

SYLVIA CHRYSOPTERA, Bonap. Syn., p. 87.

GOLDEN-WINGED WARBLER, *Sylvia chrysoptera*, Nutt. Man., vol. i. p. 411.

GOLDEN-WINGED WARBLER, *Sylvia chrysoptera*, Aud. Orn. Biog., vol. v. p. 154.

Male in spring.

Bill shorter than the head, slender, conical, compressed toward the end, tapering to an acute point; upper mandible with the dorsal line almost perfectly straight, being very slightly convex toward the end, the ridge narrow, the sides sloping at the base, rounded toward the end, the edges a little inflected, without notch, the tip acuminate; nostrils basal, oblong, operculate; lower mandible with the angle rather short and obtuse, the dorsal line straight, the sides convex, the edges inflected, the tip acuminate, the gape-line nearly straight.

Head of moderate size, ovate; neck rather short; body slender. Feet of moderate length, slender; tarsus longer than the middle toe, much compressed, covered anteriorly by seven scutella, posteriorly by two plates meeting so as to form a very thin edge; toes small, much compressed; hind toe comparatively large, lateral toes nearly equal, middle toe much longer; claws moderate, well curved, much compressed, laterally grooved, acute.

Plumage very soft and blended. Wings of moderate length, the second quill longest, the third scarcely shorter, the first and fourth about equal, the first with the outer web narrowed in its whole length, the next three toward the end; secondaries long, rather narrow, rounded. Tail rather long, nearly even, the middle feathers being scarcely a twelfth of an inch shorter than the lateral.

Bill and feet black; iris brown. The general colour of the upper parts is light ash-grey, of the lower white; the upper part of the head, and a patch on the wing, formed by the first row of small coverts and the secondary coverts, bright yellow; a band from the bill to the eye, continued under it, and enlarging behind, so as to include the ear-coverts, together with the throat, to the extent of about an inch, black; a white band from over the eyes to behind the ears, and another from the lower mandible down the side of the neck, enlarging as it proceeds; the sides under the wings very pale ash-grey. The quills and tail-feathers are brown, edged with ash-grey; the outer three feathers of the tail have a large portion of the inner web white.

Length to end of tail $4\frac{1}{4}$ inches, to end of wings $3\frac{7}{8}$; extent of wings $7\frac{5}{8}$; bill along the ridge $\frac{5\frac{3}{4}}{12}$, along the edge of lower mandible $\frac{7\frac{1}{2}}{12}$; wing from flexure $2\frac{8}{12}$; tail $2\frac{1}{4}$; tarsus $\frac{8\frac{1}{2}}{12}$; hind toe $\frac{3}{12}$, its claw $\frac{3}{12}$; middle toe, $\frac{5}{12}$ its claw $\frac{2}{12}$.

Adult Female.

The female resembles the male, but has the tints less bright, the sides of

the head and the throat grey instead of black, and the white bands on the head narrower and less extended.

Length to end of tail 4 inches, to end of wings $3\frac{5}{8}$; extent of wings $7\frac{1}{4}$.

BACHMAN'S SWAMP-WARBLER.

+ *HELINAIA BACHMANII*, Aud.

PLATE CVIII.—MALE AND FEMALE.

My friend BACHMAN has the merit of having discovered this pretty little species of Warbler, and to him I have the pleasure of acknowledging my obligations for the pair which you will find represented in the plate, accompanied with a figure of one of the most beautiful of our southern flowers, originally drawn by my friend's sister, Miss MARTIN. I myself have never had the good fortune to meet with any individuals of this interesting *Sylvia*, respecting which little is as yet known, its discoverer having only procured a few specimens of both sexes, without being able to find a nest. The first obtained was found by him a few miles from Charleston, in South Carolina, in July 1833, while I was rambling over the crags of Labrador. According to my amiable friend, it was "a lively active bird, gliding among the branches of thick bushes, occasionally mounting on the wing and seizing insects in the air in the manner of a Flycatcher. It was an old female that had to all appearance just reared a brood of young." Shortly after, several were seen in the same neighbourhood; and we may still expect an account of its manners, migration, and breeding, although not yet discovered.

BACHMAN'S WARBLER, *Sylvia Bachmanii*, Aud. Orn. Biog., vol. ii. p. 483.

Adult Male.

Bill rather long, slightly bent towards the tip, subulato-conical, extremely acute, the edges sharp and inflected. Nostrils basal, lateral, elliptical, half-closed above by an arched membrane. The general form slender. Feet of ordinary length; tarsus slender, compressed, anteriorly scutellate, sharp behind; toes free, the hind toe strong, the two lateral nearly equal, the middle one much longer; claws slender, arched, compressed, acute.

R.T.

Golden-winged Swamp Warbler.

1 Male. 2 Female.

Audubon, John James. 1841. "Golden-Winged Swamp-warbler, *Helinaia chrysophtera*, Linn. [Pl. 107]." *The birds of America : from drawings made in the United States and their territories* 2, 91–93. <https://doi.org/10.5962/p.319220>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124834>

DOI: <https://doi.org/10.5962/p.319220>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319220>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.