plates meeting so as to form a very sharp edge; toes scutellate above, free, the lateral nearly equal. Claws slender, arched, much compressed, acute.

Plumage soft and blended, the wing-coverts compact. Wings of moderate length, the third primary longest, the fourth scarcely shorter, the second less than one-twelfth shorter than the third, and exceeding the first by one-twelfth. Tail long, broad, emarginate and divaricate, the middle feathers only a twelfth and a half shorter than the longest, which is the third from the lateral, the latter being of the same length as the middle.

Bill brown above, paler beneath. Iris brown. Feet and claws light brown. Forehead and a band over the eye, proceeding down the neck, crimson; throat, fore part of breast and sides, with the rump, rich carmine, the latter paler. The upper parts greyish-brown, the head, hind neck, and fore part of the back slightly tinged with red; quills, coverts, and tail-feathers dusky and edged with pale brownish-grey, the first row of small coverts rather conspicuously tipped with a lighter tint of the same. Short bristly feathers at the base of the bill greyish-yellow; loreal space and ear-coverts, light grey; the hind part of the breast, abdomen, and lower tail-coverts yellowish-white, streaked with dusky.

Length to end of tail $6\frac{1}{4}$ inches; bill along the ridge $\frac{4\frac{1}{2}}{12}$, along the edge of lower mandible $\frac{5\frac{1}{2}}{12}$; wing from flexure $3\frac{1}{2}$; tail $3\frac{1}{4}$; tarsus $\frac{8}{12}$; hind toe $\frac{3\frac{1}{2}}{12}$, its claw $\frac{2\frac{1}{2}}{12}$; middle toe $\frac{6\frac{1}{2}}{12}$, its claw $\frac{2}{12}$.

GREY-CROWNED PURPLE-FINCH.

⁴ERYTHROSPIZA TEPHROCOTIS, *Swains.*

PLATE CXCVIII.—MALE.

This species, which in form and proportion is allied to *Erythrospiza purpurea* on the one hand, and to *E. cannabina* on the other, is of extremely rare occurrence, a single specimen only being mentioned as having been obtained by Dr. RICHARDSON on the Saskatchewan, in May 1827, from which my figure was taken.

Male, 6; wing, 4.

Saskatchewan river. Very rare. Migratory.

Grey-crowned Purple Finch.

Male.

Scoptes cyanea.

LINARIA (LEUCOSTICTE) TEPHROCOTIS, *Swainson's Grey-crowned Linnet*, Swains. and Rich. F. Bor. Amer., vol. ii. p. 265.

GREY-CROWNED LINNET, *Fringilla tephrocotis*, Aud. Orn. Biog., vol. v. p. 232.

Adult Male.

Bill short, thick, conical, its upper and lower outlines straight, the ridge and sides convex, the edges inflected, the gap-line slightly arched, but at the base a little deflected. Nostrils basal, round, concealed by short bristly feathers.

Head rather large, broadly ovate; neck short; body moderate. Feet of moderate length; tarsus of the same length as the middle toe and claw, compressed, anteriorly scutellate, with two lateral plates meeting behind so as to form a sharp edge; hind toe stout, lateral toes nearly equal. Claws rather long, arched, compressed, acute, that of the hind toe largest.

Plumage soft and blended. Wings long, pointed; the second quill longest, the first slightly shorter, the rest rapidly graduated; secondaries rounded. Tail of moderate length, emarginate, the middle-feathers a quarter of an inch shorter than the longest.

Bill black; tarsi and claws dusky brown. The general colour of the plumage is dark umber-brown; the feathers margining the bill whitish; the upper part of the head ash-grey, spotted with black anteriorly; the first row of smaller wing-coverts, the feathers of the rump and the upper tail-coverts are broadly edged and tipped with rose-red, as are the feathers of the sides and the lower tail-coverts, of which however the tint is paler. The quills, larger coverts, and tail-feathers, are dusky brown; the primary and secondary coverts edged with dull red, the quills and tail-feathers with brownish-white.

Length to end of tail 6 inches; bill along the ridge $\frac{5\frac{1}{2}}{12}$, along the edge of lower mandible $\frac{6\frac{1}{2}}{12}$; wing from flexure 4; tail $2\frac{7}{12}$; tarsus $\frac{8\frac{1}{2}}{12}$; hind toe $\frac{4}{12}$, its claw $\frac{3}{12}$; middle toe $\frac{6}{12}$, its claw $\frac{2\frac{1}{2}}{12}$.

Grey-crowned Purple Finch.

Male.

Scoptes cyanea

Audubon, John James. 1841. "Grey-Crowned Purple-Finch, *Erythropsiza tephrocotis*, Swains. [Pl. 198]." *The birds of America : from drawings made in the United States and their territories* 3, 176–177. <https://doi.org/10.5962/p.319311>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124835>

DOI: <https://doi.org/10.5962/p.319311>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319311>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.