TOWNSEND'S SURF-BIRD.

† *APHRIZA TOWNSENDI*, Aud.

PLATE CCCXXII.—FEMALE.

The remarkable bird here represented, which in form and size bears a considerable resemblance to the Knot, was procured by Mr. TOWNSEND on the shores of Cape Disappointment, and proved to be a female. Nothing is known as to the habits or range of the species. In order to exhibit its characters to the best advantage, I have figured it flying in two different aspects. The following note accompanied the specimen sent to me by Mr. TOWNSEND:—"I shot this bird, the only one I have ever seen, on Cape Disappointment, at the entrance of the Columbia river. It was sitting on the edge of the steep rocks, and the heavy surf frequently dashed its spray over it as it foraged among the retreating waves. When it started, it flew with a quick, jerking motion of its wings, and alighted again at a short distance. It was a female. The stomach was remarkably strong and muscular, and contained fragments of a small black shell-fish which adheres to the rocks in this neighbourhood."

TOWNSEND'S SURF-BIRD, *Aphriza Townsendi*, Aud. Orn. Biog., vol. v. p. 249.

Female, 11; wing, $7\frac{1}{2}$.

Cape Disappointment, Columbia river.

Female.

Bill a little shorter than the head, rather stout, compressed, tapering, straightish, being recurvate in a slight degree. Upper mandible with the dorsal line straight and a little declinate as far as the middle, then concave, and towards the end convex, the nasal groove extending to near the end, the ridge rather broad and flattened, the tip compressed and bluntish. Lower mandible with the angle rather long and narrow, the dorsal line ascending and slightly convex, the sides grooved for half their length, convex toward the end, the tip narrowed but blunt. Nostrils sub-basal, linear near the margin.

Head rather small, ovate, rounded in front. Neck of ordinary length. Body rather full. Feet of moderate length, rather stout; tibia bare at the lower part, and reticulated; tarsus roundish, with small angular scales all round, those on the fore part larger; toes four, with numerous scutella, the

Drawn from Nature by J. T. Audubon & J. B. S. Ellis

Terns and Jays - Bird.
Females.

Printed & Colored by J. T. Bowen, Phila.

first very small and placed higher than the rest; the anterior toes free to the base, distinctly margined on both edges, flat beneath, the inner considerably shorter than the outer, the third a quarter of an inch longer than the latter; claws rather small, curved, compressed, blunted.

Plumage full, soft, rather dense, on the neck and lower parts blended. Wings very long, narrow and pointed; primaries with strong shafts, narrowed towards the end, the first longest, the rest rapidly decreasing; outer secondaries incurved, obliquely rounded, inner elongated, one of them reaching to an inch and two-thirds of the tip of the longest primary when the wing is closed. Tail rather short, even, of twelve moderately broad, rounded feathers.

Bill dusky toward the end, orange at the base. Feet bluish-green, claws black. The general colour of the upper parts is a very dark or blackish-grey; the quills greyish-black; a broad band of white crosses the wing, occupying the tips of the primary coverts, the terminal third of the secondary coverts, the bases and more or less of the margins and tips of the quills, several of the inner secondaries having only a streak of dusky on the inner web, but the innermost or elongated quills are destitute of white. The shafts of the quills are also white, as are some of the feathers of the rump, the upper tail-coverts, the basal half of the tail, of which the rest is black, the feathers narrowly edged with white at the end; the black on the tail is narrower on the lateral feathers, and on the outer does not occupy much more than half an inch. The throat is greyish-white; the cheeks, sides, and fore part of the neck, and the anterior part of the breast dull grey, of a lighter tint than the back; the rest of the lower parts white, with small longitudinal oblong dark grey streaks; the axillaries and lower wing-coverts white, those at the edge of the wing dark grey, with white margins.

Length to end of tail 11 inches; bill from flexure $1\frac{11}{12}$, along the edge of lower mandible $1\frac{1}{12}$; wing from flexure $7\frac{1}{2}$; tail $3\frac{2}{12}$; tarsus $1\frac{2}{12}$; hind toe $\frac{2}{12}$, its claw $\frac{2}{12}$; middle toe $1\frac{1}{12}$, its claw $\frac{3}{12}$.

The prominence on the terminal part of the upper mandible gives the bill somewhat of the appearance of that of a Plover, but in other respects it more resembles that of the Turnstone, the plumage agrees with that of the latter bird, and the colouring is very similar to its winter dress. This species in short seems intermediate between *Tringa* and *Strepsilas*, but is much more allied to the latter, with which it agrees in form and proportions, the principal differences being in the tail, which is not rounded, but even, in the want of scutella on the tarsi, and in the form of the bill at its extremity, the upper mandible in place of being a little recurvate and depressed, having its extremity arched and the point a little decurved. Were the latter worn off, it would agree with that of *Strepsilas*. Conceiving this bird to present cha-

Drawn from Nature by J. T. Audubon & J. B. S. Ellis

Terns and Jays - Bird.
Females.

Printed & Colored by J. T. Bowen, Phila.

Audubon, John James. 1842. "Townsend's Surf-Bird, *Aphriza townsendi*, Aud. [Pl. 322]." *The birds of America : from drawings made in the United States and their territories* 5, 228–230. <https://doi.org/10.5962/p.319435>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124982>

DOI: <https://doi.org/10.5962/p.319435>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319435>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.