H A V E L L ' S T E R N .

+STERNA HAVELLII, Aud.

PLATE CCCCXXXIV.—ADULT.

I have several reasons for naming this Tern after Mr. ROBERT HAVELL, of Oxford Street, London. In the first place I consider him as one of the best ornithological engravers in England. Secondly, I feel greatly indebted to him for the interest which he has always evinced in my publication, which, I dare venture to assert, is the largest work of the kind that has hitherto appeared, and the engraving of which has cost him much trouble and anxiety. Thirdly, I consider myself entitled to express my gratitude in this manner, the individual on whom I confer the honour being more deserving of it than many to whom similar compliments have been paid.

I shot several individuals of this species out of a number congregated on the broad eddies opposite New Orleans, in 1820. They were engaged in picking up floating coleopterous insects; but after I had fired several shots, and was rowing to those which had fallen on the water, the rest flew off. Since that time it had not been my fortune to meet with any birds of the same species, until I visited Texas in the spring of 1837, when two of them were procured.

HAVELL'S TERN, *Sterna Havelli*, Aud. Orn. Biog., vol. v. p. 122.

Adult, $15\frac{1}{2}$, wing, $10\frac{3}{12}$.

From Texas to South Carolina. Common. Migratory.

Adult in winter plumage.

Bill about the length of the head, rather stout, much compressed, acute. Upper mandible with the dorsal line slightly declinato-convex, the ridge convex at the base, very narrow in the rest of its extent, the side sloping at the base, nearly erect and convex toward the end, the edges sharp and inflected, the tip very narrow. Nasal groove rather short, but with a channel surmounted by a ridge running from its anterior part to the edge of the mandible about half an inch from the tip. Lower mandible with the angle very narrow and acute, extending to beyond the middle, the outline of the crura a little concave, that of the rest ascending and straight, a slight pro-

minence or angle being formed at their junction, as in Gulls, the sides erect and slightly convex, the edges sharp and inclinate, the tip acuminate, the gap-line slightly arcuate.

Head of moderate size, ovate; neck of moderate length; body slender. Feet small; tibia bare for seven-twelfths of an inch; tarsus very short, compressed, anteriorly scutellate; toes small, slender; the first extremely small, the third longest, the fourth much longer than the second; all scutellate above, the anterior connected by reticulate webs, of which the inner is more deeply emarginate. Claws moderately arched, compressed, very slender, that of the middle toe much larger, and having its inner edge somewhat dilated.

Plumage soft, close, blended, very short on the fore part of the head. Wings very long, narrow, and pointed; primary quills tapering to an obtuse point; the first longest, the second half an inch shorter, the rest rapidly graduated; secondaries short, incurved, obliquely pointed, some of the inner proportionally longer and narrower. Tail of moderate length, deeply forked, of twelve feathers, of which the middle are rounded, and three inches and a quarter shorter than the outer, which taper to a slender point.

Bill black, with the base of the lower mandible tinged with brown, and a very small portion of the tip yellowish. Iris brown. Feet orange-yellow; claws dusky. Surrounding the eye, and extending toward the nape, is a broad band of black; the fore part of the head, the lower eyelid, and all the under parts are pure white; the hind part of the head and the nape are dusky-grey, mixed with white. The rest of the upper parts are light greyish-blue, excepting the rump, which is white, the primary coverts and quills as well as the tail-feathers and their coverts are hoary, with the shafts white; but five of the quills are dusky on the outer web, on the inner along the shaft, and on the inner margin toward the end.

Length to end of tail $15\frac{1}{2}$ inches; bill along the ridge $1\frac{7}{12}$, along the edge of lower mandible $2\frac{4}{12}$; wing from flexure $10\frac{8}{12}$; tail to end of middle feather $2\frac{8}{12}$, to end of longest feather $6\frac{1}{12}$; tarsus $\frac{11\frac{1}{2}}{12}$; hind toe $\frac{3}{12}$, its claw $\frac{2}{12}$; middle toe $\frac{10}{12}$, its claw $\frac{4\frac{1}{2}}{12}$.

This species differs from the Marsh Tern, *Sterna anglica*, in being less robust, in having the bill a little longer and much more slender, its height at the angle being $\frac{3\frac{3}{4}}{12}$, whereas in that species it is $\frac{4\frac{3}{4}}{12}$; in having the tarsus shorter and much more slender, the feet yellow instead of being black, the claws more slender, and the tail more deeply forked.

The figure in the plate, which is that of an adult bird yet in its winter plumage, has the lateral tail-feathers obliquely truncate, but this was caused by accident, for these feathers in my other specimens run to a narrow point. My specimens from Texas are also in their winter plumage. One of them

is coloured as above; but the other is a young bird, which may here be described.

The bill is somewhat shorter and more tinged with brown; the lower parts, the rump, the outer web of the lateral tail-feathers and the sides of the neck are white; the wings as in the adult, but the primaries internally margined with white, and the secondaries tipped with the same; the upper part of the head, and the rest of the upper parts, are light yellowish-brown, intermixed with greyish-blue, and there is the same black band on the side of the head as in the adult.

TRUDEAU'S TERN.

†*STERNA TRUDEAUI*, *Aud.*

PLATE CCCCXXXV.—ADULT.

This beautiful Tern, which has not hitherto been described, was procured at Great Egg Harbour in New Jersey, by my much esteemed and talented friend, J. TRUDEAU, Esq. of Louisiana, to whom I have great pleasure in dedicating it. Nothing is known as to its range, or even the particular habits in which it may differ from other species. The individual obtained was in the company of a few others of the same kind. I have received from Mr. TRUDEAU an intimation of the occurrence of several individuals on Long Island.

TRUDEAU'S TERN, *Sterna Trudeaui*, Aud. Orn. Biog., vol. v. p. 125.

Adult, 16; wing, $10\frac{1}{2}$.

Great Egg Harbour and Long Island. Rare. Migratory.

Adult in winter plumage.

Bill about the length of the head, rather slender, much compressed, acuminate. Upper mandible with the dorsal line slightly declinato-convex, the ridge rather broad and convex at the base, very narrow in the rest of its extent, the sides sloping at the base, nearly erect and convex toward the end, the edges sharp and inflected, the tip very narrow. Nasal groove rather short, but with a slight channel, surmounted by a ridge, running from its

Great Frigatebird

Adult

Drawn from Nature by J. Audubon F.R.S.F.L.S.

Lith. Printed & Col^d by J. T. Bowen, Philad^a

Audubon, John James. 1844. "Havell's Tern, *Sterna havellii*, Aud. [Pl. 434]." *The birds of America : from drawings made in the United States and their territories* 7, 103–105. <https://doi.org/10.5962/p.319546>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124981>

DOI: <https://doi.org/10.5962/p.319546>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319546>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.