first five quills are black at the end, the first on its outer web also, the fifth with a small white tip, the tips of all the other quills more or less white.

Length to end of tail 18 inches, to end of wings 20, to end of claws 17; extent of wings $36\frac{1}{2}$; wing from flexure 12; tail 7; bill along the back $1\frac{1}{2}$, along the edge of lower mandible $2\frac{2}{12}$; tarsus $1\frac{7}{12}$; middle toe $1\frac{1}{2}$, its claw $\frac{4}{12}$. Weight $1\frac{1}{2}$ lbs.

Young bird in January.

Bill and feet black. Edges of eyelids and iris as in the adult. The hind head and neck are bluish-grey, and before the eye there is a semi-lunar blackish mark, the tips of the auriculars also dark grey. Forehead, sides of the head, throat, and lower parts, white, as is the rump. Tail white, with a broad terminal band of black, the outer feather having only a spot on the inner web. The mantle is bluish-grey, but a broad band of black crosses the lower part of the hind neck, and the larger wing-coverts are of the same colour towards the end. The primary quills are black, more or less margined with white internally.

Length to end of tail 17 inches, to end of wings 19, to end of claws 17; extent of wings $36\frac{3}{12}$. Weight $14\frac{1}{2}$ oz.

THE IVORY GULL.

†*LARUS EBURNEUS*, *Gmel.*

PLATE CCCCXLV.—ADULT AND YOUNG.

Having ascertained that this beautiful species visits the southern coast of Labrador and Newfoundland every winter, I have thought it probable that it occasionally extends its rambles as far as our eastern shores, and therefore determined to include it in my Illustrations. The figures in the plate were taken from two specimens procured by Captain JAMES CLARK ROSS, one of which was an adult male, the other a young bird in its second year. Captain SABINE says that the Ivory Gulls are attracted in considerable numbers by whale blubber, are therefore usually found in company with the *Procellaria glacialis*, and are easily killed, being by no means shy. Dr. RICHARDSON informs us that they were observed breeding in great numbers on the high perforated cliffs which form the extremity of Cape Parry, in latitude 70°.

Drawn from Nature by W. J. Hudson, F.R.S. P.L.S.

Great Grebe.
1. Adult Male 2. Young second plumage.

1848. Printed & sold by J. T. Bowen, Philadelphia.

LARUS EBURNEUS, Bonap. Syn., p. 360.

LARUS EBURNEUS, *Ivory Gull*, Swains. and Rich. F. Bor. Amer.

IVORY GULL, Nutt. Man., vol. ii. p. 301.

IVORY GULL, *Larus eburneus*, Aud. Orn. Biog., vol. iii. p. 571.

Adult, 19, 41.

Accidental on the coast of the United States. Common in winter in Labrador and Newfoundland. Breeds in high latitudes.

Adult Male.

Bill shorter than the head, robust, nearly straight, compressed. Upper mandible with the dorsal line nearly straight at the base, arched and declinate towards the end, the ridge convex, the sides slightly so, the edges sharp, a little inflected, somewhat arched, the tip rather obtuse. Nasal groove rather long and narrow; nostrils in its fore part, lateral, longitudinal, linear, wider anteriorly, pervious. Lower mandible with a prominence at the end of the angle, which is long and narrow, the dorsal line slightly concave and ascending, the sides flattened, the edges sharp and inflected.

Head rather large. Neck of moderate length, strong. Body rather full. Feet of moderate length, stout; tibia bare below; tarsus somewhat compressed, covered before with numerous scutella, on the sides and behind with series of small angular scales; the hind toe very small and elevated, the fore toes of moderate length, the fourth much longer than the second, the third longest, the hind one with a single scutellum and three transverse series of scales, the rest scutellate above and connected by reticulate membranes having a concave margin, the lateral toes margined externally with a narrow membrane. Claws stout, rather large, arched, compressed, rather obtuse, that of middle toe with an enlarged inner edge.

The plumage in general is close, full, elastic, soft and blended, on the back rather compact. Wings very long, rather broad, acute, the first quill longest, the other primaries rapidly graduated; secondaries broad and rounded, the inner tapering but rounded. Tail of moderate length, even, of twelve rather broad rounded feathers.

Bill bright yellow, greenish-dusky at the base. Iris brown, edges of eyelids vermilion. Feet and claws black. The whole of the plumage is pure white.

Length to end of tail 19 inches, to end of wings $20\frac{1}{2}$; extent of wings 41; wing from flexure $13\frac{1}{2}$; tail $6\frac{1}{4}$; bill along the back $1\frac{5}{12}$, along the edge of lower mandible 2; tarsus $1\frac{7}{12}$; middle toe $1\frac{2}{12}$, its claw $\frac{5}{12}$. Weight 20 oz.

Young of the second year, killed in September.

After the second moult, the bill is pale yellow at the end, dusky at the base for two-thirds of its length; the edges of the eyelids vermilion, the iris brown, the feet black. The plumage is white; the forehead and sides of the

head mottled with leaden-grey; most of the wing-coverts have towards the end a spot of greyish-black, and the quills, large coverts, and tail-feathers are similarly marked, the markings on the tail forming a subterminal bar.

THE COMMON AMERICAN GULL.

†*LARUS ZONORHYNCHUS*, *Richardson*.

PLATE CCCCXLVI.—ADULT MALE, AND YOUNG IN WINTER.

No country can afford greater facilities for the migration of water-birds than the United States of America. Even the Gulls are enabled to traverse their whole extent from north to south, and in the contrary direction, without suffering from want of food or of proper resting places. The Gull that has been bred in Labrador, or still farther north, can reach the Gulf of Mexico without being seriously incommoded by the storms that now and then rage along the Atlantic coast. The broad waters of the St. Lawrence leads it to our great lakes, from which hundreds of streams conduct it to the head waters of the Ohio or the Mississippi, by following the windings of which it at length arrives in the warm regions of the Mexican Gulf, on whose waters the traveller can spend the winter. That these advantages are embraced by many species of Gull, there can be no doubt; and should you, as I have done, repeatedly visit our broad lakes, or the great rivers just mentioned, you would find there, at particular seasons, not only this species, but several others, as well as various kinds of Terns, but none of the genus *Lestris*. Lake Erie supplies with food the *Larus marinus*, *L. argentatus*, *L. atricilla*, and some others, as well as the Great, the Arctic, the Roseate, and the Black Terns, all of which pass at times over to the Ohio, and from thence to the ocean. During these inland movements, the birds seem to be peculiarly attracted by certain places, at which they remain for awhile. Thus, at the Falls of the Ohio, some species remain for weeks, and wherever much shipping occurs on that river or the Mississippi, Gulls are sure to be seen gleaning the garbage that has been thrown overboard, or seizing such fishes as rise incautiously to the surface of the water. In the months of September and October, Gulls and Terns might almost be said to abound on

W.E.H.

Great Grebe.

1. Adult Male 2. Young second year male.

Audubon, John James. 1844. "The Ivory Gull, *Larus eburneus*, Gmel. [Pl. 445]." *The birds of America : from drawings made in the United States and their territories* 7, 150–152. <https://doi.org/10.5962/p.319557>.

View This Item Online: <https://www.biodiversitylibrary.org/item/124981>

DOI: <https://doi.org/10.5962/p.319557>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319557>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.