The Morphology of the Reproductive Structures in the Podocarpaceae.¹

BY

EDMUND W. SINNOTT.

With Plates V-IX and nine Diagrams in the Text.

THE construction of a natural classification for the Coniferales, based upon comparative studies among its many living and fossil forms, has been the object of all recent morphological investigations which have dealt with members of the group. A wealth of new facts concerning this great plant order has consequently been added to our knowledge, and although in most of its families at least the main outlines of the anatomy of the sporophyte and of the structure and development of the gametophytes are now understood, yet among certain groups so little has been done that, with regard to almost everything but external features, comparative ignorance still prevails.

This is especially true in the case of the Podocarpaceae, an important and clearly defined family, the natural position and relationships of which have been the cause of much discussion and difference of opinion among botanists.

Its members belong predominantly to the Southern Hemisphere, and are represented north of the Equator by only a few species in southern and eastern Asia and the West Indies. The family includes six genera: *Podocarpus*, of sixty species, distributed throughout most of the Southern Hemisphere and in India, China, Japan, and the West Indies; *Dacrydium*, of sixteen species, from Australasia, the East Indies, and southern Asia; *Saxegothea*, monotypic, from the Andes; *Microcachrys*, also monotypic, from Tasmania; *Pherosphaera*, of two species, from Australia; and *Phyllocladus*, of five species, from New Zealand, Tasmania, and the Philippine Islands.

The almost entire restriction of the Podocarpaceae to the Southern Hemisphere doubtless explains the neglect accorded them at the hands of the morphologists, who up to the present have had little but fragmentary and poorly preserved material with which to work. Our knowledge of the

¹ Contributions from the Phanerogamic Laboratories of Harvard University, No. 53.

reproductive features of the family, and especially of the female gametophyte and formation of the embryo, are therefore very incomplete. Observations on the strobilar anatomy of the Podocarpineae date back to 1869, when Van Tieghem (19) published a brief general account of the structure and vascular supply of the female cone. This was amplified by Strasburger (14, 15) in 1872 and 1879, and in more recent years Thomson (17), Tison (18), Stiles (11, 12, 13), Miss Gibbs (4), and Miss Robertson (10) have investigated the sporophyll in several genera. The structure of the mature pollen-grain is known for most of the genera from the work of Jeffrey and Chrysler (5), Burlingame (1), Coker (2), and Miss Young (20), and the development of the male gametophyte has been studied, but in only a comparatively few cases. Our knowledge of the female gametophyte and of the development of the embryo is confined to the observations of Miss Kildahl (6) and Miss Young (21) on *Phyllocladus alpinus*, of Coker (2) on *Podocarpus coriaceus*, and of Stiles (13) and Miss Gibbs (4) on several other species of *Podocarpus*. The last two writers have recently published long papers which contain considerable information as to the reproductive morphology of the family. The results of Mr. Stiles, though accurate and well stated as far as they go, are based on first-hand material of only *Microcachrys*, two of the four sub-genera of *Podocarpus*, and one of the two main divisions of *Dacrydium*. Even in these, few new data are brought forward as to the structure of the female gametophyte and the development of the embryo. Miss Gibbs's paper on the female strobilus of *Podocarpus*, though embracing observations on a large number of forms in all the sub-genera, is mainly concerned with histological details, and lays little emphasis on broad anatomical comparisons between members of the various groups. Considerable information is set forth as to the female gametophyte and embryo, but this is fragmentary, and is concerned almost entirely with the very early stages of the gametophyte and with the structure of the mature embryo. The author draws no conclusions whatever as to relationships.

During the year 1910-11 the present writer was fortunate in being able to visit Australia and New Zealand as Frederick Sheldon Travelling Fellow of Harvard University, and while there collected abundant material for a study of the vegetative and reproductive structures in several of the genera and a large number of the species of the Podocarpineae.¹ Such

¹ These include *Podocarpus Totara*, *P. Hallii*, *P. nivalis*, *P. spinulosus*, *P. elatus*, *P. macrophyllus*, *P. ferrugineus*, *P. spicatus*, *P. vitiensis*, and *P. dacrydioides*; *Dacrydium Bidwillii*, *D. intermedium*, *D. Colensoi*, and *D. cupressinum*; and *Phyllocladus glaucus*, *P. trichomanoides*, and *P. alpinus*.

Anatomical material was preserved in 70 % alcohol or 4 % formalin, but that destined for embryological investigation was first killed in chrom-acetic acid (usually about 1 % chromic acid and 1 % acetic acid in water) or in a mixture of three parts of a saturated solution of corrosive sublimate in 50 % alcohol and one part of a 2 % solution of potassium bichromate; after which it was carefully

of the results as deal with the morphology of the reproductive organs are embodied in the present paper, which has for its object a description of the anatomical features of the staminate and ovulate strobili and a history of the development of the male and female gametophytes and of the embryo, together with a discussion, in the light of the facts set forth, of the relationships between the various genera and of the affinities of the family as a whole to the other great groups of the Coniferales.

THE STAMINATE STROBILUS.

The male cone of the Podocarpaceae exhibits no great diversity of structure and conforms to the general coniferous type. The axis contains a cylinder of small vascular bundles, outside which is a ring of cortical canals. The single bundle departing to each microsporophyll leaves behind it a gap in the cylinder. The sporophylls are always bisporangiate, thus agreeing precisely with those found among the Abietineae, and differing strongly from the peltate multisporengiate sporophylls of the Araucarineae.

THE OVULATE STROBILUS.

The female strobilus of the Podocarpaceae is distinguished from that of all other Conifers save the Taxineae by the exhibition of two well-marked tendencies—towards a reduction in the number of sporophylls, and towards a fleshy condition of a part or the whole of the cone at maturity. In *Podocarpus* and *Dacrydium*, reduction has resulted in most cases in an abbreviation of the axis to a short, thickened stalk, bearing a few sterile, scale-like bracts at its base and one or two ovuliferous appendages above; but in *Saxegothea*, *Microcachrys*, *Pherosphaera*, and *Phyllocladus* the fertile scales are usually more numerous, and are arranged in more typical cones. In all the family, the bract or scale subtends but a single ovule, which is inverted in *Podocarpus* and certain species of *Dacrydium*, but is nearly or quite erect at maturity in the other genera. Outside the ovule, in addition to the usual single integument, there is an 'outer integument', 'arillus' or 'epimatium', which may entirely envelop the seed or be reduced to a small basal sheath. The origin and significance of this epimatium, and the morphological nature of the strobilar appendage on which it is borne, are disputed questions and have an important bearing on our theories as to the relationships of the family.

The genus *Podocarpus*, by far the largest and most widely distributed of the Podocarpaceae, shows much diversity of structure in its various species, and is separated into four main sub-genera on the character of the washed in water or 50 % alcohol, respectively, and run up slowly into 70 % alcohol for preservation. The ordinary precautions used where careful fixation is desired were observed, such as killing the material directly in the field and slicing off the sides of the ovules with a razor, or stout knife in the case of species with a stony integument, to ensure rapid penetration.

female strobilus. *Eupodocarpus*, comprising the great majority of species, possesses a cone reduced to one or two ovules on a short axis which ripens into a bright fleshy 'receptaculum.' In *Stachycarpus* the axis is sometimes very short, bearing a single ovule, and is sometimes elongated, with several, but it is never thickened into a receptaculum. The fleshy character is here present in the epimatium, which becomes pulpy and edible, the seed being protected from injury by an exceedingly hard integument. *Nageia* is somewhat intermediate between these two sub-genera, for its axis is shortened and bears several fleshy bracts. The broad, parallel-veined leaves are the best distinguishing feature. In *Dacrycarpus* the cone is reduced to a single ovule, and the subtending bract, instead of being entirely free from the ovule as in all the rest of the genus, is fused for almost or quite its full length to the dorsal face of the epimatium. The short axis ripens into a fleshy receptaculum. Species of all these sub-genera were investigated.

Five members of *Eupodocarpus* were available for study: *P. Totara*, *P. nivalis*, *P. Hallii*, *P. spinulosus*, and *P. elatus*. The short cone axis in these species bears at its apex a single pair of opposite bracts, and in the axil of one or both is an inverted ovule (Pl. VI, Figs. 7 and 10; Diagram 1, A). A pair of sterile bracts at right angles to the fertile ones are often present, and there are sometimes signs of still a third pair. The slender cone-stalk is provided with a ring of vascular bundles and of cortical canals. This ring expands as it enters the enlarged receptaculum, and on approaching the apex becomes flattened into an ellipse (Diagram 1, B). From each end of the ellipse departs a bundle to one of the two fertile bracts, and each of these bract bundles is closely followed in turn by two large strands which carry away from the sides of the gap all the vascular tissue of the cylinder save the small traces to the sterile scales (Fig. 9; Diagram 1, C). Each pair of large bundles supplies an ovule, and its two component strands, as they pass upward through the cortex, gradually approach each other by their adaxial ends, xylem outward, and form a broadly V-shaped bundle with orientation inverse to that of the bract supply. After the early separation of the bract, this large strand enters the base of the ovule, where its subsequent behaviour varies somewhat according to the species.

In *P. Totara* and *P. nivalis* it divides again into two strands (Diagram 1, D and E), which remain close together and pass upward along the dorsal side of the epimatium (Fig. 8; Diagram 1, F). Upon reaching the chalaza, into which each sends a small concentric bundle (Diagram 1, G), they enter the apical 'knob' and soon die out. One or two weak branches sometimes depart towards the ventral side of the epimatium.

Conditions are similar in *P. Hallii* and *P. spinulosus*, save that the single basal bundle either remains undivided or breaks up into three parallel strands.

DIAGRAM I. *Podocarpus Totara*. A, vertical section through strobilus; B-G, successive transverse sections from receptaculum to chalaza. *P. elatus*. H, transverse section through middle of epimatium and ovule. Integument shaded, xylem black.

In all these species the nucellus for about half its length is free from the integument, which at maturity is divided into a narrow stony layer outside and a wider soft one within. It is fused throughout with the epimatium, which ripens into a leathery coat and is well provided with mucilage canals or sacs.

P. elatus has a much larger ovule and its thick epimatium is supplied by a more complex vascular system. The single basal strand, formed by the fusion of two, divides into three which separate considerably as they ascend. At the chalazal end these strands enter an anastomosing complex of seven main bundles, formed by the three we have just traced, which supply the dorsal portion of the epimatium, and by four others which pass down its ventral side and end blindly near the micropyle (Diagram 1, H). These ventral bundles probably represent continuations of the two lateral

DIAGRAM 2. *Podocarpus spicatus*. A, vertical section through portion of strobilus; B-F, successive transverse sections from axis to chalaza.

members of the dorsal series. They usually fork somewhat and are often connected by small strands. The integument is thin and is fused throughout its entire length to the stout epimatium, but the inner margin of the latter is so fully occupied by distended resin sacs or spaces that the ovule proper is virtually suspended from it by thin strands of tissue (Diagram 1, H). The epimatium ripens into a thick firm coat.

Two members of the sub-genus *Stachycarpus* were investigated—*P. spicatus* and *P. ferrugineus*.

P. spicatus, unlike any other member of the genus, with a very few exceptions, possesses a long loose cone of many ovules somewhat distant

from one another, and each in the axil of a small bract (Fig. 16; Diagram 2, A). The vascular cylinder of the axis is a ring of small bundles, each with its cortical canal (Fig. 19; Diagram 2, B). Immediately after the departure of the strand which supplies the bract two larger ones are given off, one from each side of the gap (which soon closes), and as these pass outward through the cortex they approach each other by their adaxial ends, as in *Eupodocarpus*, and thus display an orientation inverse to that of the bract bundle (Diagram 2, C). The bract immediately becomes free and these two strands, which lie next each other but do not fuse, pass upward into the dorsal side of the epimatium (Diagram 2, D), and on ascending diverge until they lie about 90 degrees apart on the circumference of the ovule. They converge again considerably as they approach the chalazal end, where each sends a small bundle into the base of the ovule (Diagram 2, F) and then enters the apical knob, turns at a sharp angle, and descends into the ventral side of the epimatium without further branching. A cross-section through the median region of the ovule will thus show four bundles in the epimatium, nearly equidistant from one another (Diagram 2, E), the xylem of each facing outward and often tending to surround the phloem entirely. The nucellus is free for nearly half its length from the integument, which is intimately fused, save at its very tip, with the epimatium. The latter is plentifully supplied with large canals and becomes fleshy at maturity, but the integument ripens into an exceedingly hard and stony protective shell.

P. ferrugineus is closely related to *P. spicatus*, and the anatomy of the female strobilus in the two species is similar. The cone axis of *P. ferrugineus*, however, has been reduced to a very short branch bearing a few small scales and terminated by one, or occasionally two, inverted ovules (Figs. 15 and 17; Diagram 3, A). This short axis contains a ring of bundles and of canals, as do the other species, and a small strand is given off to each of the spirally arranged sterile bracts (Diagram 3, B). At the insertion of the ovule, however, one small bundle and its canal enter the fertile bract, and the remainder of the vascular ring divides into two large strands which orient themselves inversely to the bract bundle, and side by side enter the epimatium of the ovule (Pl. VII, Fig. 20; Diagram 3, C and D). In the rare cases where two ovules are present they are each supplied by a pair of bundles. The two strands now pass upward close together to the chalaza, where each sends a small bundle into the base of the ovule (Diagram 3, F) and then turns at an abrupt angle and passes downward, dying out near the micropyle. These two basipetal extensions go down on either side, directly opposite each other, and each lies just outside the lateral ridge of the integument. A median transverse section therefore shows two large dorsal bundles, close together, and two smaller lateral ones (Pl. VI, Fig. 18; Diagram 3, E). The xylem in all faces outward, and in the dorsal bundles, at least, often surrounds the phloem entirely. Many small strands, apparently of phloem,

enter the integument from the lateral bundles, and in some cases completely encircle the nucellus. The lower portion of the nucellus is attached laterally to the integument (Fig. 18; Diagram 3, E), but is otherwise free throughout. The integument, which becomes stony at maturity, is fused with the fleshy epimatium everywhere except along a rather narrow strip on its dorsal and ventral faces.

Only very young ovules of *P. vitiensis*, of the sub-genus *Nageia*, were available for study, but the general arrangement of the vascular tissues could be made out and agrees with that described by Miss Gibbs for the species.

DIAGRAM 3. *Podocarpus ferrugineus*. A, vertical section through strobilus; B-F, successive transverse sections from axis to chalaza.

The reduced cone is similar to that of *P. ferrugineus* and bears a few sterile basal bracts and a single terminal ovule (Fig. 11). At the apex of the cone axis a bundle leaves the cylinder to supply the large fertile bract, in the lamina of which it branches into several parallel veins. The remainder of the vascular ring enters the base of the ovule, where it immediately divides into four bundles. These four—the two median ones close together, the two lateral somewhat divergent—pass up the dorsal face of the epimatium, and at the chalazal end of the ovule enter an anastomosing vascular complex. From this they emerge in the downward direction as two small ventral strands and two larger ones, exactly lateral. Several small bundles enter

the base of the ovule from the chalazal complex. In the general disposition of the vascular supply to its ovule, *P. vitiensis* is intermediate between *P. elatus* and *P. ferrugineus*.

The only other species of *Podocarpus* under investigation, *P. dacrydioides*, is a member of the section *Dacrycarpus*, which is distinguished from the rest of its genus by the fact that the bract is fused to the dorsal face of the ovule. The cone is very much reduced and its short thick axis, becoming fleshy at maturity, bears two opposite bracts, in the axil of one of which and adnate to it is the inverted ovule (Figs. 12, 13, and 14; Diagram 4, A). The vascular cylinder of the axis gives off a strand to each of the two bracts

DIAGRAM 4. *Podocarpus dacrydioides*. A, vertical section through strobilus; B-F, successive transverse sections from receptaculum to chalaza.

(Diagram 4, B and C), and the two large bundles which remain approach each other, xylem outward, and enter the base of the epimatium (Diagram 4, D). At first these two lie near the bract bundle (Diagram 4, E), but they gradually diverge, and on approaching the chalaza separate rapidly, passing across the ovule, one on either side, to its ventral face (Diagram 4, F). Here they again converge, send downward a few weak bundles, and then pass upward a little way before dying out. The bract bundle disappears near the end of the bract, which in the young cone projects some distance beyond the ovule (Figs. 13 and 14) and is readily distinguishable from it, but which at maturity is so intimately fused with the leathery epimatium as to be almost

indiscernible. The nucellus is free only at its very apex from the integument, which is fused with the epimatium throughout.

Although the four sub-genera of *Podocarpus* differ from one another considerably in the structure of the female strobilus, they agree in possessing an inverted ovule with integument and epimatium closely united. The genus *Dacrydium*, however, includes one group of species in which the ovule is inverted and another in which at maturity it is almost erect, but in all cases the integument is entirely free from the epimatium and the nucellus from the integument. The ovulate cones of four species of *Dacrydium* were

DIAGRAM 5. *Dacrydium Bidwillii*. A, vertical section through strobilus; B-F, successive transverse sections from axis to chalaza.

investigated—*D. Bidwillii*, *D. cupressinum*, *D. Colensoi*, and *D. intermedium*. Only in the first two were mature ovules obtainable.

D. Bidwillii, with two other New Zealand species, agrees with *Podocarpus* in the possession of an inverted ovule. The much reduced cone consists of a very short branch with a number of bracts at its base much resembling the scale-like vegetative leaves and one or two axillary inverted ovules at its apex (Pl. VII, Fig. 21; Diagram 5, A). The vascular supply of the cone, as in *Podocarpus*, is a ring of bundles and canals from which a small strand departs to each bract (Diagram 5, B). This is followed, in the case of the fertile bracts, by two bundles which fuse by their adaxial ends and

thus inversely oriented, depart into the base of the epimatium (Diagram 5, C and D). The two component parts of this strand soon separate and diverge considerably as they proceed up the dorsal face of the ovule (Diagram 5, E). At the chalaza each sends down a small concentric bundle into the base of the ovule (Fig. 22; and Diagram 5, F) and then dies out. The nucellus is entirely free from the integument, as is the integument from the epimatium (Diagram 5, A). The latter is strongly ribbed and well provided with large sac-like canals. It ripens dry and hard, save for a thickened corky ring at the micropylar end, which acts as an absciss layer.

DIAGRAM 6. *Dacrydium cupressinum*. A, vertical section through strobilus; B-E, successive transverse sections from axis to ovule.

Dacrydium Bidwillii, from its close resemblance in strobilar anatomy to *Podocarpus*, may therefore well be considered an intermediate form between the latter genus and the other members of *Dacrydium*.

The rest of the genus *Dacrydium* show a similarly reduced condition of the cone, with a few sterile bracts at the base and a few fertile ones above. The young ovules, which were examined only in *D. Colensoi* and *D. intermedium*, are distinctly inverted, and the epimatium, with a knob or knee at its chalazal end just as in most species of *Podocarpus*, is entirely free from the integument, but at an early stage encloses it completely (Fig. 23). In subsequent growth, however, the outer coat is outstripped by the rest of the

ovule, which is eventually forced up into a partially erect position (Diagram 5, A). At maturity the bract is difficult to make out, and the epimatium appears as merely a thin sheath around the lower third of the ovule. The vascular system of the mature cone, investigated only in *D. cupressinum*, is at its base very similar to that of *D. Bidwillii* (Diagram 6, B). The base of the ovule has been left very near the apex of the cone, and the two bundles to the ovule, which fuse and become inversely oriented as in other cases, pass directly into the chalaza after a very short course in the base of the epimatium (Diagram 6, C and D). The integument is entirely free from the nucellus (Diagram 6, E), and since it is now the only protective coat of the ovule, it is much firmer than in the other members of the Podocarpaceae which we have examined, and has become differentiated into two layers, a narrow but very firm outer one of heavy-walled stone cells, and a less rigid, thinner-walled one within. These seem to correspond with the hard and soft layers in the ripe integument of *Eupodocarpus*.

The resemblance in structure and orientation to the seed of *Podocarpus* which is displayed by the young ovules of this section of *Dacrydium* suggests that we have here to do with an instance of recapitulation, and that the erect and naked condition of the ovule at maturity is not primitive, but has been recently acquired.

This description of the anatomical features of the female strobilus of *Podocarpus* and *Dacrydium* agrees in most particulars with the observations of Van Tieghem, Strasburger, Stiles, Miss Gibbs, and others. The marked structural resemblance between the cone of these genera and that of the Abietineae is noteworthy and will be discussed more fully under our consideration of the relationships of the Podocarpaceae.

Saxegothea and *Microcachrys* were not investigated by the writer, but good accounts of the strobilar anatomy of the two genera have been published by Thomson (17), Tison (18), and Stiles (11).

The female cone in *Saxegothea* consists of a considerable number of spirally arranged fertile bracts above and of numerous sterile ones below, which pass gradually into the vegetative leaves. The whole structure is thus more cone-like in appearance than anything we have yet described in the family. Each fertile scale bears near its base a single ovule which is semi-erect from the first and is provided with a thin epimatium entirely separate from the integument. Tison calls attention to the dissimilarity in structure between the tissue just below the insertion of the ovule (and of which the epimatium is obviously a continuation) and that of the rest of the bract. From the vascular supply of the cone axis, which is formed by a ring of bundles and is surrounded by a row of cortical canals, a single vascular strand passes off to each bract whether sterile or fertile (Diagram 8, E). In the case of the latter, however, two bundles, one from each end of the foliar

strand, are soon cut off, and rotating through an angle of 180° , assume an orientation inverse to that of the parent bundle. In the simplest cases, these two bundles, without further division, pass into the base of the ovule. According to Tison, a third median bundle is usually formed, and in several cases this 'upper series' was found to be composed of five bundles, only two of which entered the ovule. The bract supply also often divides into a series of strands, of which only one persists to the end.

DIAGRAM 7. *Phyllocladus glaucus*. A, vertical section through portion of strobilus; B, transverse section. *Cephalotaxus*. C, vertical section through portion of strobilus; D, transverse section.

Attention will be directed later to the resemblance between the megasporophyll of *Saxegothea* and those of other Conifers, such as *Athrotaxis*, in which reduction has undoubtedly taken place.

Conditions in *Microcachrys* are very similar to those in *Saxegothea*, but are more simple. The cone-like female strobilus bears numerous fertile scales which somewhat resemble the small appressed vegetative leaves and

are continuous with them. A poorly developed epimatium surrounds the partially erect ovule. A single bundle departs from the axis to the fertile bract, and from its ventral side is given off a solitary strand with inverse orientation, which soon divides into two and enters the base of the ovule (Diagram 8, D).

Three species of *Phyllocladus* were investigated by the writer: *P. glaucus*, *P. trichomanoides*, and *P. alpinus*. In the last two, the female cone consists usually of one or two fertile bracts in a rather spherical head, but in *P. glaucus* there are from ten to twenty bracts, and they are arranged spirally. In all three species the short stout bract bears at its base a single erect ovule, the lower half of which is surrounded by a symmetrical papery epimatium, late in developing (Diagram 7, A). The vascular system of the cone axis consists of a cylinder of bundles which sends forth a single strand to supply each bract (Diagram 7, B). From the sides of the gap, as in *Podocarpus*, arise two bundles which orient themselves inversely and pass directly into the base of the ovule. The integument is entirely free from the nucellus and from the epimatium, and is differentiated into two layers, an outer sclerotic one and an inner one less firm in texture.

Discussion of the morphological significance of the strobilar anatomy in the Podocarpaceae will be reserved until after an account of the development of the male and female gametophytes and of the embryo.

MALE GAMETOPHYTE.

The important facts in the history of the male gametophyte of the Podocarpaceae are now comparatively well known, and the main result of the present investigation has therefore been to confirm previous observations and to extend them to a few more species. Material for study was available of the early stages of the male gametophyte in *Podocarpus japonicus* and *P. macrophyllus*; of mature or nearly mature pollen, and usually of tube formation, in *P. Totara*, *P. spinulosus*, *P. spicatus*, *P. ferrugineus*, *P. dacrydioides*, *Dacrydium intermedium*, and *D. cupressinum*, and of almost everything from archesporium to the formation of male cells in *P. Hallii* and *P. elatus*.

The sporogenous cells are thin walled and have large nuclei. They are first differentiated when the sporophyll is very small and increase rapidly in number as the sporangium grows larger. Around the sporogenous tissue is eventually developed a narrow tapetum, and the sporangium wall outside this consists of a few rows of thin-walled cells and of a very stout, much thickened epidermis. The microspore mother-cells are eventually rounded off, and each divides into a tetrad of spores.

The development and structure of wings on the microspore was carefully observed, for their presence is one of the most important features of the pollen of the Podocarpaceae, and serves to distinguish the family from

all other Conifers save the Abietineae. Two large wings on each spore were found throughout the sub-genera *Eupodocarpus* and *Stachycarpus*. The occurrence of three wings in *Podocarpus dacrydioides*, which had previously been reported, was found to be the normal condition for this species, though four were occasionally present. The wings here are somewhat smaller in proportion to the grain than in the other sub-genera. The mature pollen of *Dacrydium intermedium* shows two well-developed wings, but in *D. cupressinum* they are very much reduced in proportional size, and have a wrinkled and shrunken appearance (Pl. V, Fig. 6 and Pl. VII, Fig. 24). In this species, however, young pollen-grains with but a single nucleus were occasionally found in the nearly ripe sporangia, and these were distinguishable by the possession of two wings large in proportion to the size of the spore and of the normal *Podocarpus* type (Fig. 5). The more rapid growth of the spore finally results in their partial obliteration. The pollen of *Microcachrys* as described by Thomson has normally three wings, but occasionally four, five, or six. In *Saxegothea* they are altogether lacking. The microspore of *Phyllocladus* shows two wings, irregular in shape and much reduced. It seems entirely probable, as we shall endeavour to show, that the wings in the Podocarpaceae are the exact homologues of similar structures in the Abietineae, and that their absence or poor development in certain genera and species is due to reduction.

The details of gametophytic development agree with those reported in previous accounts. Two prothallial cells are cut off and persist, giving rise to a varying amount of vegetative tissue. In the mature grain of *Podocarpus* this usually consists of from six to eight cells, and there are in addition a stalk and a body-cell and a tube nucleus. All of these except the body-cell are eventually represented by free nuclei, among which the tube and stalk nuclei are usually indistinguishable. In *Dacrydium cupressinum* there are from four to six, but in *D. intermedium* never more than four, prothallial cells and vegetative nuclei.

The germination of the pollen-grain was observed in *P. Totara*, *P. Hallii*, *P. elatus*, *P. ferrugineus*, *P. spicatus*, *P. dacrydioides*, and *Dacrydium cupressinum*. Into the tube as it penetrates the nucellus passes the procession of free nuclei, which here lose their spherical shape and become considerably elongated. The body-cell, conspicuous by its size, is the last to leave the pollen-grain and migrates slowly down through the nucellar tissue (Pl. V, Fig. 4). Just before fertilization its nucleus divides into two, of which one is naked, and the other, destined to unite with the egg, is surrounded by a dense body of cytoplasm (Pl. IX, Fig. 50). In *Stachycarpus* these male cells are buried close to the neck of the archegonium by the growing endosperm (Pl. V, Fig. 2).

THE FEMALE GAMETOPHYTE AND EMBRYO.

Our knowledge of the female gametophyte and embryo of the Podocarpaceae has up to the present been practically confined to the results of investigations by Miss Young (21) on *Phyllocladus alpinus*, by Coker (2) on *Podocarpus coriaceus*, and by Stiles (13) on *P. latifolius* and *P. macrophyllus*, and to the fragmentary observations of Miss Gibbs (4) on several other species of the genus. It is now possible to add to this an almost complete account of the history of the macrogametophyte and embryo of *Podocarpus Totara*, *P. nivalis*, *P. ferrugineus*, *P. spicatus*, *P. dacrydioides*, and *Dacrydium cupressinum*, together with considerable information concerning *P. elatus*, *P. spinulosus*, *P. vitiensis*, and *D. Bidwillii*.

In *Eupodocarpus* only a few months elapse between the appearance of the young female cones and the ripening of the seed, but the length of the period, besides being subject to a considerable amount of specific and individual variation, depends also on climatic influences. It is very much shorter in the case of *P. nivalis* from the Southern Alps of New Zealand than in *P. elatus* from sub-tropical Queensland. The development of *P. Totara* in the North Island of New Zealand may be taken as fairly typical. Here the female cones are noticeable early in October, and pollination occurs about the middle of the month. The ovules reach nearly two-thirds of their mature size by the latter part of November, when fertilization takes place, and the seeds ripen in February.

While the ovule is very small the megaspore mother-cell becomes differentiated from the rest of the nucellus by its greater size (Pl. VII, Fig. 25). Its very large nucleus soon divides, as do the nuclei of the two resulting cells, and a linear tetrad of megaspores is thus formed, although there are sometimes but three owing to the failure of one of the daughter cells to divide. Three of the four spores, usually those towards the micropyle, become abortive, and the single remaining one germinates to form the gametophyte. The occurrence in the mature ovule of two embryo-sacs, from the germination of two spores, was observed rather frequently in most of the species.

In all members of this sub-genus a group of uninucleate, densely protoplasmic cells makes its appearance around the megaspore mother-cell, and later becomes a conspicuous layer of 'spongy tissue' surrounding the spores and the developing embryo-sac (Fig. 26). Whether or not this corresponds strictly to a true tapetum, its function is surely to nourish the growing spore. The absence of spongy tissue has been reported by Coker in *P. coriaceus*, but its presence has been noted in all other members of the sub-genus investigated by other writers.

The details of spore germination are very much as in other Conifers. The nucleus soon divides and the resultant nuclei, which in their turn

multiply rapidly, come to lie against the wall, the centre of the gametophyte being filled by a large vacuole. For the immediately succeeding stages, information is available only in the cases of *P. Totara* and *P. Hallii*. In these species the vacuolated sac becomes lined with hundreds of free nuclei before wall-formation begins. The first walls isolate each nucleus from its neighbours and leave it in a distinct compartment, but in free communication with the vacuole. The nucleus lies at the inner end of this compartment, and is connected by protoplasmic fibrils with the edges of the inwardly growing walls, which eventually meet at the axis of the gametophyte. The sac is thus filled with radiating tubular 'alveoli' which are rarely interrupted by transverse walls. The division of each alveolus into a row of cells, however, takes place rapidly after the cessation of centripetal growth, and the gametophyte thus becomes a body of thin-walled uninucleate cells arranged in rows radiating from its centre. The spongy tissue disintegrates as the embryo-sac reaches its full size, but a very well-developed megaspore membrane, thick basally but becoming thinner over the apex, is laid down around the gametophyte (Pl. VI, Fig. 7).

The history of the female gametophyte and embryo of *Eupodocarpus* from this point onward was studied in *P. Totara*, *P. Hallii*, and *P. nivalis*, and is practically identical in the three species.

The archegonia appear soon after the sac is filled with tissue, and occur in an apical group of three to six (Fig. 7). As many as fourteen have been noted by Stiles in *P. latifolius*. They are arranged roughly in a circle, but are rarely or never grouped, always being separated from one another by one or more rows of cells.

Each archegonium arises from a superficial cell, which soon divides periclinally into a primary neck-cell and a primary central cell. The neck-cell is unchanged for some time, but the central cell, growing rapidly, becomes much elongated, and is filled with thin, vacuolate contents. The nucleus remains at the upper end, just below the neck. No further changes take place until the archegonium reaches its full size, when three deeply staining centres of radiating protoplasmic strands appear in the central cell, one at the base, one near the middle, and one just below the nucleus at the apex (Pl. VII, Fig. 27). Between the lowest two of these 'asteroids' arises a large vacuole. The primary neck-cell has meanwhile divided anticlinally or obliquely into an irregular group of cells. One row of jacket cells, distinguishable from the surrounding gametophyte tissue by their more dense contents and large nuclei, one or two in number, becomes differentiated around the central cell.

The nucleus of the central cell now undergoes division into egg nucleus and ventral canal nucleus, and although the actual mitosis was not observed, the two small resulting nuclei were several times seen in the upper portion of the cell. The egg nucleus remains for only a very brief time in this

position, but almost immediately drops to the centre of the archegonium just above the large basal vacuole (Fig. 28). The three asteroids now disappear. The ventral canal nucleus may immediately go to pieces, may persist unchanged for a time (Fig. 29), or may more rarely increase to nearly or quite the size of the egg nucleus itself. A definite ventral canal cell was never observed, nor were there indications of the subsequent activity of the ventral canal nucleus reported by Coker.

While the archegonia are thus reaching their mature development the other cells of the gametophyte are also dividing, and the embryo-sac at the time of fertilization has attained two-thirds or more of its final size. The nuclei of the endosperm cells are at first large, but after repeated cell-divisions decrease considerably in size. A narrow cone-shaped patch of tissue extending from the archegonial region to a little below the centre of the embryo-sac is composed of smaller and squarer cells with more protoplasmic contents than the others, and with several large nuclei. The multinucleate condition extends also to a few of the other endosperm cells, but at this stage most of them possess only a single nucleus.

The archegonia are now ready for fertilization and the pollen-tubes, which for over a month have been coming down through the nucellus, expand over the top of the embryo-sac and often burrow into it to some extent. A branch of the tube, entering and breaking through the neck of an archegonium, is closely followed by the two male nuclei. The functional one, surrounded by a layer of dense cytoplasm, is about half the size of the egg nucleus itself, which it immediately approaches, and to which it becomes closely pressed. At this time the second male nucleus and several of the accompanying prothallial nuclei may be observed in the upper portion of the archegonium (Pl. V, Fig. 1). The contents of both sexual nuclei are granular, and the two are very similar in other details. The male, which is much flattened against the female, is for some time separated from it by a limiting membrane, but this eventually breaks down and the resulting fusion nucleus soon prepares to divide.

The first mitosis takes place in the position of the egg nucleus, and the two small daughter nuclei, accompanied by most of the dense cytoplasm of the egg, migrate to the base of the archegonium and increase greatly in size. Each soon divides again (Pl. VII, Fig. 30), as do the four which result, so that eight rather large free nuclei, irregularly arranged, fill the lower portion of the archegonium. Each of these again divides, and the resulting sixteen become separated from one another by the first-formed walls of the pro-embryo (Fig. 31). The cells are arranged in three tiers: a single binucleate cell at the very base, which will form the embryo, a suspensor tier of from seven to nine cells above this, and a rosette tier at the very top of a few cells which soon go to pieces. The suspensors now begin to elongate (Pl. VIII, Fig. 32) and to push the embryo cell, protected by a thick cellulose

cap (Pl. VII, Fig. 31), down into the endosperm. The details of proembryo formation are thus somewhat different from those reported by Coker for *P. coriaceus*, where there are fourteen suspensors and fourteen rosette nuclei.

The cells of the 'cone' into which the embryo is now pushed are packed with starch. Most of the remaining tissue of the embryo-sac becomes multinucleate at this time. The endosperm soon reaches its full size, and often grows up over the archegonia to some extent. Several of the archegonia usually remain unfertilized, and their nuclei may either go to pieces directly or first become broken up amitotically into many unequal fragments.

The embryo cell, which is pushed far into the endosperm before it shows signs of multiplication, divides first by a longitudinal wall and then by transverse ones. Instances of budding or of single suspensors giving rise to embryos were only rarely observed. One embryo eventually outstrips the others and is the only one to reach maturity. All the cells of the endosperm now become filled with starch, and in their midst lies the developing dicotyledonous embryo (Pl. VIII, Fig. 33).

The marked resemblance between *Eupodocarpus* and the Abietineae in the development of the female gametophyte and embryo, and the equally strong dissimilarity between this sub-genus and the Araucarineae, deserve emphasis, and are of much importance in determining the affinities of the Podocarpaceae.

Podocarpus dacrydioides, the only member of the sub-genus *Dacrycarpus* under investigation, resembles *Eupodocarpus* in the completion of its entire reproductive cycle in a single season. The young female cones, in the North Island of New Zealand, first become noticeable early in October and are pollinated about the middle of the month. Fertilization probably occurs at this latitude in late November or early December, but all subsequent stages of this species were obtained in the extreme south of the South Island, where the season is undoubtedly much later, and here fertilization was found to take place in the latter part of January and the fruit to mature in February and March.

The megaspore mother-cell appears while the ovule is still very small, and is surrounded by a zone of spongy tissue which is not so well developed as in *Eupodocarpus*. The mother-cell gives rise to a linear tetrad of which usually only one spore germinates, although the frequent occurrence of two embryo-sacs in the ovule indicates that this is not always the case. The early stages of the gametophyte resemble those of *Eupodocarpus*, and consist of an enlarging vacuolate sac lined with nuclei. The subsequent centripetal growth of endosperm could not be traced, for this had already taken place in the next later material examined. Most of the young endosperm cells have but one nucleus, except in the cone-shaped region below the archegonia, where they are multinucleate. A megaspore membrane is present around the embryo-sac, but is much less well developed than in *Eupodocarpus*.

The archegonia are from five to twelve in number, and occur in a circle around the apex of the gametophyte, but are never directly adjacent to one another (Pl. VI, Fig. 12). Each arises from a superficial cell which divides periclinally into primary neck-cell and primary central cells. The latter, which rapidly becomes elongated, has at first a very thin and foam-like contents filled with many small vacuoles, but no structures comparable to the 'asteroids' of the young central cell of *Eupodocarpus* were observable. One row of uninucleate, or binucleate, densely protoplasmic jacket cells becomes differentiated around the archegonium.

The small nucleus remains in the upper portion of the central cell until it has divided into egg and ventral canal nuclei, after which the egg nucleus drops to about the middle of the archegonium and becomes surrounded by a conspicuous layer of dense cytoplasm (Pl. VIII, Fig. 34). The ventral canal nucleus, even if it is always formed, must usually go to pieces immediately. In the comparatively few cases where its presence could be determined, it was almost always as large as the egg nucleus itself, and the two were lying close together (Fig. 35). The many small vacuoles now coalesce into a single large basal one, and the cytoplasm of the egg becomes of uniform density throughout. Meanwhile the primary neck-cell has divided first anticlinally and then periclinally into a neck two or three tiers of cells in height. The endosperm grows up over the archegonia to such an extent as to leave the neck at the bottom of a small pit, or even sometimes to close it over entirely and prevent the access of a pollen-tube.

In the details of fertilization, formation of the embryo, and history of the endosperm this species agrees almost entirely with *Eupodocarpus* (Figs. 36, 37, and 38), but the embryo-sac is much wider and more nearly spherical, and its cells, beginning with a few towards the outside at the time of fertilization and increasing rapidly in number until they include almost all the endosperm, become completely filled with a mucilaginous contents (Fig. 38).

The female gametophyte of *Podocarpus dacrydioides* differs chiefly from that of *Eupodocarpus* in the poorer development of spongy tissue and megaspore membrane, in the absence of asteroids in the central cell, in the presence of a dense cytoplasmic sheath around the egg nucleus, and in the occurrence of mucilaginous contents in the endosperm cells.

The sub-genus *Stachycarpus*, however, is distinct from the two preceding groups in the structure and development of its female gametophyte and embryo. *Podocarpus spicatus* and *P. ferrugineus*, the only species examined, were found to be very similar.

The reproductive cycle covers a period of nearly eighteen months. Young female strobili appear early in October, and are pollinated two or three weeks later, but the embryo-sac, which attains considerable size, is not ready for fertilization until a year from the next January. Ripe fruit is produced in March.

The megaspore mother-cell is distinguished early by its rapid increase in size and by its large nucleus. It gives rise to a linear tetrad of spores, of which the one nearest the base germinates (Pl. VIII, Fig. 39). Two spores occasionally develop into gametophytes. Spongy tissue seems to be entirely absent (Pl. VI, Fig. 17 and Pl. VIII, Fig. 39), for the cells immediately surrounding the spores and the young sac possess no dense protoplasmic contents and are not distinguishable in any way from the other tissue of the nucellus. Except for this absence of spongy tissue the young gametophyte conforms to the usual coniferous type.

The next stages obtainable, shortly after the endosperm had completed its centripetal growth, showed it to be composed of uninucleate cells and surrounded by a strongly developed megaspore membrane. The embryo-sac of *P. spicatus* is approximately spherical in shape and about half as large as that of *P. ferrugineus*, which is slightly obovate and nearly twice as long as broad.

The archegonia appear early, while the sac is less than half its final size, and are usually two in number, though three sometimes occur. Each archegonium arises from a superficial cell, which divides into neck-cell and central cell. The latter becomes greatly elongated and the mature archegonia are often 2 mm. in length, perhaps the largest among vascular plants. The nucleus remains near the neck and the contents of the central cell are thin, vacuolate and without asteroids. A single layer of jacket cells becomes sharply differentiated, and its members, uninucleate at first, soon contain several nuclei. The primary neck-cell divides anticlinally into a rosette of from eight to twelve cells, and this single tier is the final condition of the neck.

Meanwhile the endosperm has increased greatly in size and at the apex has grown up around the necks of the archegonia and enclosed the ends of the waiting pollen-tubes and the male cells which are destined to effect fertilization (Pl. V, Fig. 2 and Pl. VIII, Fig. 40). Any archegonium towards the neck of which no pollen-tube has by this time approached is now entirely sealed over by the sterile gametophytic tissue and prevented from being fertilized. A cone-shaped region below the archegonia, especially long and narrow in *P. ferrugineus*, is early differentiated as in the other species by its smaller and more rectangular cells and by the many nuclei which they contain. The remaining cells of the embryo-sac become multinucleate before fertilization, and the number of nuclei in each cell seems to be considerably greater than in the other two sub-genera.

As the central cell approaches maturity its nucleus divides into egg and ventral canal nuclei (Figs. 2 and 41). The latter may immediately go to pieces or may persist for some time, and in rare cases increase considerably in size. The egg nucleus sinks but slightly below the neck, always remaining in the upper quarter of the archegonium, and now grows very large (Figs. 2 and 40). It has loosely granular contents and is usually surrounded

by a cytoplasmic sheath. The contents of the egg itself become much more dense and contain a few rather small vacuoles, but nothing corresponding to the large basal one of the other species. Dense angular bodies, probably protein in nature, are scattered through the cytoplasm. The wall of the egg is much thicker than in other members of the genus and is full of large sieve-like pits opposite the jacket cells. There are now two or three rows of the latter instead of only one.

The act of fertilization was not observed, but the larger of the two male nuclei, which is surrounded by a cytoplasmic body and is doubtless functional, is about half the size of the egg nucleus. The figure of the first mitosis is very small, and occurs within the nucleus (Fig. 42), which now loses its membrane and becomes irregular in shape but remains in its original position until after the second mitosis. The four nuclei which result from this lie close together and surrounded by a sheath of dense cytoplasm drop slowly towards the bottom (Fig. 43), followed by much of the contents of the egg. Above them develops a large vacuole (Fig. 44), separating the basal, proembryonic region of the archegonium from its upper sterile portion. Nuclei often are present in this upper part, and though some of them may perhaps be derived from the ventral canal or second male nuclei, it seems much more likely that they are all jacket-cell nuclei, for these are often seen breaking through into the archegonium as it begins to go to pieces.

The four nuclei of the proembryo now divide and eventually give rise to about sixteen, which are crowded irregularly into the very narrow base of the archegonium (Pl. IX, Fig. 45). At this time walls are first formed. Some of the cells immediately undergo division, and the regular number and arrangement found in the proembryos of other species is not observable here. The rosette tier is very poorly developed, and almost immediately goes to pieces. The suspensors are usually nine in number and often cut off cells from their upper ends. The embryogenous tier consists not only of a large binucleate terminal cell, as in the other members of the genus, but of a group of cells between it and the suspensors. This group is small at first, but its members immediately begin to divide, and in *P. spicatus* may number from twenty to forty before the suspensors elongate to any extent (Fig. 46). In *P. ferrugineus* there are rarely more than eight or twelve cells in this tier (Fig. 47) until it has been pushed far into the endosperm. In the former species, the terminal cell, long, pointed, and often serving for a time as a protecting cap, is usually sloughed off. In *P. ferrugineus* the apical cell is blunter, but is provided with a thickened tip and almost always persists. The embryos show no signs of budding and are forced far down into the sac, where they begin a rapid growth. The later development of the embryo could not be followed, but it doubtless becomes dicotyledonous, as in the rest of the family. Starch is laid down in the axial region immediately after fertilization and is soon deposited throughout the whole endosperm.

The female gametophyte and embryo of *Stachycarpus* thus differ in several important particulars from those of the other sub-genera of *Podocarpus*. The reproductive cycle is extended through two seasons; there is a very slight development of spongy tissue; the number of archegonia is much reduced, but their size is greatly increased; the neck is composed of but a single tier of cells; the first two embryonic mitoses take place in the upper portion of the archegonium, and the embryo almost from the first is composed of a body of cells instead of one. Several of these features are shared by the genus *Dacrydium*, but a still more striking resemblance is presented by the female gametophyte and embryo of *Cephalotaxus*.

The only material at hand for a study of the female gametophyte in the sub-genus *Nageia* was some very young ovules of *P. vitiensis*. The embryo-sac in these was still very small, but was surrounded by a layer of spongy tissue, distinct but not as strongly developed as in *Eupodocarpus*.

In the genus *Dacrydium* material was available for the study of the early stages of the female gametophyte in *D. Colensoi* and *D. intermedium*, two closely related species. The later history was investigated in *D. cupressinum*, and some information was also obtained as to the structure of the mature archegonium and the development of the embryo in *D. Bidwillii*.

In that portion of the genus represented by this last species the reproductive cycle is apparently completed in a single season, for in the Southern Alps in February there seemed to be but one generation of ovules, and these contained young or mature embryos. In *D. cupressinum* and its related species, however, tiny ovules which are just being pollinated are visible on the same branches with those of the previous year, which have reached their full size and in which fertilization and the development of the embryo is taking place. The life-history of this species therefore extends through two seasons, as in *Stachycarpus*.

The growth of the megaspore and its germination into a linear tetrad occur in very much the same manner in *Dacrydium* as in *Podocarpus*, but in the former genus there is a less conspicuous development of spongy tissue. The vacuole, lined with free nuclei, increases in size and is very early filled by the centripetal growth of endosperm tissue, which occurs before the embryo-sac has reached half its final size. The endosperm cells at first contain but a single nucleus. A cone of multinucleate and more densely protoplasmic cells soon appears below the archegonial region, and by the time of fertilization there are several nuclei in every cell of the embryo-sac. The prothallial tissue often grows up over the archegonia.

The latter appear while the sac is still very small, and in *D. cupressinum* are three in number at the apex of the gametophyte, each arising from a superficial cell and separated from the others by sterile tissue. The initial cell soon divides into primary neck and central cells. The former

divides anticlinally, but the mature neck is never more than one tier in thickness. The central cell, which elongates greatly and becomes filled with thin, vacuolate contents, is surrounded by a single row of uninucleate jacket cells. Its cytoplasm gradually grows more dense and displays at least one 'asteroid', which is associated with the nucleus in the upper portion of the archegonium (Fig. 48). A large basal vacuole also develops. The nucleus of the central cell, while still just below the neck, divides into ventral canal and egg nuclei. The former eventually goes to pieces, but the latter drops a little way into the cytoplasm and increases greatly in size, until it almost fills the top of the archegonium.

The male nuclei resemble those of *Podocarpus*, one being naked and the other surrounded by a mass of cytoplasm. Fertilization was not observed, but the development of the proembryo was followed and agrees with that described for *Eupodocarpus*. The young embryo, however, as it is carried down by the suspensors invariably buds or divides, and single suspensors often develop embryos (Fig. 49).

There are two archegonia in the embryo-sac of *D. Bidwillii*, and they are somewhat sunken in the endosperm (Pl. V, Fig. 3). Each is surrounded by one layer of jacket cells and has a neck but a single tier of cells in thickness. The egg nucleus is very large. Fertilization and the development of the proembryo were not observed, but the young embryos present in almost all the ovules examined much resemble those of *D. cupressinum*, and show a similar tendency to become subdivided.

Dacrydium resembles *Eupodocarpus* and *Dacrycarpus* in the development of the proembryo, and *Stachycarpus* in the small number of archegonia, the single tier of neck-cells, the large egg nucleus, and, in most species, the length of the reproductive cycle.

Our knowledge of the female gametophyte and embryo of *Phyllocladus* is due to the investigations of Miss Kildahl (6) and Miss Young (21) on *P. alpinus*.¹ The reproductive cycle in the genus extends through only a few months. Spongy tissue is conspicuous and there is a strong megaspore membrane. The archegonia are usually two in number, and are sunken in the endosperm. The neck is a rosette only one tier of cells in thickness. A ventral canal nucleus is present. There are at least eight nuclei in the proembryo before wall formation and the mature embryo is dicotyledonous. The endosperm cells are all multinucleate.

The structure of the female gametophyte and embryo of *Phyllocladus*, as far as it is known, agrees more closely with that of *Dacrydium*, especially in the reduced number of archegonia and the structure of the neck, than with that of any other member of the family.

¹ The writer was able to collect material in New Zealand illustrating the female gametophyte of *P. glaucus*, *P. trichomanoides*, and *P. alpinus*. The results of an investigation of these three species will be published in the near future.

AFFINITIES OF THE PODOCARPINEAE.

(a) Relationship to Abietineae and Araucarineae.

We must now discuss briefly the theories which have been proposed as to the affinities of the Podocarpineae, and consider what light is thrown on the phylogeny of the family by the present study of its reproductive structures.

The view of Celakovsky and his school that the Taxaceae, including the Podocarpineae, are closely related to *Ginkgo*, and are among the most primitive of the living Coniferales, has become entirely discredited since our knowledge of the female gametophyte in the Gymnosperms has grown more complete.

A much more reasonable theory, however, and one which apparently has a very wide acceptance at the present time, has been put forward by Thomson, Tison, Stiles, and others. These writers consider the Podocarpineae to have arisen from the araucarian Conifers through forms closely resembling *Saxegothea* and *Microcachrys*, and base their conclusions on the resemblance between these genera and the Araucarineae in the external features of the female cone, the development of the male gametophyte, the structure of the ovule, and the vascular anatomy of the ovuliferous appendage. They regard *Dacrydium*, and more especially *Podocarpus*, as advanced and specialized members of the family.

From this hypothesis the writer entirely dissents. After considering certain objections to the arguments on which it is based, he will endeavour to construct a line of evidence in support of the view that the genus *Podocarpus* is the most primitive member of the family; that it is nearly related to the Abietineae; that from ancient forms close to it the other genera of the Podocarpineae, and probably the Taxineae as well, have arisen through divergent lines of ascent, and therefore that instead of being derived from the typical araucarian Conifers the Podocarpineae may much more readily be traced back to the ancestral abietineous stock from which both families, Araucarians and Podocarps, along a somewhat parallel line of development, have been evolved.

The fact that the female strobilus of *Saxegothea*, *Microcachrys*, and *Pherosphaera* is well developed and composed of a considerable number of fertile bracts, in comparison to its reduced condition in most other members of the family, has been one of the main arguments brought forward to show the very primitive constitution of these three genera. The sub-genus *Stachycarpus* of *Podocarpus*, however, includes some species in which there are many fertile bracts, and others, closely related, where the female cone is reduced to a single ovule. *Phyllocladus glaucus* has a well-developed strobilus with from ten to twenty ovules, but there are only one or two in

the other New Zealand species of the genus. The degree of strobilar development may be extremely variable and is an unreliable criterion in determining affinities.

The gradual transition between fertile bracts and vegetative leaves in *Saxegothea* and *Microcachrys* has been compared with similar conditions in *Araucaria*, but since the same phenomenon occurs in *Cunninghamia*, *Cryptomeria*, and to a certain extent in other Conifers, it loses its significance.

The fact that the three genera in question which have well-developed cones consist of very few species and are limited in distribution has been emphasized by Stiles as an instance of the general principle that such restricted types are primitive. Dominance and widespread distribution in a group of plants are considered *prima facie* evidence for its recent origin, since the possession of more perfect adaptations gives it superiority over less favoured and therefore moribund ancestors. This argument is open to considerable criticism, for there is good reason to believe that many groups dominant to-day, such as the genus *Pinus*, are comparatively primitive. Unspecialized and plastic organisms, 'generalized types,' are more easily adapted to changing conditions, and therefore are often more widely distributed than those which are more 'advanced' and which through their higher degree of specialization and complexity are confined to a limited environment.

The occurrence in the pollen-grains of both Araucarineae and Podocarpineae of two primary prothallial cells which later undergo repeated division has also been cited as evidence of the affinity of these two families. This resemblance in the male gametophyte appears indeed most easily traceable to a common ancestry but to one which must have been abietineous in affinity, for the occurrence in *Ginkgo* and the Abietineae of pollen with but two prothallial cells makes it extremely probable that this is the primitive condition for all the Coniferales, and that the subsequent increase in prothallial tissue which takes place in the Podocarpineae and Araucarineae is a comparatively recent development and not the persistence of a primitive structure.

The entire absence of wings in the pollen of *Saxegothea* (8) and their somewhat irregular development in *Microcachrys* have led Professor Thomson (16) and others to believe that the winged condition in the Podocarpineae has arisen entirely independently from that in the Abietineae, and that the early stages in its development are represented in *Microcachrys*, where three, four, five, or six wings may be formed. The occurrence of both winged and wingless pollen in *Tsuga*, and the entire absence of wings in *Larix* and *Pseudotsuga* alone among the Abietineae, however, are almost certainly the results of reduction and loss, for these genera are surely not the most primitive members of their family. A similar explanation

in the case of the Podocarpineae seems much better than the one Professor Thomson proposes, and must clearly be correct if we consider *Podocarpus* as primitive and *Saxegothea* as specialized. The evidence from *Dacrydium cupressinum* bears out the reduction theory, for in this species there are two or three well-developed wings on the young spore, but these become withered and almost obliterated in the mature pollen, which thus approaches the condition found in *Saxegothea*. *Phyllocladus* is obviously a reduced and specialized genus, and its pollen wings show a correspondingly great reduction. *Podocarpus dacrydioides* with its three or four wings is intermediate between typical *Podocarpus* of two wings and *Microcachrys*, where the irregular development of these structures resembles more the behaviour of a vanishing character than of one just making its appearance. It seems therefore much more in accord with all the facts to consider the pollen of *Saxegothea* and *Microcachrys* as reduced rather than primitive in structure, and to regard the whole male gametophyte of *Podocarpus* as what it so obviously seems to be, a structure closely related to that of the Abietineae.

Certain features of the ovule common to Podocarps and Araucarians have been brought forward as indications of relationship between the two groups. The fact that cone scales bearing but a single ovule are almost entirely confined to these families is perhaps significant, but certain species of *Juniperus* show a similar condition, and it is well known that several ovules to each scale were commonly possessed by the ancient Araucarineae.

The stigma-like nucellar outgrowth described by Tison for *Saxegothea* has been compared to the slight protrusion of this portion of the ovule in *Araucaria*, but among other Podocarps anything comparable to it is entirely absent, and it seems a generic peculiarity with little significance.

Dacrydium, *Saxegothea*, and *Microcachrys*, in common with the Araucarineae, possess a nucellus entirely separate from the integument, and this has sometimes been considered a primitive feature and evidence for the relationship of the two groups. But the degree of fusion between integument and nucellus is a very variable character, as is shown, for example, by the sub-genus *Stachycarpus*, which has one species (*P. spicatus*) with the nucellus free only at the very tip, and another (*P. ferrugineus*) where it is almost entirely distinct from the integument. Freedom of the nucellus has also been attained in such widely separated forms as *Cunninghamia*, *Cryptomeria*, *Sciadopitys*, *Callitris*, and *Juniperus*, and its occurrence in the two families under consideration may therefore be best explained as the result of somewhat parallel but quite independent development.

The anatomy of the cone scale, however, has been most emphasized by supporters of the araucarian origin of the Podocarpineae, and since evidence derived from this source furnishes some of the strongest arguments for the writer's contentions as well, the subject is worthy of careful consideration.

The earliest accounts of vascular anatomy in the female cone of Conifers are in general more clear and comprehensive than those published more recently. Van Tieghem (19), in 1869, investigated the 'female flower' in everything from Cycadales to Gnetales, and was the first to apply anatomical criteria to the question of the morphology of the ovuliferous scale. Strasburger, in a number of his works, notably 'Angiospermen und Gymnospermen' and 'Coniferen und Gnetaceen', extended and amplified Van Tieghem's observations. A notable paper by Radais (9), and one which for some unaccountable reason seems to have escaped entirely the attention of subsequent writers on the subject, gives a clear account of some very careful and extensive investigations on the anatomy of the female cone, both young and mature, among the Abietineae, Taxodineae, and certain Araucarineae. The results obtained by these writers agree in emphasizing the fundamental similarity of the vascular supply to the strobilar appendage among all living Gymnosperms above the Cycadales. In every case the sterile leaf or bract subtends an axillary structure which is sometimes fused entirely with the bract, and is often much reduced, and which bears the ovule or ovules. This double appendage (leaf and scale) is supplied by a double vascular system from the central cylinder of the axis.

In *Ginkgo* a normal double foliar strand departs from the stele and passes into the subtending leaf. From each side of the gap thus caused springs a bundle, and these two, each of which soon bifurcates, enter the megasporophyll or axillary ovuliferous stalk, which is thus obviously a branch provided with a vascular cylinder of four bundles.

In the Abietineae (Diagram 8, A) conditions are very similar save that the subtending structure is a bract, not a foliage leaf, and that the axillary ovuliferous scale is not so evidently a branch. A strand precisely resembling the foliar bundle supplies the sterile bract. After its departure two bundles arise from the cylinder, one from each side of the gap, and approach each other by their adaxial ends, xylem outside and phloem inside, thus assuming an orientation which is inverse to that of the bract supply. The latter passes without division into the free bract, but the two scale bundles during their course through the cortex fuse into an arc or partial cylinder, which divides later into three parts. The small median bundle undergoes no further branching, but the two large lateral ones each give off a strand to an ovule and then by repeated division form the wide vascular arc which enters the lamina of the scale. The cone scale is thus precisely like an axillary branch in the origin of its vascular supply. A striking similarity is evident between the strobilar anatomy of the Abietineae and that of *Podocarpus* (Diagram 8, A and B).

In the Taxodineae and Cupressineae the bract and scale are more or less completely fused with one another, but where the latter is dominant, as in *Sequoia*, the vascular supply is double at its origin, and consists

DIAGRAM 8. Series illustrating the vascular structure of the strobilar appendages, from con-axis to ovule or beyond, in certain Conifers. A, Abietineae; B, *Podocarpus elatus*; C, *Podocarpus dactyloides*; D, *Microcachrys*; E, *Saxegothea*; F, *Athrotaxis*.

of a single bundle for the bract, followed by two others, derived from the sides of the gap and inversely oriented, which usually fuse, divide again into three, and then form a wide arc in the scale.

In *Cryptomeria*, *Athrotaxis*, and *Cunninghamia*, however, only a single bundle leaves the axis to supply the double appendage, but this single bundle, normally oriented, consists of the bract and scale systems fused. Its two ends soon bend upward, adaxially, and in *Cryptomeria* a ring of separate bundles is eventually formed. Only the median lower one enters the free end of the bract, and all the rest, forming a wide arc precisely as in *Sequoia*, represent the scale series. In *Athrotaxis* (Diagram 8, F) and *Cunninghamia* only a few small bundles, sometimes only one, are cut off from the adaxially curving ends of the original single strand and become oriented inversely. These bundles, which are obviously all there is left of the scale or axillary system, merely supply the ovules. The lamina of the appendage, now almost entirely bract, is provided with a long row of normally oriented bundles derived from the main part of the original single strand.

Early writers considered that a condition very similar to that of *Cunninghamia* was present in *Araucaria* and *Agathis*, and that the scale system was here reduced merely to the ovular bundles, the bulk of the 'scale' being supplied by a wide arc of strands, normally oriented, which were derived from the main part of the original bract bundle. Certain recent investigators, notably Professor Thomson, have maintained that the scale in the Araucarineae is a single (aplosporophyllous) structure and not a double one, and that it is entirely homologous with the microsporophyll and the vegetative leaf. He calls attention to the fact that in this family there are only two series of bundles, one to the 'scale', and the other, with inverse orientation, to the ovule, but that in the Abietineae, Taxodineae, and Cupressineae, which he considers double-scaled or diplosporophyllous, there are three series: one to the bract; one, inverse to this, to the scale; and a third, with another inversion of orientation, to the ovule. The Podocarpaceae, according to this theory, have been derived from the Araucarineae through forms similar to *Saxegothea*, for conditions in this genus resemble those in *Agathis*, the bulk of the appendage being supplied by a row of normally oriented bundles derived from a single strand, and the upper series consisting of little or nothing beside the ovular supply. *Dacrydium* and *Podocarpus* are derived from this by the increase in comparative size of the ovule and by the 'basipetal development' of its vascular system, which eventually attains the dignity of an independent insertion on the axis quite distinct from that of the supply to the 'scale'. The Podocarps and Taxads are therefore regarded as simple-scaled forms which have developed their female strobilus by an amplification of that of the male, and which are sharply separated from those Conifers where the female cone is an axis bearing axillary ovuliferous short shoots. The Coniferales are on this supposition divided into two great groups which must have been separate almost since the assumption of the strobilar habit.

A monophyletic conception of the order, however, seems much more in accord with what we know of the history of the Coniferales, and they have been considered by almost all investigators to be a natural group descended from a single source. If this is true, their ancestors must either have been single-scaled forms, from which the double-scaled families of to-day have arisen, or else the cone with double scales is primitive, and has given rise to that with a single scale by reduction.

The former view has been adopted by several writers, notably by Stiles in a recent paper, and is apparently shared by Miss Gibbs as well. The primitive cone among the Podocarpaceae, and apparently among the Araucarineae also, is believed to have consisted of many spirally arranged sporophylls, in the axil of each of which was a single erect ovule. *Pherosphaera*, among living Conifers, seems most closely to approach this hypothetically primitive condition. The first sign of the ovuliferous scale was the appearance of an epimatium, inconspicuous and poorly developed at first, as in *Microcachrys*, *Saxegothea*, and some species of *Dacrydium*, but becoming larger until in *Podocarpus*, the most recent genus of all, the ovule is borne on the epimatium, which has been carried up on a stalk and is almost free from the 'scale'. In some such way as this the free ovuliferous scale of the Abietineae is supposed to have arisen, and the family is derived from a podocarpean-araucarian plexus. The living Araucarineae have departed from their *Pherosphaera*-like prototype only in possessing an inverted instead of an erect ovule.

This view of coniferous phylogeny will doubtless appeal strongly to those who believe that the Araucarineae are the most primitive members of the order, but the adoption of it necessitates an entire overthrow of the brachyblast theory of the ovuliferous scale of the Abietineae, which has until recently met with almost unanimous acceptance. This theory is well supported by facts, for, in addition to the mass of evidence derived from abnormalities, the vascular structure of the upper scale resembles precisely in its origin an axillary shoot. Much more cogent evidence than has yet been presented will be necessary to convince many persons that the ovuliferous scale of the Abietineae is only a glorified epimatium, and that its independent vascular supply represents merely a basipetal development of the ovular bundles. The homology between ovuliferous scale and epimatium is now pretty generally admitted, but the presence of the epimatium is logically accounted for only if it is derived from an axillary ovuliferous scale. To consider it a fortuitous outgrowth from the ventral face of the bract is not to explain it satisfactorily.

The other alternative, which regards as most primitive a cone with distinct double scales, resembling that of the living Abietineae, appears to the writer much more in agreement with all the facts. According to such a conception the epimatium is a vestige of the ovuliferous scale, not its

primordium, and the single-scaled series have arisen through either the fusion of the two appendages or the abortion of one of them. In the ascending series from the Abietineae to the Taxodineae and Cupressineae we may observe the gradual fusion of bract and scale, which usually results in the complete dominance of the scale and the almost entire disappearance of the bract and its vascular supply. In a few cases, however, notably *Athrotaxis* and *Cunninghamia*, where it is the bract which becomes dominant, the scale is so much reduced that its vascular system is represented by only a few small strands, sometimes only one, given off from the ventral face of the bract bundle, and serving merely to supply the ovules (Diagram 8, F). This condition of affairs is so precisely similar to that which we find in *Saxegothea* on the one hand, and in certain of the Araucarineae on the other, that it provides an illuminating suggestion, noted by earlier investigators, as to how the features of strobilar anatomy in these two groups have arisen. The hypothesis that in *Saxegothea* and the Araucarineae, as well as in *Cunninghamia*, the simple scale has been evolved by reduction from a more complex double one seems to the writer to be the most plausible yet suggested. Eames (3) in a contemporaneous paper brings forward strong evidence for the derivation of the araucarian cone-scale from a primitively double structure, and the results of anatomical investigations on the cones of *Saxegothea*, *Microcachrys*, and *Phyllocladus* seem to indicate that the simple scale in these forms has been evolved from the double strobilar appendage of *Podocarpus*, and that this genus is the most primitive of the family and stands close to the Abietineae.

This hypothesis, which instead of deriving the Podocarpineae from the Araucarineae considers them to have descended from the primitive Abietineae through forms somewhat resembling those found in the living genus *Podocarpus*, is supported by abundant evidence from our comparative study of the reproductive structures of the families in question.

In both *Podocarpus* and the Abietineae the female cone is composed of subtending sterile bracts, each provided with a single strand, and almost or quite free from the axillary ovuliferous structure. In both, the vascular supply of the latter arises as two separate bundles, each derived from a side of the bracteal gap, which approach each other by their adaxial ends, become oriented inversely to the bract bundle, and fuse. In both, this single fused bundle tends strongly to divide into three when it enters the base of the ovuliferous scale or epimatium. In most species of *Podocarpus* two distinct ovular bundles are given off from the main supply just as in the Abietineae, and instead of displaying inverse orientation these are usually concentric, a condition which Radais also observed in the ovular bundles of the Abietineae. In fact, it seems doubtful if the inversion in orientation of the ovular supply, 'the second inversion,' is at all common.

The main cone of *Podocarpus* and of its family is almost identical with

that of the Abietineae, for in both groups the microsporophyll bears but two sporangia, a very different condition from that obtaining in the Araucarineae, where the pollen-sacs are often very numerous. In fact, the Abietineae and Podocarpineae are the only Conifers which never have more than two microsporangia to a sporophyll.

These two families are also distinguished from all others of the Coniferales by the possession of winged pollen-grains, and attention has already been called to the strong resemblance between the whole male gametophyte of *Podocarpus* and that of the Abietineae.

Now that the investigations of Eames (3) in *Agathis australis* have given us a much fuller knowledge of the araucarian female gametophyte and embryo, we are able to compare the reproductive histories of the Araucarineae and Podocarpineae, and to observe their many points of difference. In the Araucarineae the megaspore membrane is poorly developed, the archegonia are very numerous and distributed over the whole upper portion of the embryo sac, and the fertilized egg develops *in situ* and gives rise to a proembryo furnished with an elaborate protective cap, such as is found nowhere else. Among the Podocarpineae, on the other hand, the megaspore membrane is conspicuous, the archegonia are fewer in number and gathered in an apical group, and the products of the fertilized egg sink directly to the bottom of the archegonium, where they produce a proembryo very similar to that found among the Abietineae and their allies. Indeed it is to the type occurring among these dominant Northern Hemisphere Conifers that the whole female gametophyte of *Podocarpus* and its family conforms, and it shows little similarity to that of the Araucarineae.

It is highly improbable that so many strikingly similar reproductive structures, both sporophytic and gametophytic, should have arisen in two independent groups and along two quite distinct lines of evolution. We are, therefore, forced to adopt the hypothesis that the Podocarpineae, and especially the genus *Podocarpus*, are more nearly related to the Abietineae than to any other family of the Coniferales, and since a great body of evidence is accumulating in support of the view that the Abietineae are the most ancient members of the Coniferous Order, *Podocarpus* must be considered as a relatively primitive type and the oldest genus of its family. The resemblances between the Podocarpineae and the Araucarineae are best explained as an inheritance of structures and tendencies from a very ancient group, ancestral to both and closely abietineous in affinities, from which the modern Araucarians have widely departed. In fact, it is perhaps more logical to consider the Araucarineae as derived from, rather than having given rise to, the Podocarpineae.

This hypothesis as to the phylogeny of the Podocarpineae has the advantage of explaining more facts than does any other which has been put forward.

It is in precise harmony with the view, now supported by an abundant evidence from many sources, that the Abietineae are the most primitive of all Conifers.

It assigns a logical position to the Podocarpaceae among the Coniferales, and does away with the necessity of dividing this otherwise homogeneous order into two sharply distinct series.

It explains the wide diversity in structure found among the members of the genus *Podocarpus* as the natural state of affairs in a primitive and generalized genus which has not yet become fixed or stereotyped.

It offers a reasonable explanation for the presence of the epimatium, as a structure homologous with the ovuliferous scale of the Abietineae, well developed and displaying its true nature in *Podocarpus*, but much reduced in the other genera. To consider the epimatium as a normal second integument which later expands into the ovuliferous scale, or to regard it as merely a ventral outgrowth of the sterile bract, are both theories which fail to explain it adequately in the presence of all the facts.

Finally, instead of so often appealing to parallel development to solve the riddle, it offers a reasonable explanation for the striking similarity between *Podocarpus*, which is considered primitive, and the Abietineae, and on this basis makes possible the construction of a clear and consistent natural classification for the various genera and species of the Podocarpaceae.

Numerous intermediate forms connect the widely differing sub-genera of *Podocarpus* with one another, but on a study of their reproductive structures alone it is impossible to determine definitely which group of species is the most primitive. *Eupodocarpus*, on the whole, may be so considered, for in the structure of its male and female gametophytes it resembles the Abietineae more closely than do any of the others. The poor development of spongy tissue and the small number of archegonia in *Stachycarpus*, with the almost complete freedom of the nucellus in one of its species, shows that this sub-genus has progressed considerably from the primitive abietineous condition. *Dacrycarpus*, with its fused bract and three-winged pollen, is obviously the most specialized member of the sub-genera.

A rather close series unites the parent genus with *Dacrydium* through *D. Bidwillii*, for this species, although it possesses the typical subulate leaves of its genus, has nearly the strobilar structure of *Eupodocarpus*, save that its nucellus, integument, and epimatium have all attained complete freedom from one another. *Dacrydium* proper has progressed much further, and only during a very young condition of the cone, when the epimatium, bearing an apical knob, completely surrounds the ovule, is the resemblance to *Podocarpus* at all marked. The female gametophyte of *Dacrydium* by the decrease in number and increase in size of its archegonia shows clearly a more advanced condition than does that of *Eupodocarpus*.

In *Podocarpus dacrydioides* the three-winged pollen-grain and the com-

plete fusion of ovule and epimatium to the sterile bract indicate a definite step in the direction of *Microcachrys* and *Saxegothea*. This species is also set off from the rest of the genus by its subulate leaves, numerous mucilage cells in the endosperm, and poor development of spongy tissue and megaspore membrane.

Microcachrys is close to *P. dacrydioides* in the possession of subulate leaves, three-winged pollen, and a single bundle in the bract (Diagram 8, C and D), but it has advanced, like *Dacrydium*, in the freedom of epimatium from integument. The upper vascular system has likewise grown very weak.

Pherosphaera is the last step in reduction, and here all traces of the epimatium have been lost, and the ovule has become axillary and erect, both of which features seem undoubtedly modern and not primitive.

In *Phyllocladus* the wings of the pollen-grain are disappearing, and the ovule has also attained a perfectly erect position and is surrounded symmetrically at its base by an epimatium obviously homologous with that in *Dacrydium* proper. The integument of the ovule in these two genera is divided in precisely the same way into a stout outer layer and a more soft inner one, and is free entirely from the nucellus. The female gametophyte, especially in the reduced number of archegonia and the structure of the neck, is very similar in both genera. All the evidence seems to indicate that *Phyllocladus*, though radically modified in vegetative structures, has undergone the same reproductive development as the other recent members of the family, especially *Dacrydium*, and that it is closely related to them.

Podocarpus dacrydioides, *Dacrydium*, *Microcachrys*, *Pherosphaera*, and *Phyllocladus* are with a very few exceptions Australasian, and seem to represent two or three main lines of evolution from the genus *Podocarpus*. They show a tendency towards extreme leaf-reduction, a modification or loss of the pollen wings, an erect position of the ovule, freedom of the nucellus, freedom and reduction of the epimatium, and fusion of the epimatium with the bract.

Very close to these forms stands the Chilean *Saxegothea*, which differs from them in having much larger leaves and a more complex vascular system in the bract, where the single strand has divided into three or five (Diagram 8, E). This dominance of the bract is precisely comparable to a similar state of affairs in *Cunninghamia*, *Athrotaxis* (Diagram 8, F), and the Araucarineae. A small free epimatium and the absence of winged pollen also show the advanced position of the genus.

The classification of the Podocarpineae here proposed (Diagram 9) is radically different from that recently put forward by Stiles, which is based on the hypothesis we have already discussed, that *Saxegothea* and *Pherosphaera* are the most primitive members of the family. The complete separation by this author of § *Dacrycarpus*, *Microcachrys*, and *Pherosphaera*

from one another is not in agreement with the many similarities between these three groups, and *Podocarpus* and *Dacrydium*, though placed wide apart in the scheme, seem in reality closely related to one another through such species as *Dacrydium Bidwillii*. The derivation of *Phyllocladus* from *Pherosphaera* is surely a novel solution to the vexed question of the relationship of this interesting genus.

The Podocarpineae, therefore, on the hypothesis set forth in the present paper, form a well-marked natural alliance. A fairly close series connects

DIAGRAM 9. Illustrating the phylogeny of the Podocarpineae.

the various members with one another and their interrelationships are clearly explicable if we consider the genus *Podocarpus*, standing at one end of the line, as the most primitive type. If the series is reversed, however, and the whole family is derived from the Araucarineae through forms like *Saxe-gothea*, the close resemblance between *Podocarpus* and the *Abietineae* must either be regarded as a most amazing case of parallel development, or else the affinity of these two groups must be recognized, and all other Conifers as well be derived from the Araucarineae. This view denies the brachyblast character of the ovuliferous scale by considering the double-scaled forms to have arisen from single-scaled ones, and also disregards all the

evidence which points to the fact that the Abietineae and not the Araucarineae are the most ancient Conifers.

The close series of forms from *Podocarpus* to *Saxegothea* is very suggestive as offering a key to the evolutionary development of the modern Araucarineae. If the epimatium suffered progressive reduction and final elimination and the bract were thus strongly dominant, as has happened in the line culminating in *Pherosphaera*, the resulting cone scale, with a somewhat amplified vascular system as in *Saxegothea*, would much resemble that of a miniature *Agathis*. Should the adnate sterile bract of such a form as *Podocarpus dacrydioides* become much more strongly developed in proportion to the size of the ovule, as in *Cunninghamia* and *Saxegothea*, and if the apical knob of the epimatium were drawn out into a ligule-like structure, a state of affairs would result precisely comparable to that in *Araucaria*. Steps are therefore easy to imagine by which the living Araucarineae have arisen from an ancient 'double-scaled' group, close to the primitive Podocarpaceae, and thus ancestral to both families, but which had only recently sprung from the ancient Abietineae.

(b) Relationship to Taxineae.

The relationship between the Podocarpaceae and the Taxineae has also been a subject of controversy, but from the prevailing reduction of the female cone to a single ovule in both families and the common possession of a 'second integument' and a fleshy fruit, they have usually been placed near each other and included together as the Taxineae.

Cephalotaxus, by the strong development of its ovulate strobilus, the typically coniferous structure of its female gametophyte, and the simple condition of its microsporophyll, which, instead of bearing numerous sporangia, has but two or three, is clearly shown to be more primitive than *Torreya* or *Taxus*, the other members of the Taxineae. It also displays a notable similarity in strobilar anatomy, gametophytic history, and embryonic development, to certain species of the Podocarpaceae.

The female cone is a short axis with a number of bracts, in the axil of each of which are borne two erect ovules (Diagram 7, C). The vascular system precisely resembles that of the Podocarpaceae and Abietineae, for the cylinder of bundles gives off one strand to each bract and two more, from the sides of the gap, and becoming inversely oriented, to the ovuliferous structures, one to each ovule (Diagram 7, D). The gross anatomy of the cone closely resembles that of *Phyllocladus*.

In the structure of the ovule, also, *Cephalotaxus* shows a strong resemblance to *Podocarpus*, and especially to its sub-genus *Stachycarpus*. The epimatium in both instances is full of canals, ripens into a fleshy coat, and is entirely fused with the integument, which becomes stony at maturity. Attention was called to the fact that in *Podocarpus spicatus* two bundles,

wide apart, run down through the epimatium on its adaxial face. In *P. ferrugineus* these two bundles, after leaving the chalazal end, pass down exactly opposite one another, one on either side of the ovule and each just at the inner margin of the epimatium. This latter condition, obviously less primitive than that in *P. spicatus*, is precisely what we find in the ovule of *Cephalotaxus* (and of *Torreya* as well), where two opposite and lateral bundles pass upward from the chalaza towards the micropyle just at the limit of the two integuments. In *Podocarpus*, of course, there are the two dorsal strands which represent the vascular system of the ovuliferous scale between the axis and the attachment of the ovule, but if we imagine that this part of the scale, after being greatly shortened, as in *Dacrydium*, has disappeared entirely, as in *Phyllocladus*, and that the ovule has become erect while at the same time retaining its integumentary and vascular structures, we obtain a close approximation to conditions in *Cephalotaxus*. A study of the little known New Caledonian genus *Acropyle* should prove very interesting in this connexion, for here the cone has been reduced to a fleshy receptaculum bearing a single erect ovule which has an inner stony and, apparently, an outer leathery integument. This seems to provide an intermediate condition between *Podocarpus* and the Taxineae.

The resemblance between *Cephalotaxus* and the sub-genus *Stachycarpus* of *Podocarpus* is further emphasized by the structure of the female gametophyte and embryo, for in both the reproductive cycle extends through two seasons; spongy tissue is almost completely absent; the archegonia are only two or three in number and are very long and narrow; they are eventually deeply sunken in the endosperm; the neck is composed of but one tier of cells; the first two sporophytic mitoses occur in the upper portion of the archegonium, and wall formation occurs at the sixteen-nuclear stage, and is followed by rapid and irregular cell-division which gives rise to a proembryo different from that of almost all other Conifers. It is composed of four tiers: a single terminal cell, more or less protective in function; a group of cells immediately behind this, which is to give rise to the embryo; a suspensor tier, and a poorly developed rosette.

In opposition to this evidence from sporophytic anatomy and the female gametophyte, it may be urged that the occurrence in *Cephalotaxus* of two ovules to a bract instead of one is a strong objection to the derivation of this genus from forms close to *Podocarpus*. The other two members of the Taxineae have but one ovule to a bract, and although the occurrence of a single ovule is constant throughout the Podocarpaceae (two nucelli were found in one epimatium of *P. spicatus*), the number varies greatly in other coniferous families. It seems very probable that the multiplication of sporangia which is so evident in the male cone of the Taxineae may have affected the female strobilus as well. The occurrence of two megasporangia to each megasporophyll in *Ginkgo* and the Abietineae makes it also very

probable that this is the primitive state of affairs. To this the biovulate condition of *Cephalotaxus* may conceivably be a reversion.

Stachyotaxus, as described by Nathorst (7) from the Rhaetic, is worth mentioning here, though it probably shows nothing more than an interesting analogy. Its foliage is distinctly taxineous in habit, and the cones, long and lax like those of *P. spicatus*, are composed of small bracts each of which bears near the end of its lamina two erect ovules, both apparently invested by a basal cup-like 'epimatium'. This fossil, apparently a combination of *Podocarpus*, *Dacrydium*, and *Cephalotaxus*, might easily be regarded as an ancestral form for the Taxineae, but its ancient geological horizon, the absence of all information save that from impressions, and our slight acquaintance even with these, make quite uncertain any conclusions as to its affinities. *Stachyotaxus* at least demonstrates the antiquity of tendencies exhibited by the living Taxineae.

The resemblance in reproductive structures between *Cephalotaxus* and the Podocarpaceae are sufficiently numerous and important to warrant our assuming a rather close relationship between them, and since *Cephalotaxus* is the most primitive member of its family, we are justified in considering the Taxineae as a whole to have arisen from somewhere among the ancient members of the Podocarpaceae.

Further investigations, both anatomical and embryological, upon many other species of the Podocarpaceae are necessary before there will be at hand a body of facts sufficiently comprehensive to enable us to construct with certainty a true natural classification for the whole family. Information obtained by the writer from a general study of the anatomy of the vegetative organs of the Podocarpaceae emphatically confirms the conclusions based on the present investigation of their reproductive structures and furnishes further evidence for the affinity of *Podocarpus* with the Abietineae on the one hand and with the Taxineae on the other.

We must recognize that in the families under consideration, as in so many other groups of plants, very similar structures have been developed along quite distinct lines of evolution, and we must guard against accepting this similarity as indubitable evidence of identical origin. The whole body of facts derived from all possible sources, rather than a few arbitrarily set up as unfailing criteria, must be considered if we are to arrive at sound phylogenetic conclusions.

SUMMARY.

1. The male strobilus of the Podocarpaceae is essentially uniform in structure throughout the family, and consists, as in the Abietineae, of spirally arranged bisporangiate sporophylls.

2. The female strobilus shows a tendency to become fleshy in whole or part at maturity. Reduction in number of sporophylls has occurred

throughout, though this reduction has not been uniformly progressive, but subject to great variation in degree within the same group.

3. In typical *Podocarpus* the free bract is provided with a single bundle from the vascular cylinder of the cone and subtends the epimatium, an axillary structure exactly homologous with the ovuliferous scale, and which is supplied by two bundles with inverse orientation, each arising from one side of the bract gap. These eventually send two small strands into the chalaza. The ovule is inverted and is completely enveloped by the epimatium. The integument is adnate to the epimatium without and to the nucellus within.

4. From this condition, closely resembling that of the Abietineae, there is an almost complete series through *Podocarpus dacrydioides* and *Dacrydium* to *Microcachrys*, *Pherosphaera*, *Saxegothea*, and *Phyllocladus*, in which are exhibited the following tendencies: the bract becomes fused to the epimatium and its single bundle may divide into a series of strands; the vascular system of the scale is reduced to little but the ovular supply, and its two bundles become fused for some distance to that of the bract, from the sides of which they eventually depart and bend upward adaxially, assuming an inverse orientation; the ovule swings from an inverted to an almost erect position; the epimatium becomes much reduced in proportion to the bract, and, instead of enclosing the ovule entirely, degenerates into a mere basal sheath or quite disappears; the nucellus, integument, and epimatium become entirely free from one another.

5. The male gametophyte is characterized by the presence of two primitive prothallial cells, which give rise subsequently to more or less vegetative tissue. At germination all the cells are represented by free nuclei save the body cell, the nucleus of which divides into the two male nuclei, one naked and the other surrounded by a protoplasmic body.

6. In typical *Podocarpus* there are two well-developed wings on the pollen-grain, but in its sub-genus *Dacrycarpus* there are three or four, a condition which also obtains in *Microcachrys*. In *Dacrydium* the young spore has always two large wings, but in certain species these are almost obliterated at maturity. The pollen of *Phyllocladus* has two much-reduced wings, but in that of *Saxegothea* there are none at all.

7. The length of the reproductive cycle is limited to a single season in § *Eupodocarpus*, § *Nageia*, § *Dacrycarpus*, *Dacrydium Bidwillii*, *Phyllocladus*, and apparently in *Saxegothea* and *Microcachrys*. It extends over two seasons in § *Stachycarpus* and in most of *Dacrydium*.

8. Spongy tissue is best developed in § *Eupodocarpus*, and is somewhat reduced or entirely absent in the other members of *Podocarpus* and *Dacrydium*. The germination of the spore and development of the embryo-sac occur as in most Conifers. A megaspore membrane is well developed everywhere save in § *Dacrycarpus*, where it is somewhat reduced. The

archegonia are apical. They vary in number from two or three in § *Stachycarpus* and *Dacrydium* to ten or more in certain species of *Podocarpus*, and are with rare exceptions quite distinct from one another and not grouped. The neck may be either a single-tiered rosette or an irregular group of cells. In § *Eupodocarpus*, and to some extent in *Dacrydium*, asteroids, or centres of radiating fibrils, occur in the young archegonium. A large basal vacuole is everywhere present, save in § *Stachycarpus*.

9. At fertilization the male nucleus is about half the size of the female, to which it becomes closely appressed and is finally fused. In all forms but § *Stachycarpus* only the first sporophytic mitosis takes place in the position of the fertilized egg, and the two resultant nuclei drop to the bottom and divide into sixteen. Walls are now laid down and a proembryo results of three tiers: a terminal binucleate embryogenous cell, a suspensor tier, and a poorly developed rosette. In § *Stachycarpus* the first two sporophytic mitoses occur in the upper portion of the archegonium, and the four nuclei, dropping to the base, give rise to a proembryo of four tiers: a single terminal cell more or less protective in function and often sloughed off; a group of from eight to forty cells which will form the embryo; a suspensor tier, and a rosette.

10. The endosperm cells in all species become multinucleate about the time of fertilization. The embryo remains intact except in *Dacrydium*, where it usually buds or divides. At maturity the embryo is dicotyledonous.

11. The close resemblance of *Podocarpus* to the Abietineae in the development of the male and female gametophytes and the embryo, as well as in the anatomy of the staminate and ovulate strobili, warrants the hypothesis that the Podocarpaceae have been derived from the Abietineae through forms somewhat resembling *Podocarpus*. Certain points of similarity between Podocarpaceae and Araucarineae suggest that these two families may have both arisen from an ancient group, closely abietineous in affinity. The advantages of this hypothesis are that it is in harmony with the view that the Abietineae are the most ancient Conifers; that it assigns a logical position to the Podocarpaceae; that it accounts for the wide variation in the genus *Podocarpus*; that it offers a reasonable explanation of the origin of the epimatium, and that it makes possible the construction of a natural classification for the Podocarpaceae on the assumption that the family represents three or four main lines of ascent from its most primitive genus, *Podocarpus*.

12. The striking resemblance in strobilar anatomy, in the structure of the female gametophyte, and in the development of the embryo, between *Stachycarpus*, a sub-genus of *Podocarpus*, and *Cephalotaxus*, the most primitive genus of the Taxineae, suggests that the latter family has arisen from some ancient member of the Podocarpaceae. The structure of the

living *Phyllocladus* and that of the fossil *Stachyotaxus* indicate possible steps by which the taxinean evolution may have taken place.

The writer wishes to thank most heartily his many friends in the several Botanic Gardens, in various other departments of the public service, and elsewhere throughout Australia and New Zealand for their active assistance in securing material, and he is especially indebted to Mr. T. F. Cheeseman, F.L.S., F.Z.S., of the Auckland Museum, for invaluable aid as to localities and identifications. This investigation was carried on in the Phanerogamic Laboratories of Harvard University.

BIBLIOGRAPHY.

1. BURLINGAME, L. L.: The Staminate Cone and Male Gametophyte of *Podocarpus*. Bot. Gazette, vol. xlvi, 1908, pp. 161-78.
2. COKER, W. C.: Notes on the Gametophytes and Embryo of *Podocarpus*. Bot. Gazette, vol. xxxiii, 1902, pp. 89-107.
3. EAMES, A. J.: The Morphology of *Agathis australis*. Ann. Bot., vol. xxvii, 1913.
4. GIBBS, L. S.: On the Development of the Female Strobilus in *Podocarpus*. Ann. Bot., vol. xxvi, 1912, pp. 515-73.
5. JEFFREY, E. C., and CHRYSLER, M. A.: The Microgametophyte of the Podocarpaceae. Amer. Nat., vol. xli, 1907, pp. 102-7.
6. KILDAHL, N. JOHANNA: The Morphology of *Phyllocladus alpinus*. Bot. Gazette, vol. xlvi, 1908, pp. 339-48.
7. NATHORST, A. S.: Über *Pallissya*, *Stachyotaxus* und *Palaeotaxus*. Kungl. Svenska Vetenskap. Handl., Band xliii, 1908, No. 8.
8. NORÉN, C. O.: Zur Kenntnis der Entwicklung von *Saxegothea conspiciua*. Svensk. Bot. Tidsskr., vol. ii, 1908, pp. 101-22.
9. RADAIS, M.: L'Anatomie comparée du fruit des Conifères. Ann. Sci. Nat., Bot., sér. 7, t. xix, 1894, pp. 165-368.
10. ROBERTSON, AGNES: Some Points in the Morphology of *Phyllocladus alpinus*. Ann. Bot., vol. xx, 1906, pp. 259-65.
11. STILES, W.: The Anatomy of *Saxegothea conspiciua*. New Phytologist, vol. vii, 1908, p. 209.
12. ——— (With BROOKS, F. T.): The Structure of *Podocarpus spinulosa*. Ann. Bot., vol. xxiv, 1910, pp. 305-18.
13. ———: The Podocarpeae. Ann. Bot., vol. xxvi, 1912, pp. 443-515.
14. STRASBURGER, E.: Die Coniferen und die Gnetaceen. 1872.
15. ———: Die Angiospermen und die Gymnospermen. 1879.
16. THOMSON, R. B.: On the Pollen of *Microcachrys tetragona*. Bot. Gazette, vol. xlvii, 1909, pp. 26-9.
17. ———: The Megasporophyll of *Saxegothea* and *Microcachrys*. Bot. Gazette, vol. xlvii, 1909, pp. 345-54.
18. TISON, A.: Sur le *Saxegothea*. Mém. Soc. Linn. Normandie, t. xxiii, 1909, pp. 139-60.
19. VAN TIEGHEM, P.: Anatomie comparée de la fleur femelle et du fruit des Cycadées, des Conifères, et des Gnétacées. Ann. Sci. Nat., Bot., sér. 5, t. x, 1869, pp. 269-304.
20. YOUNG, MARY S.: The Male Gametophyte of *Dacrydium*. Bot. Gazette, vol. xlv, 1907, pp. 189-96.
21. ———: The Morphology of the Podocarpaceae. Bot. Gazette, vol. 1, 1910, pp. 81-100.

DESCRIPTION OF FIGURES IN PLATES V-IX.

Illustrating Mr. Sinnott's paper on the Podocarpaceae.

PLATE V.

- Fig. 1. *Podocarpus Totara*. Archegonium at time of fertilization, showing fusion nucleus, second male nucleus, and prothallial nuclei.
 Fig. 2. *P. ferrugineus*. Mature archegonium and male cells, just before fertilization.
 Fig. 3. *Dacrydium Bidwillii*. Mature archegonium sealed over by endosperm.
 Fig. 4. *Podocarpus spicatus*. Germinating pollen.
 Fig. 5. *Dacrydium cupressinum*. Microspore.
 Fig. 6. *D. cupressinum*. Mature pollen-grain with reduced wings.

PLATE VI.

- Fig. 7. *Podocarpus Totara*. Longitudinal section of mature ovule. $\times 12$.
 Fig. 8. *P. nivalis*. Transverse section of mature ovule. $\times 15$.
 Fig. 9. *P. Totara*. Transverse section of receptaculum, with bract traces departing. $\times 15$.
 Fig. 10. *P. spinulosa*. Longitudinal section of young strobilus. $\times 15$.
 Fig. 11. *P. vitiensis*. Longitudinal section of young ovule. $\times 10$.
 Fig. 12. *P. dacrydioides*. Longitudinal section of mature ovule. $\times 12$.
 Fig. 13. *P. dacrydioides*. Longitudinal section of very young ovule, megaspore stage. $\times 20$.
 Fig. 14. *P. dacrydioides*. Longitudinal section of young ovule, free nuclear stage. $\times 15$.
 Fig. 15. *P. ferrugineus*. Longitudinal section of young ovule. $\times 8$.
 Fig. 16. *P. spicatus*. Longitudinal section of young ovule and part of cone axis. $\times 9$.
 Fig. 17. *P. ferrugineus*. Longitudinal section of very young ovule, just after spore germination. $\times 15$.
 Fig. 18. *P. ferrugineus*. Transverse section of ovule, showing vascular girdles around nucellus from lateral bundles. $\times 10$.

PLATE VII.

- Fig. 19. *P. spicatus*. Transverse section of cone axis, supply to bract and ovule departing at left. $\times 30$.
 Fig. 20. *P. ferrugineus*. Transverse section through very base of ovule, showing single trace to bract and two inverted bundles to epimatium. $\times 60$.
 Fig. 21. *Dacrydium Bidwillii*. Longitudinal section of ovule. $\times 15$.
 Fig. 22. *D. Bidwillii*. Transverse section through chalazal region, showing two bundles in epimatium and two departing from these into base of ovule. $\times 30$.
 Fig. 23. *D. intermedium*. Longitudinal section of young ovule, entirely surrounded by epimatium. $\times 35$.
 Fig. 24. *D. cupressinum*. Mature pollen-grain with reduced wings. $\times 700$.
 Fig. 25. *P. elatus*. Megaspore mother-cell. $\times 160$.
 Fig. 26. *P. Totara*. Small embryo-sac at free nuclear stage, surrounded by spongy tissue. $\times 60$.
 Fig. 27. *P. Totara*. Young archegonium with 'asteroids' in central cell. $\times 60$.
 Fig. 28. *P. Totara*. Mature archegonium. $\times 60$.
 Fig. 29. *P. Totara*. Mature archegonium with persistent ventral canal nucleus. $\times 140$.
 Fig. 30. *P. Totara*. Proembryo of four nuclei. $\times 160$.
 Fig. 31. *P. nivalis*. Sixteen-celled proembryo with protective cellulose cap, just after wall formation. $\times 160$.

PLATE VIII.

- Fig. 32. *P. Totara*. Embryos from two archegonia, with suspensors just beginning to lengthen. $\times 120$.

- Fig. 33. *P. Hallii*. Nearly mature embryo. $\times 30$.
 Fig. 34. *P. dacrydioides*. Mature archegonium. $\times 60$.
 Fig. 35. *P. dacrydioides*. Mature archegonium with persistent ventral canal nucleus. $\times 125$.
 Fig. 36. *P. dacrydioides*. First sporophytic mitosis. $\times 60$.
 Fig. 37. *P. dacrydioides*. Fusion nucleus. $\times 160$.
 Fig. 38. *P. dacrydioides*. Portion of endosperm, showing mucilage cells and young embryo.
 $\times 20$.
 Fig. 39. *P. ferrugineus*. Functional megaspore with two abortive ones at end. $\times 160$.
 Fig. 40. *P. ferrugineus*. Two archegonia, one with proembryo at base, the other unfertilized.
 $\times 30$.
 Fig. 41. *P. spicatus*. Apex of archegonium, with egg nucleus and ventral canal nucleus. $\times 160$.
 Fig. 42. *P. spicatus*. First sporophytic mitosis. $\times 160$.
 Fig. 43. *P. ferrugineus*. First four nuclei of proembryo in upper part of archegonium, surrounded by cytoplasmic sheath. $\times 160$.
 Fig. 44. *P. spicatus*. Archegonium with first four nuclei of proembryo dropping towards the base, and vacuole forming above. $\times 30$.

PLATE IX.

- Fig. 45. *P. spicatus*. Proembryo of sixteen nuclei at base of archegonium. $\times 160$.
 Fig. 46. *P. spicatus*. Proembryo soon after wall formation, with suspensors, embryo, and apical cell. $\times 125$.
 Fig. 47. *P. ferrugineus*. Two young embryos. $\times 160$.
 Fig. 48. *Dacrydium cupressinum*. Two young archegonia. $\times 60$.
 Fig. 49. *D. cupressinum*. Young embryo beginning to divide. $\times 160$.
 Fig. 50. *Podocarpus dacrydioides*. Male cells and prothallial nuclei in pollen-tube. $\times 160$.

3.

2.

1.

4.

5.

6.

7.

8.

9.

10

11.

12.

13.

14.

15.

16.

17.

18.

19.

20.

21.

22.

23.

24.

25.

26.

27.

28.

29.

30.

31.

32

33.

34

35.

36.

37.

38.

39.

40.

41

42.

43.

44.

45.

46.

47.

48.

49.

50.

Sinnott, Edmund W. 1913. "The morphology of the reproductive structures in the Podocarpaceae." *Annals of botany* 27, 39–82.

<https://doi.org/10.1093/oxfordjournals.aob.a089452>.

View This Item Online: <https://www.biodiversitylibrary.org/item/237410>

DOI: <https://doi.org/10.1093/oxfordjournals.aob.a089452>

Permalink: <https://www.biodiversitylibrary.org/partpdf/319956>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.