ARNOLD ARBORETUM
HARVARD UNIVERSITY LIBRARY
NEW YORK
BOTANICAL
GARDEN.BULLETIN
OF
POPULAR INFORMATION

JAMAICA PLAIN, MASS.

JULY 16, 1915

Cornus Amomum, the Silky Cornel, which has been much used in the Arboretum, has been in flower here for several days. In cultivation it is not a satisfactory plant unless it can be given sufficient room for its wide-spreading branches to extend freely and spread over the ground. When crowded by other plants the branches become erect and it loses its real beauty and value. To be seen at its best this Cornel should have a clear space with a diameter of not less than twenty feet in which to spread. It is well suited for the front of groups of trees and shrubs, and there is no better shrub to plant by the margins of ponds and streams where its long branches can hang gracefully over the water. Its purple stems are attractive in winter, and the bright blue fruits which ripen in the autumn add to the value of this native shrub. In the Cornel Group, at the junction of the Meadow and Bussey Hill Roads, there is a good specimen of this plant, and its value for planting near water can be seen on the borders of the small pond in the rear of the Cornel Group.

Cornus asperifolia. This Cornel flowers a week or ten days later than the Silky Cornel. It is a widely distributed plant from western New England to Texas, and under favorable conditions becomes a tree occasionally forty feet high. At the north, however, it is a tall, broad shrub with erect and slightly spreading branches. The flower-clusters are small but are produced in great profusion and are followed by white fruits. There is a large specimen among the other Cornels near the junction of the Meadow and Bussey Hill Roads.

Cornus paucinervis. It was hardly to have been expected that this Chinese shrub could flourish in New England for it grows naturally not much above the sea-level central China in a climate where the

Orange thrives, and rarely ascends to elevations of three thousand feet. It is a shrub five or six feet tall with erect stems, small, narrow pointed leaves with only two or three pairs of prominent veins, small clusters of white flowers and black fruits. There is a good specimen of this plant among the Chinese shrubs on the southern slope of Bussey Hill which in a few days will be covered with flowers. Shrubs which flower here late in July are not very common and this Cornel promises to be a useful addition to the list.

Late flowering Barberries. Three species of *Berberis* from western China are now in flower, *B. aggregata*, *B. Pratii*, and *B. subcaulialata*. These plants will probably become popular for they are the latest of the Barberries to flower. They are all erect-growing, tall shrubs with small yellow flowers in drooping clusters which are followed by red fruits. There are plants in the Shrub Collection and with the Chinese shrubs on the southern slope of Bussey Hill.

Hydrangea radiata. A form of *Hydrangea arborescens* (var. *grandiflora*), with large globose heads of sterile flowers, has become immensely popular in this country since its discovery a few years ago in one of the western states, and it can now be seen in many suburban gardens. A much more beautiful American species, however, is *Hydrangea radiata*, which is now in flower in the Shrub Collection. It is a native of mountain slopes in North and South Carolina, and is a round-topped shrub with large leaves very dark green above and silvery white below, and broad heads of flowers surrounded by a ring of white neutral flowers. It is one of the handsomest of all the Hydrangeas which are perfectly hardy in this climate, and although once a popular garden plant it is now rarely found in collections.

Hydrangea paniculata. More conspicuous now in the collection is the early-flowering form of *Hydrangea paniculata* (var. *praecox*). The most generally planted form of *Hydrangea paniculata* is that in which all the flowers are sterile, known as *Hydrangea paniculata grandiflora*. This plant produces large clusters of white flowers which turn rose color in fading, and will not be in bloom for several weeks. The variety *praecox*, which is one of the forms of the wild plant, has ray flowers surrounding the clusters of sterile flowers. There are two or three forms of the variety *praecox* in the collection differing in the size of the flower-clusters and in the size of the ray flowers. The handsomest and earliest of these was raised from seeds collected by Professor Sargent in Hokkaido where it grows into a small tree sometimes twenty or thirty feet tall.

Colutea arborescens. This and related species are now among the most beautiful plants in the Shrub Collection as they are still covered with yellow flowers which are mixed with the large, inflated, rose-colored or pink pods to which these plants owe their common name of Bladder Sennas.

Aesculus parviflora. The last of the Horsechestnuts to flower, *Aesculus parviflora*, will soon be in bloom. It is a tall, round-topped, shapely shrub well suited to plant in large masses or as a single specimen. In good soil and when uncrowded by other plants it soon spreads

1915. "Cornus paucinervis." *Bulletin of popular information - Arnold Arboretum, Harvard University* 1(13), 49–50. <https://doi.org/10.5962/p.320399>.

View This Item Online: <https://www.biodiversitylibrary.org/item/226132>

DOI: <https://doi.org/10.5962/p.320399>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320399>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

IMLS LG-70-15-0138-15

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.