for fully six weeks. *Gordonia* is related to the *Camellia*, and the pure white flowers which vary from three to three and a half inches in diameter, although more cup-shaped, resemble single *Camellia* flowers. This small tree was discovered in 1765 near Fort Barrington on the Altamaha River; it has entirely disappeared, however, as a wild plant, and it has only been preserved by the specimens cultivated chiefly in the neighborhood of Philadelphia where it was sent by its discoverers. There are well established plants on Hickory Path near Centre Street and on Azalea Path, and in these sheltered positions the plants are doing well and have not suffered at all in recent severe winters. This *Gordonia*, however, grows more rapidly and to a larger size in the middle states and there are many good specimens in Pennsylvania gardens.

Abelia grandiflora on Hickory Path near Centre Street is still well covered with flowers. These resemble in shape the flowers of some of the Honeysuckles; they are white faintly tinged with rose color, and their delicate beauty is set off by the small, dark green and lustrous leaves. *Abelia grandiflora* is a slender shrub with arching stems from three to four feet high and is thought to be a hybrid between two Chinese species. Until the introduction by the Arboretum of some of the species of this genus from western China it was believed to be the hardiest of the *Abelias*. In the Arboretum it suffers in severe winters but in sheltered positions it flowers well every year and the flowers continue to open during nearly two months. This *Abelia* has become an exceedingly popular plant in the gardens of the southern states and is cultivated with more or less success as far north as New York.

The Chinese Buddleias. The flowers are still in good condition on the different forms of *Buddleia Davidii* or, to use its more common name, *B. variabilis*. This plant has one-sided, pointed, many-flowered clusters which curve downward from arching stems and are thickly covered with small, blue-purple, fragrant flowers. In some of the forms of this plant are found perhaps the most beautiful of all summer and autumn flowering shrubs, and although only recently brought to the United States and Europe by Wilson they are already largely planted in this country where they have received the name of Summer Lilacs. Here at the north *Buddleia Davidii* is not perfectly hardy, and the stems are killed to the ground by cold, but new stems spring up and as the flower-clusters are produced at the ends of branches of the year this severe pruning improves the flowers. Few plants in their season are better suited to supply cut flowers, and for this purpose as well as for garden decoration this *Buddleia* in its various forms has proved one of the most useful shrubs of recent discovery.

The period for flowers in the Arboretum. It has been shown by the Arboretum that flowers can be found here in Massachusetts on trees and shrubs growing in the open ground during every month of the year with the exception of December. There will still be flowers on the native Witch Hazel in November and before the end of January the flowers will be open on the Witch Hazel of southern Missouri and Arkansas, *Hamamelis vernalis*. These will soon be followed by the

flowers of the Japanese and Chinese Witch Hazels which last well through February and do not suffer from the severest cold eastern Massachusetts ever has to endure. Then before the end of March the flowers of some of the Willows open and begin the long procession which only ends in November. The introduction of the Asiatic Witch Hazels has added greatly to the interest of northern gardens in winter and they are bright and cheerful winter companions. They might, therefore, well find a place near every country home and in small city yards. The flowers of the Chinese species, *Hamamelis mollis*, are larger and of a brighter yellow color than the flowers of the other Witch Hazels and this promises to be the best worth cultivation of them all. It is a perfectly hardy vigorous shrub, grows rapidly, and begins to flower when only a few feet high. Specimens of all the species of *Hamamelis* are planted in the group of these plants on the Meadow Road, and the best specimen of *Hamamelis mollis* is on Hickory Path near Centre Street.

Evonymus alatus. The leaves of many of the trees and shrubs of eastern Asia turn to as brilliant colors in the autumn as those of the related species of eastern North America. Usually, however, the leaves of the Asiatic species change color later than those of the American species, and in Japan the best color effects are in November or a month later than here. There are, of course, many exceptions to this general rule. The leaves of *Evonymus alatus*, for example, were brilliant in the Arboretum ten days ago and are already beginning to fall. This is a shapely shrub six or eight feet tall and ten or twelve feet broad, distinguished by the corky wings of the branches. The flowers, as in all the species of this genus, are not conspicuous and the fruit is smaller and less showy than that of many of these plants. The great value of this hardy Japanese Burning Bush is therefore found in the autumn coloring of the leaves, which assume a deep rose color of exquisite beauty and unlike that produced by any other hardy plant in cultivation.

Acer ginnala. The leaves of this Maple also turn and fall by the middle of October, and equal or surpass in their autumn scarlet those of any American plant. *Acer ginnala* is a small shrubby tree with deeply dentated leaves, sometimes thirty feet high, and very common along forest borders near Vladivostok and in other parts of eastern Siberia. The flowers are produced in rather compact clusters and, unlike those of other Maples, are distinctly fragrant. This Maple is one of the first Siberian trees introduced by the Arboretum and it is now gradually finding its way into general cultivation in this country.

Nyssa sylvatica. There is perhaps no more beautiful object this week in the Arboretum than the group of these trees variously known as Sour Gum, Tupelo and Pepperidge. The scarlet and orange colors of the leaves of the Sour Gum in October are probably not surpassed by those of any other American tree and their beauty is increased by the lustre of the leaves which adds to their autumn brilliancy. The Tupelo is a common and widely distributed tree, occurring from Maine to Florida, Missouri and Texas. At the north, especially near the coast, it is usually found near the borders of swamps and ponds, and is a low,

1915. "The period for flowers in the Arboretum." *Bulletin of popular information - Arnold Arboretum, Harvard University* 1(15), 58-59.

<https://doi.org/10.5962/p.320429>.

View This Item Online: <https://www.biodiversitylibrary.org/item/226132>

DOI: <https://doi.org/10.5962/p.320429>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320429>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

IMLS LG-70-15-0138-15

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.