fectly hardy here but it blooms so early that in about nine years out of ten the flowers after they open are spoiled by frost. This year not a flower-bud has been injured and the plants are now in flower.

Forsythias. Many flower-buds of these plants have been killed as they were two years ago. All the flower-buds in the lower part of the great Forsythia bed on the slope above the Bussey Hill Road are entirely killed, while on the plants in the upper part of this bed, although many buds have suffered, there are open flowers enough to make this part of the bed conspicuous. In the lower ground of the general Shrub Collection the flower-buds of all the species and varieties of Forsythia have been entirely killed with the exception of those of the Albanian *F. europaea* which are uninjured. If the flower-buds of this species are better able to support cold than those of the other species, which are all natives of eastern continental Asia, the European plant should be better known and more generally cultivated. One of the last plants discovered in Europe, it is a vigorous, fast-growing shrub with more erect stems and branches than are usually found in the other species of this genus. The flowers are not quite so showy as those of *F. suspensa* and its hybrids and varieties, but they are always abundantly produced and of good color. This plant is still rare in gardens and it is doubtful if it can be found in American nurseries.

Magnolias. The Japanese shrubby species, *Magnolia stellata*, is usually the first of the Magnolias to flower in the Arboretum and the petals are sometimes browned by a late frost. This year many of the flower-buds have been killed and only a few flowers much reduced in size are now open on these beautiful and usually satisfactory plants. Some flowers are now open on the northern form of the Japanese *M. kobus* (var. *borealis*). This is a shapely tree, with small drooping flowers which open before the appearance of the leaves, and good foliage, but it never flowers here very freely and this year there are fewer flowers than usual. As a flowering tree for this climate it is inferior to the white-flowered Chinese species, *M. conspicua*, or as it now to be called, *M. denudata*, and its hybrids, which also flower before the appearance of the leaves and are not yet in flower.

Witch Hazels. The earliest of the Witch Hazels, *H. vernalis*, a native of southern Missouri and of Arkansas, opened its first flowers during the last week of December and these were followed in January and February by the flowers of the Chinese and Japanese species which were not affected by a temperature of several degrees below zero. In the size and color of its flowers and in foliage the Chinese *H. mollis* is the handsomest of these winter-flowering shrubs. It is hardy, fast-growing and free-flowering, and might well find a place in every garden where it can be seen from the windows of country or city houses. Unfortunately this shrub is still difficult to obtain from American nurseries.

The Cornelian Cherry (*Cornus mas*) is one of the earliest trees or arborescent shrubs to flower here. The flowers are light yellow and

are borne in clusters in the axils of the unfolding leaves, and although individually small are produced in such numbers that the branches are covered with them. They are followed by bright red, lustrous, oblong fruit the size of a small olive. The flower-buds and the flowers of this little tree are not injured by cold. The habit of this plant is good. The foliage is dark green and abundant and the fruit, although somewhat hidden by the leaves, is handsome. The Cornelian Cherry is a native of most European countries and of western Siberia, and has been an inhabitant of gardens for more than three hundred years. In the United States it was probably more generally planted in the first half of the last century than it is at present, although there are not many early flowering trees hardy in this climate which are better worth a place in the garden. In the Arboretum it can be seen in the Cornel Group near the junction of the Meadow and the Bussey Hill Roads.

Two Japanese plants. The leafless branches of *Euptelea polyandra* are now covered with flowers. The conspicuous part of these is the large bright orange-red anthers which hang on short filaments in axillary clusters from one end of the branches to the other. For its peculiar and early flowers this plant is well worth a place in New England gardens, and the large green leaves are handsome and abundant. There are several fine specimens of this large shrub or small tree on the lower side of Azalea Path and this week they are well worth examination. Another species, *E. Franchetii*, introduced by Wilson from western China, appears to be perfectly hardy in the Arboretum. The plants are still too small, however, to flower. The related *Cercidiphyllum japonicum*, which is the largest deciduous-leaved tree in Japan and now becoming common in American collections, is also beginning to flower. The flowers are inconspicuous but the red color of the unfolding leaves makes the tree attractive at this season. There is a group of this tree on the two sides of the Meadow Road not far beyond the Administration Building.

Corylopsis. This is a genus of the Witch Hazel Family, distributed, with several species, from the western Himalayas through western and central China to Japan. These plants have leaves which in general appearance resemble those of the Witch Hazel, and drooping spikes of fragrant yellow flowers which appear before the leaves, and terminate on what, later in the season, become short leafy branchlets. Several of these plants are hardy in this climate, but the flower-buds are often killed by intense cold, or, if they are not killed, the flowers open so early that they are destroyed by late frosts. This year the flower-buds have not been much injured, and the flowers of three species are now open. One of the best known of these plants is the Japanese *C. spicata*. It is a shrub with spreading branches four or five feet high, with yellow flowers, about half an inch long, and produced in from six to twelve-flowered spikes. There are plants of this shrub on Hickory Path near Centre Street. In the collection of Chinese shrubs on the southern slope of Bussey Hill two distinct and beautiful species introduced

1916. "The Cornelian Cherry." *Bulletin of popular information - Arnold Arboretum, Harvard University* 2(1), 2-3. <https://doi.org/10.5962/p.320461>.

View This Item Online: <https://www.biodiversitylibrary.org/item/216951>

DOI: <https://doi.org/10.5962/p.320461>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320461>

Holding Institution

Harvard University Botany Libraries

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.