western China. If they prove permanently successful here this *Daphne* will be a delightful addition to our early-flowering dwarf shrubs.

Crataegus Arnoldiana. This Hawthorn was never fuller of flower-buds than it is this spring and these will soon be open. This tree was discovered growing naturally on a wooded bank in the Arboretum; it grows also on the banks of the Mystic River in West Medford, Massachusetts, and near New London, Connecticut. It belongs to the Molles group of Hawthorns, which are trees distinguished by their large size, by their large early flowers which usually open with the unfolding of the leaves, and by the large, often edible, scarlet or rarely yellow fruits. There are several species found from the valley of the St. Lawrence River in the Province of Quebec to Texas. The species are, however, most numerous in the region west of the Mississippi River, and are almost entirely wanting in the southeastern states. *C. Arnoldiana* is one of the handsomest of the species of this group which is hardy here. The brilliant red fruit ripens late in August and falls in September earlier than that of the other species. In winter this tree is easily recognized by its upright growing, distinctly zigzag branches which are more thickly covered with spines than those of many of the related species. The largest trees of this Thorn can be seen on the left-hand side of the Valley Road just inside the Centre Street Gate and in front of the Platanus Collection. There are also several of these trees in front of the group of White Oaks, also on the left-hand side of the Valley Road. At the South Street entrance there are large plants of three other species of the Molles Group, *C. mollis* from the Ohio-Illinois region, *C. arkansana* from central Arkansas, and *C. submollis*, a Canadian and New England tree. These will all be in flower in a few days.

Malus Sieboldii, var. calocarpa. This Japanese Crabapple is one of the handsomest in the Arboretum both in spring and autumn. It is a broad tree-like shrub or small tree with only slightly lobed leaves, pink and white flowers fully an inch in diameter and brilliant scarlet lustrous fruits which are half an inch in diameter and more beautiful perhaps than those of any other Crabapple. This beautiful plant was raised from seed presented to the Arboretum in 1890 by Dr. William Sturgis Bigelow of Boston. It can be seen in the front row of the Crabapple Collection at the eastern base of Hemlock Hill. It has been found that *Sieboldii* is the oldest and therefore the proper name for the Chinese and Japanese Crabapple which up to this time has been known in gardens as *Malus* or *Pyrus toringo*.

It is interesting to note that Wilson found in western China the wild, single-flowered form of the beautiful Crabapple with rose-colored, semi-double flowers, *Malus Halliana*, or the Parkman Crab, which first came to this country from Japan, and was long believed to be a Japanese species. It is the Kaido of Japanese gardens. To the double-flowered form, which is the one generally cultivated in eastern gardens, the name *Malus Halliana Parkmanii* has been given; it is flowering well in the Arboretum this year.

1916. "Crataegus Arnoldiana." *Bulletin of popular information - Arnold Arboretum, Harvard University* 2(4), 16–16. <https://doi.org/10.5962/p.320490>.

View This Item Online: <https://www.biodiversitylibrary.org/item/216951>

DOI: <https://doi.org/10.5962/p.320490>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320490>

Holding Institution

Harvard University Botany Libraries

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.