Syringa Sweginzowii is flowering well here this year as it has for two or three years. The leaves are dark dull green and sharply pointed, and the flowers are borne in long narrow clusters; they are delicately fragrant, half an inch long, with very slender corolla-tubes, and are flesh-colored in the bud, becoming nearly white after the buds open. This species flowers freely even as a small plant and is well worth a place in a collection of Lilacs.

Syringa yunnanensis, although it is a native of southwestern China, is quite hardy in the Arboretum where it is flowering now for the fourth year. It is related, like most of the species of western China, to *S. villosa* and is a tall shrub of open habit with glabrous leaves pale on the lower surface and long narrow clusters of light flesh-colored or pink flowers. Geographically interesting, this plant is probably of less value as a garden plant than *S. Sweginzowii*.

The Tree Lilacs. As the flowers of the late-flowering group of Lilacs fade the earliest flowers of the so-called Tree Lilacs begin to open. There are three of these Lilacs which all bear large clusters of white or yellowish white flowers with a corolla shorter than the stamens, while in other Lilacs the corolla is longer than the stamens which are hidden in its throat. The flowers of the Tree Lilacs all have the disagreeable odor of the flowers of the Privet, and the leaves fall in the autumn without change of color. The first of these plants to flower, *S. amurensis*, a native of eastern Siberia as its name implies, is a shrub in habit twelve or fifteen feet high with dark close bark, broad thick leaves dark green above and pale below, and short, broad, unsymmetrical flower-clusters. *S. pekinensis* from northern China flowers next. This is also shrubby in habit, sometimes twenty or thirty feet tall and broad, with stout, spreading stems covered with yellow-brown bark separating readily into thin plates like that of some of the Birch-trees, dark green, narrow, pointed leaves and short and unsymmetrical flower-clusters usually in pairs at the ends of the branches. This species holds its leaves later in the autumn than the others, and produces great quantities of flowers every year, the other species usually flowering abundantly only every other year.

The last of the Tree Lilacs to flower, *S. japonica*, is a native of northern Japan, and is really a tree sometimes forty feet high with a tall straight trunk covered with lustrous brown bark like the bark of a Cherry-tree, a round-topped head of upright branches, broad, thick, dark green leaves, and erect, mostly symmetrical flower-clusters from twelve to eighteen inches long. This is one of the handsomest of the small trees which bloom here at the end of June or early in July.

Cornus Kousa is a small tree which in eastern Asia enlivens the forests as *Cornus florida* enlivens the forests of eastern North America, and *Cornus Nuttallii* those of our Pacific states. The three species have the large white or creamy white bracts under the flower-clusters which make the inflorescence so conspicuous, but the Asiatic tree differs from the American trees by the union of the fruits into a globular fleshy head, while the fruits of the American trees are not united together. *Cornus Kousa* is a small tree rarely exceeding twenty feet in height, and the floral bracts are narrower, more pointed and not as pure white as those of the American trees. It is valuable, however,

because it flowers three or four weeks later than *C. florida*. *C. Kousa* is a native of central Japan and was found in western China by Wilson. The Japanese and Chinese plants are both now in flower, the former on Hickory Path near Centre Street and the latter with the Chinese shrubs on the southern slope of Bussey Hill. The floral bracts of the Chinese plant in the Arboretum overlap and are broader and less pointed than those of the Japanese form, and the inflorescence, which is three inches across,, is much handsomer than that of the Japanese plant. The handsomest form of the Japanese tree which has been seen in this neighborhood is growing in Mount Auburn Cemetery where it has become an object of much beauty and interest.

Indigofera Potaninii has been raised at the Arboretum from seeds collected in the Province of Kansu by William Purdom during his collecting trip in northern China for the Arboretum. It is a beautiful little shrub now in bloom for the second year in the collection of Chinese shrubs on the southern slope of Bussey Hill. As it grows here it is three to four feet high, with a single stem and slender erect branches. The flowers are bright rose color, half an inch long, in long-stalked, erect and spreading racemes from two to three inches in length, from the axils of leaves on branches of the current year, and as the branches lengthen new flower-clusters appear and the plant remains in bloom for a long time. The flowers are of the same color but are larger than those of *Indigofera amblyantha* which Wilson found in western China and which until *I. Potaninii* bloomed was considered the handsomest shrubby species which could be grown here.

Cornus rugosa. Attention is called to the value of this common native shrub for the decoration of parks and gardens where, like many other eastern American trees and shrubs, it is rarely seen. *C. rugosa*, or *C. circinata* as it is still best known, is a shrub sometimes ten feet high which with plenty of space spreads into broad thickets. The young branches are green blotched with purple, becoming purple as they grow older. The leaves are broad, sometimes nearly circular, and dark bluish green; the flowers are ivory white, in compact clusters, and are followed in the early autumn by bright blue or nearly white fruits. This Cornel has been much planted in the Arboretum and is greatly improved by good cultivation. It can be seen in the Cornel Group at the junction of the Meadow and the Bussey Hill Roads; and the large individual plants and the great clumps on the right-hand side of the Bussey Hill Road beyond the Lilacs, and the masses among the Hickories in the group of these trees show the value of this shrub in park planting when great compact masses of foliage are needed.

Rosa multiflora, var. cathayensis is in flower on the southern slope of Bussey Hill. This is a climbing Rose with single pink clustered flowers and the Chinese representative of the better known white-flowered *Rosa multiflora*. This variety *cathayensis* is one of the most beautiful plants of its class, and is interesting as the wild plant from which have been derived the Crimson Rambler, such old-fashioned garden plants as *Rosa multiflora carnea* and *R. multiflora platyphylla*, long popular in England under the name of the Seven Sisters Rose.

1917. "Cornus Kousa." *Bulletin of popular information - Arnold Arboretum, Harvard University* 3(11), 43-44. <https://doi.org/10.5962/p.320726>.

View This Item Online: <https://www.biodiversitylibrary.org/item/216959>

DOI: <https://doi.org/10.5962/p.320726>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320726>

Holding Institution

Harvard University Botany Libraries

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.