New England, middle and middle western states must make his selection from native species, and from the species of northern Japan, Korea, northern China, Siberia, the Caucasus and eastern and northern Europe. That is, the largest and some of the handsomest and most interesting trees in the world cannot be successfully grown in the United States except in the south, and in western Washington, Oregon and California where the climate is better suited to the successful cultivation of conifers perhaps than that of any other part of the world, with the exception possibly of New Zealand where conifers from all parts of the world have grown with astonishing rapidity and vigor, and in the case of some species to a larger size than individuals of the species attain in their native lands.

In spite of the unusual and prolonged cold of the winter which followed a dry summer, the cold spring and the drought which lasted from April to September the Arboretum conifers are not in bad condition, and it is now possible to discuss with more confidence the value of many exotic species than it has been before. Only one species has been entirely lost from the effects of the severe winter. This is the blue-leaved form of *Cedrus atlantica*, a native of the mountains of Algeria. There was only one specimen in the Arboretum where it has been growing for many years in a sheltered position in the middle of a Pine grove. This beautiful tree sometimes grows fairly well south of Cape Cod, but there is little hope that it will live for more than a few years at a time in Massachusetts.

Some of the species of the northeastern states have suffered more than any of the exotic species, and several plants of the Red Spruce (*Picea rubra*) were killed or so badly injured that it was necessary to destroy them. If any coniferous tree should be hardy here it is the Red Spruce which grows on some of the high mountains of New England and close to the seashore of Maine and New Hampshire where it is fully exposed to the gales from the Atlantic. The Red Spruce, although there are now a number of healthy individuals in the Arboretum, does not take very kindly to cultivation and always grows slowly. Another eastern American tree, the short-leaved Pine (*Pinus echinata*) was injured by the winter. This tree finds its northern home on Staten Island, New York, and there have been a number of trees raised from seeds collected at this northern station growing in the Arboretum for twenty years. These all lost their leaves and several were killed; the others produced new leaves in June and now look nearly as well as ever. One specimen of this Pine raised here in 1879 from Missouri seeds also lost all its leaves but is now in comparatively as good health as it was a year ago. Several plants of the White Cedar of the eastern states (*Chamaecyparis thyoides*) lost their tops and were a good deal injured by the winter although none were killed. This plant has not taken kindly to the conditions the Arboretum affords it, but it is surprising that it is not more hardy here, as within twenty miles of Boston there are hundreds of acres of low ground covered with forests of this tree.

Cedar of Lebanon. The Cedars of Lebanon raised here from seeds gathered on the Anti-Taurus, which have been growing in the Arboretum for sixteen years and which have not before been injured by heat

or cold, in early spring lost all their leaves which had been killed by the excessive cold of the winter; they soon put out a new growth, however, and although the branches are now less densely covered with foliage than in other years the trees are in good health. Among the Spruces and Firs the Grecian and Roumanian form of *Abies cephalonica* (var. *Apollinis*) suffered the most, and although the plants are still alive they can never grow into good trees. Some small plants of *Abies cephalonica* were killed, but the large plants of this Fir in the collection are in good condition, although this tree was badly injured in other collections in Massachusetts and New York. *Abies cilicica*, which has been for many years considered one of the hardiest and handsomest of the Firs which can be grown in the northeastern states, has suffered seriously in other collections, but in the Arboretum it was little injured by the winter and is now in good condition. *Abies amabilis* from the Cascade mountains of Oregon, although always a slow-growing, shabby looking tree in cultivation, lost a good many leaves in the spring but is now in its usual health. *Abies grandis* from the northwest coast, planted in sheltered and exceptionally favorable positions, is uninjured, but for general use in New England this handsome tree should not be depended on. The Sugar Pine of the California Sierras (*Pinus Lambertiana*) and the Chinese White Pine (*P. Armandi*) lost a good many leaves but now look as well as usual. The Japanese Black Pine (*P. Thunbergii*) suffered more in the loss of its leaves, but the buds were uninjured and the trees, although somewhat disfigured, are recovering.

Among the long established trees here which are not native in New England and which show no evidence of having just passed through the most serious experience of their lives, and may therefore be considered suitable for cultivation in the northern states, are all the forms of the Norway Spruce (*Picea Abies*), the Balkan Spruce (*P. omorica*), the Caucasian Spruce (*P. orientalis*), the Siberian *Picea obovata*, *Picea Schrenkiana* from Chinese Turkestan, all the Japanese species, and the species of the Rocky Mountains *P. pungens*, *P. Engelmannii* and the western form of *P. canadensis*. The Firs not already mentioned which have not been injured are the Rocky Mountain form of *Abies concolor*, which is the most satisfactory of all Firs in the northeastern states, the Caucasian *Abies Nordmanniana*, the Japanese *A. homolepis*, (or *brachyphylla*) and *A. Veitchii*. Like the eastern American Balsam Fir (*A. balsamea*), the Rocky Mountain *A. lasiocarpa* and the Fir of central Siberia (*A. sibirica*) are perfectly hardy here, but are short-lived shabby trees in cultivation, and are not worth planting in eastern North America. The Korean *A. holophylla* was first raised at the Arboretum twelve years ago and it is still one of the rarest of all conifers in cultivation. Fortunately Wilson sent from Korea a year ago a supply of seeds of this tree; these germinated well and there are now many seedlings in this country and Europe. The twelve-year-old plant has grown well in the Arboretum; it has not suffered from cold or heat and promises to be a good tree here. The Douglas Spruce (*Pseudotsuga mucronata*) raised from seeds gathered in Colorado, has been growing in eastern Massachusetts for nearly fifty years and promises to live long here and grow to a large size. Numerous specimens of the Carolina Hemlock (*Tsuga caroliniana*) have been uninjured by the cold and drought of the year. This is one of the handsomest of

all cone-bearing trees which can be grown in this part of the country. One plant of the Japanese *Tsuga densiflora* was killed during the winter, but several others were uninjured. Small plants of the Japanese *T. Sieboldiana* have lived in an exceptionally sheltered position, but there is little hope that this beautiful tree, which is more southern in its range than the other Japanese Hemlock will ever live long in Massachusetts. A small plant of the Hemlock of the Northwest coast of North America (*T. heterophylla*), the largest and handsomest of all Hemlock trees, was uninjured in a sheltered position. There is not much probability, however, that this tree will live for more than a few years in this part of the country. The Chinese Hemlock (*T. chinensis*) was injured by the winter and probably will never be very successful here.

Pines. The European and Asiatic* *Pinus sylvestris*, the so-called Scotch Pine, the Austrian and other forms of the European Black Pine (*P. nigra*), the forms of the European *P. montana*, and the Swiss and Siberian forms of the Stone Pine (*P. Cembra*) have not been injured. The Japanese White Pine (*P. parviflora*), the Japanese Red Pine (*P. densiflora*) and the Korean form of this tree which the Japanese botanists call *Pinus gracilis*, seem able to support the New England climate without injury. The Korean Nut Pine (*P. koraiensis*) which has produced seeds in the Arboretum for several years, and the Lacebark Pine of northern China (*P. Bungeana*) are uninjured. Of the Pines of western North America only *Pinus monticola*, *P. ponderosa* var. *scopulorum*, and *P. Jeffreyi* grow successfully in the east, and these are uninjured, as are the eastern American *P. pungens* and *P. virginiana*. The northern *Pinus Banksiana*, which just reaches northern New England, with a doubtful station on Nantucket, grows well in the Arboretum but not as well as it grows much further north. Although killed last winter in some New England collections, the Japanese Umbrella Pine was little injured in the Arboretum.

None of the Arbor Vitae in the large collection of these trees here suffered with the exception of the Chinese *Thuya orientalis* which is never a very hardy or satisfactory tree in this part of the country. It is of particular interest that plants of the western Arbor Vitae, the so-called Red Cedar of the northwest (*T. plicata*), raised here from seeds gathered in Idaho, have been uninjured, for this is one of the great conifers of the world. In a sheltered position several plants of the California Incense Cedar were little injured by the winter. There has been little injury to the Junipers, and the Larches and the Chinese *Pseudolarix* have not suffered.

New Chinese Conifers. It is too soon to say much about the new conifers introduced by Wilson from northern China. All the forms of *Pinus sinensis* are growing well and appear to be hardy. All the Spruces have also done well with the exception of *Picea Sargentiana* which has suffered from cold and will probably not be hardy here. The Chinese Firs grow less well than the Spruces and only *Abies Delavayi* gives much promise of success.

1918. "Cedar of Lebanon." *Bulletin of popular information - Arnold Arboretum, Harvard University* 4(17), 66–68. <https://doi.org/10.5962/p.320887>.

View This Item Online: <https://www.biodiversitylibrary.org/item/216946>

DOI: <https://doi.org/10.5962/p.320887>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320887>

Holding Institution

Harvard University Botany Libraries

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.