Malus cerasifera. This is another of the early flowering Crabapples and is believed to be a hybrid between *M. baccata* and *M. prunifolia*. Planted in good soil and allowed sufficient room for development it will grow into a large shapely tree with a broad, round-topped, irregular head of spreading often drooping branches. The flowers are fragrant and larger than those of the other Asiatic Crabapples, with pure white or occasionally greenish petals; and the fruit, which varies in size on different plants, is globose and dull red.

Malus micromalus, which is also an early flowering plant, is one of the least known of the Crabapples. It was first sent to Europe from Japan by Von Siebold in 1856 under the name of "Kaido," a name which in Japan belongs to *Malus Halliana*. In Japan this tree is rare and known only in gardens, and by Japanese botanists is believed to have been introduced into their country from China and to be a hybrid possibly of *M. baccata* with *M. spectabilis*. The habit of this plant is more pyramidal than that of other Crabapples and this habit makes the plants conspicuous in the collection. The largest plants are covered this year with their small, pale pink, delicate flowers which will be followed by light yellow fruit, often rose color on one cheek. A plant of *Malus micromalus* first came to the Arboretum from the Paris Museum in 1888 and the plants now growing here are descendants of that plant. It is still one of the rarest of the Asiatic Crabapples in western gardens.

Malus Halliana, with its form *Parkmanii* which has double flowers, is perhaps the most distinct of all Crabapples in the color of its rose-red flowers. It is a shapely small tree, with erect and spreading stems forming a narrow vase-like head, and dark green leaves. The globose reddish fruit is not larger than a small pea. The Parkman Crab was among the first Japanese trees to reach this country direct, having been sent by Dr. George R. Hall in 1862 to Boston where it was first planted in Mr. Francis Parkman's garden on the shores of Jamaica Pond. This Crabapple is a favorite in Japanese gardens where it is known as "Kaido," but has not been found in a wild state. Whatever its origin the Parkman Crab is one of the most distinct and beautiful of the small trees which flower here during the early days of May.

Malus theifera from central and western China is closely related to Hall's Crab. It is one of Wilson's introductions through seeds sent to Veitch in 1900 and in 1907 to the Arboretum where it is now twelve feet high. It has upright, spreading, rather zigzag branches which are densely studded with short spurs which bear numerous clusters of flowers rose-red in the bud, becoming pale and almost white when fully expanded. In central China the peasants collect the leaves and from them prepare the palatable beverage which they call red tea. From this fact the specific name is derived.

Malus floribunda, by many persons considered the most beautiful of Crabapples, was introduced into Holland by Von Siebold in 1853 from Nagasaki, Japan. The place where it grows wild still remains unknown, although probably it is one of the high mountains of Kyushu. Japanese botanists and nurserymen confuse it with the Parkman Crab, and

1919. "Malus micromalus." *Bulletin of popular information - Arnold Arboretum, Harvard University* 5(4), 14–14. <https://doi.org/10.5962/p.320914>.

View This Item Online: <https://www.biodiversitylibrary.org/item/239291>

DOI: <https://doi.org/10.5962/p.320914>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320914>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.