Wilson has not seen it in Japanese gardens. It is a broad, round-topped, treelike shrub sometimes twenty-five feet tall with stout branches and slender arching and pendent branchlets. The clustered flowers are white when fully expanded, rose-red in the bud, and as they open in succession the two colors make a beautiful contrast. The fruit is about the size of a pea, yellowish or yellowish brown; from some plants it falls in the early autumn, on others it remains on the branches during the winter or until devoured by birds who are particularly fond of it. Several plants with persistent fruit are growing close to the Administration Building in the Arboretum, and during the winter are filled with numerous species of birds, including pheasants who are fond of these Crabapples. A hybrid between *M. floribunda* and perhaps *M. cerasifera* appeared in the Arboretum among a lot of seedlings of *M. floribunda* in 1883 and has been named *M. Arnoldiana*. It has the habit and abundant flowers of *M. floribunda*, but the flowers and fruit are nearly twice as large. It is a handsomer plant than *M. floribunda* and one of the most beautiful of the Crabapples in the Arboretum.

Malus Sieboldii is another of the species introduced from the gardens of Japan into Europe by Von Siebold in 1853. It is a low, dense shrub of spreading habit with the leaves on vigorous branchlets three-lobed, small flowers white tinged with rose in color, and small yellow fruits. A good specimen may be seen on the left hand side of the Forest Hills Road. Von Siebold's Crab is really a dwarf form of a species common on the Korean Island of Quelpaert, and on the mountains of central Japan and Hokkaido, to which the name var. *arborescens* has been given. This is a tree often thirty feet or more tall, with ascending, wide-spreading branches, twiggy branchlets and minute fruit yellow on some and red on other individuals. Although the flowers are small, they are produced in immense quantities, and this species has the advantage of flowering later than the other Asiatic Crabapples. Another variety of Von Siebold's Crab (var. *calocarpa*), raised in the Arboretum from seed sent in 1890 from the Nikko mountains of Japan by Dr. W. Sturgis Bigelow of Boston, has bright red fruits each half an inch in diameter. When in fruit this is the handsomest of the Japanese Crabapples.

Malus Sargentii from salt marshes in the neighborhood of Muroran in northern Japan, where it was discovered by Professor Sargent in 1892, has qualities which give it a field of usefulness peculiarly its own. This species is a dwarf with rigid and spreading branches, the lower branches flat on the ground; it is well suited for covering slopes and banks. The flowers are in umbel-like clusters, saucer-shaped, round and of the purest white, and are followed by masses of wine-colored fruit which is covered by a slight bloom and unless eaten by birds remains on the plants well into the spring.

Malus spectabilis, cultivated by the Chinese from time immemorial and introduced from Canton in 1780, was the first of the Asiatic Crabapples cultivated in Europe. Like several other species it is not yet known in a wild state but is probably of hybrid origin. It is a tree

from twenty-five to thirty feet high, with a wide vase-shaped crown made of numerous spreading and ascending branches and short branchlets. The flowers are pale pink, more or less semidouble and very fragrant: and the fruits are pale yellow, nearly globose, and about three-quarters of an inch in diameter.

Malus Scheideckeri is supposed to be a hybrid between *M. spectabilis* and some unknown species, possibly *M. micromalus*. It is a small pyramidal tree with small flowers produced in great abundance, and is well worth a place in every collection of these trees.

Malus prunifolia var. *rinki*, the Apple cultivated in Japan for its fruit before the advent of foreigners and of Chinese origin, has been mentioned in former Bulletins, notably that of May 15, 1916. The wild type of this apple was discovered by Wilson in central China in 1907. From seeds sent to the Arboretum plants were raised and one of them is now blooming for the first time; it is on Bussey Hill, in the collection of Chinese Apples, Pears and Cherries. This is now a small tree about ten feet high, with flowers like those of the common Apple and fully an inch and a half across. The fruit of *rinki* is longer than broad, yellow with a reddish cheek or entirely red, and the persistent calyx is raised, and not depressed as in the common Apple. This is the wild parent of the race of Apples long cultivated in the Orient, and since it fruits freely in the hot moist valleys of central China equally as well as in the cold regions of northern Korea it may prove of value to pomologists in breeding new races of Apples.

Space does not permit even a brief mention of all the species and hybrids of all the Asiatic Crabapples in the Arboretum collection. Among them, however, are trees suitable for the avenue, park or garden, shrubs for lawn borders and the slopes of banks, all absolutely hardy in the coldest parts of New England, and all to be depended upon to produce in spring blossoms in profusion. The plants grow quickly in good soil, love to have the breezes blow freely through their branches, and many of them begin to flower and produce fruit when only a few years old. In collections like that of the Arboretum they hybridize freely, and the species can only be propagated by grafting or budding.

Asiatic Quinces. In the Shrub Collection many varieties of these plants are growing and are now in full bloom. The flowers vary from white, flesh, pink and salmon to scarlet and fiery crimson, and no group of plants has more vivid blossoms. The numerous forms are derived from the Chinese *Chaenomeles lagenaria* (better known perhaps as *Pyrus japonica*), and the true Japanese *C. japonica* (*C. Maulei*) and have been cultivated for many centuries in the gardens of China and Japan.

The earliest of the Lilacs to bloom, the Chinese *Syringa affinis* and *S. oblata*, are in flower and the collection will probably be at its best about May 24th. The red-flowered Japanese Azalea (*Rhododendron Kaempferi*) is just opening its flowers on Azalea Path, and Fothergillas, many Spiraeas, Pearl Bushes, many Honeysuckles, Barberries and other interesting plants will be in bloom when this Bulletin reaches its Boston readers.

1919. "Malus spectabilis." *Bulletin of popular information - Arnold Arboretum, Harvard University* 5(4), 15–16. <https://doi.org/10.5962/p.320920>.

View This Item Online: <https://www.biodiversitylibrary.org/item/239291>

DOI: <https://doi.org/10.5962/p.320920>

Permalink: <https://www.biodiversitylibrary.org/partpdf/320920>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.