however, will be needed to settle this question. One of the Spruces of northern Japan, *Picea jezoensis*, and its southern form (var. *hondoensis*) have grown miserably in Massachusetts up to the present time and give little promise of ever being valuable in this climate. The rare Tiger-tail Spruce (*Picea polita*) grows rapidly and is a perfectly hardy tree, but often begins to lose its lower branches before it is thirty feet tall. The northern *Picea Glehnii*, introduced by the Arboretum from seeds obtained by Professor Sargent in Hokkaido in 1892, now promises to be a handsome tree in this climate. The handsomest of the Japanese Spruces, judged by the few trees cultivated in Massachusetts, is *Picea bicolor*, or as it is more commonly called, *P. Alcockiana*. This tree, which is rare in Japan, was discovered in 1862 during the first ascent of Fuji-san by Europeans. Seeds were collected at this time and sent to Europe and it is probable that the few large trees of this Spruce cultivated in the United States and Europe were raised from these seeds. The trees cultivated in Europe under this name are usually *Picea jezoensis* var. *hondoensis*, and the Arboretum knows only five of these trees in the United States, two in the Hunnewell Pinetum, one also planted by Mr. Hunnewell in the grounds of the Town Hall at Wellesley, and two on the Phillips Estate in North Beverly, Massachusetts. In the Arboretum there are only small unsatisfactory grafted plants, and for years the Arboretum has tried without success to obtain a supply of seeds from Japan, for, judging by our present knowledge of the behavior of Spruces in this climate, *Picea bicolor* promises to be the handsomest which can be grown in this part of the country.

Abies. Fir-trees, like the Spruces, are widely distributed with many species through northern and elevated regions of the Northern Hemisphere, growing rather further south than the Spruces, as one Fir-tree grows in Mexico, one in Spain, one in northern Africa and several in southeastern Europe and Asia Minor. Many of the Firs are large and handsome trees, but the genus has not contributed much to the beauty of our northern plantations. Many of the handsomest and most interesting species are not hardy here, and several of the others are not presentable for more than a few years. Judging from the results which have now been obtained with these trees there are only two Firs which can be depended on to retain their beauty here for more than fifty years. These are the western American White Fir (*Abies concolor*), especially the form which grows on the mountains of southern Colorado, and the Japanese *Abies homolepis* or *brachyphylla*, a splendid tree with dark green leaves white on one surface and large purple cones. The variety of this tree with green cones (var. *umbellata*) has grown more rapidly in the Arboretum than the purple-coned tree, but it is a tree of more open habit and with lighter green leaves, and is less valuable as an ornamental tree. *Abies cilicica* from Asia Minor and *A. cephalonica* from southeastern Europe have grown well in Massachusetts for many years; although they have now nearly recovered, these two trees suffered severely in the cold winter of 1917-18.

Thuja, the name of the Arbor Vitaes, is a small genus confined to eastern and western North America, Japan, Korea and north China.

1919. "Abies." *Bulletin of popular information - Arnold Arboretum, Harvard University* 5(17), 67–67. <https://doi.org/10.5962/p.321003>.

View This Item Online: <https://www.biodiversitylibrary.org/item/239291>

DOI: <https://doi.org/10.5962/p.321003>

Permalink: <https://www.biodiversitylibrary.org/partpdf/321003>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.