ARNOLD ARBORETUM
HARVARD UNIVERSITY BULLETIN
OF
POPULAR INFORMATION

JAMAICA PLAIN, MASS.

JUNE 30, 1922

English Elms. There is still apparently much confusion in popular understanding in this country in regard to the trees known here as "English Elms," and it does not seem to be generally understood that there are four distinct species of Elm-trees now growing naturally in England, and that among the trees sometimes cultivated are hybrids of these species which also in the popular mind pass as English Elm-trees.

Ulmus procera. This is the name now adopted for the tree which is generally known as English Elm in Boston where it has proved one of the best foreign trees ever planted in Massachusetts. It has been growing here for more than a century, and nearly one hundred years ago Major Paddock had a nursery at Milton for the propagation and sale of this tree. Probably no tree, native or foreign, which has been planted in the neighborhood of Boston has grown to such a large size. The Paddock Elms, which stood on Tremont Street in front of the Granary Burying Ground, were of this species, as were the great Elms on the Tremont Street Mall of the Common which were killed by the Subway. The Elm-trees on each side of the Shaw Monument opposite the State House are of this species, and there are still large specimens in the suburbs of the city. This is the common Elm-tree of southern England where it grows usually in hedge-rows, although it has been largely planted in parks. It often grows one hundred feet tall with a massive stem covered with dark deeply furrowed bark, spreading or ascending branches which form a comparatively narrow oval head, and slender branchlets thickly covered during their first year with down.

The leaves are broadly oval or ovate, oblique at base, dark green and rough on the upper surface and covered below with soft down; they are from two to three inches long with about twelve pairs of veins, and their stalks are only about one-fifth of an inch in length. This tree very rarely ripens fertile seeds in England or in this country, but it produces suckers in great numbers and is propagated entirely by means of these. As this tree so rarely produces seeds few varieties are known, but a small-leaved Elm (var. *viminalis*) is believed to be a seedling of it. Of this little Elm there are forms on which the leaves are blotched with white and with yellow.

Ulmus foliacea, or nitens. This is another English Elm which differs from the last in its paler bark, in its smooth or nearly smooth branchlets, that is without a covering of down and in its leaves which are smooth and shining on the upper surface, only slightly downy below early in the season and from two to three and a half inches long. This tree produces fertile seeds in abundance and seedlings are raised in European nurseries. It is widely distributed over central and southern Europe and grows also in northern Africa and eastern Asia. Several geographical forms are recognized; the most distinct of these are the Cornish and the Guernsey Elms which are trees of medium size with erect growing branches which form a narrow pyramidal head. Plants of these two forms are not always hardy in Massachusetts. Another form, common in Hertfordshire, is a large tree with wide-spreading and pendulous branches and at its best, although not so tall is almost as handsome as our American White Elm (*U. americana*). Another form (var. *umbraculifera*) from Persia and Armenia is interesting from its compact globose head. This tree might perhaps be made useful in formal gardens. On many trees of *Ulmus foliacea* the branches are furnished with corky wings (var. *suberosa*), and the so-called English Elms with such branchlets occasionally seen in this country are usually of this variety. The seedling trees of this Elm which have been imported from European nurseries vary in habit, in the size of their leaves and in their hardiness; and the unhealthy and generally unsatisfactory Elm-trees which have been planted in considerable numbers in eastern Massachusetts during the last twenty years are in nine cases out of ten seedling forms of *U. foliacea*.

Ulmus glabra. This is another widely distributed European Elm which is often called Scotch Elm or Wych Elm by English-speaking people. This is a tree with a trunk and branches which remain smooth for many years. It can always be recognized, too, by the large obtuse buds covered by pale brown hairs and by its dark dull green leaves abruptly pointed or three-lobed at the apex, oblique and unsymmetrical at the base, rough above, downy below, and from four to six inches long with stalks shorter than those of other Elm-trees. This tree does not sucker but produces fertile seeds in great quantities, and more abnormal seedling forms of this tree have been raised than of any other Elm. The well-known Camperdown Elm is a form of this tree with regularly pendulous branches which is often planted in suburban gardens to make natural arbors; another form (var. *pendula*) has horizontally spreading pendulous branches which form an unsymmetrical, flat-topped head.

1922. "Ulmus procera." *Bulletin of popular information - Arnold Arboretum, Harvard University* 8(13), 49–50. <https://doi.org/10.5962/p.321258>.

View This Item Online: <https://www.biodiversitylibrary.org/item/216933>

DOI: <https://doi.org/10.5962/p.321258>

Permalink: <https://www.biodiversitylibrary.org/partpdf/321258>

Holding Institution

Harvard University Botany Libraries

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.