tined to play an important part in the decoration of parks and gardens in the northeastern United States where few *Rhododendrons* or other broad-leaved evergreen plants can be grown. They are as hardy as the hardiest of the *catawbiense* hybrids, and flowering two or three weeks later than these prolong the flowering time of hybrid *Rhododendrons* into July, that is to the time when the conspicuous flowering of trees and shrubs is not abundant.

Schizophragma hydrangeoides, now that it has at last, after forty years of failure, found a place that suits it on the east side of the Administration Building, is growing rapidly and promises to cover as much space as the great plant of the Japanese Climbing Hydrangea which is its neighbor. It is already half way to the top of the building, and its value as a flowering plant in July is now shown by its conspicuous flower-clusters. The Japanese *Schizophragma* now grows as rapidly as the Climbing Hydrangea and clings as firmly to a brick wall. The leaves are smaller, more circular in shape, more coarsely toothed and darker and duller in color. The inflorescence, which is terminal on short lateral branchlets, which stand out from the stems, is interesting but not perhaps as showy as that of the Hydrangea, for instead of the surrounding ring of neutral flowers there are only two neutral flowers to each of the divisions to the large compound cluster of perfect flowers; these neutral flowers are snow white, ovate, often an inch or more long, and hang on long slender stems an inch in length. *Schizophragma hydrangeoides* seems to be a rare plant in American and European gardens, and in this country *Hydrangea petiolaris* is often sold for it. The Chinese species, *S. integrifolia* introduced by Wilson, has not yet found a place in the Arboretum which suits it, and has not proved hardy here. It is a handsomer plant than the Japanese species with much larger sterile flowers.

Decumaria barbara is another climbing plant of the Saxifrage Family which is now flowering in the Arboretum Nursery. It grows naturally by the banks of streams and in swamps from southeastern Virginia to central and western Florida, western Louisiana, and western Tennessee, often climbing up the trunks of trees by its aerial roots to the height of thirty feet. This handsome and interesting plant has dark green and lustrous leaves, small, white, fragrant flowers in large, terminal, compound clusters, and capsular, urn-shaped fruit with a persistent style and stigma. It is rare in gardens, certainly in those of the northeastern United States, but there is an old and well established plant in that of Mr. N. T. Kidder in Milton, Massachusetts, which flowers every year. It is now growing well and promises to become established in the Arboretum. The Chinese species introduced by Wilson is growing well in a cold pit here but has not yet flowered.

Itea virginica, another plant of the Saxifrage Family, was in flower last week in the Shrub Collection. It is a shrub two or three feet high, with simple, alternate, minutely serrate, deciduous leaves and small white flowers in terminal erect racemes. This interesting little plant is widely distributed from New Jersey to Florida and Louisiana, and northward to Missouri and southern Illinois, ascending on the Car-

1922. "Decumaria barbara." *Bulletin of popular information - Arnold Arboretum, Harvard University* 8(14), 54-54. <https://doi.org/10.5962/p.321267>.

View This Item Online: <https://www.biodiversitylibrary.org/item/216933>

DOI: <https://doi.org/10.5962/p.321267>

Permalink: <https://www.biodiversitylibrary.org/partpdf/321267>

Holding Institution

Harvard University Botany Libraries

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.