garian species which often does not bloom here until after the middle of June.

Syringa oblata was the first Lilac from eastern China to reach England where it was sent some sixty years ago by Robert Fortune who had found it in a Shanghai garden. It reached the United State certainly as early as 1869 and perhaps earlier. It is a round-topped shrub with heart-shaped leaves which, unlike those of other Lilacs, are thick and coriaceous and in the autumn turn scarlet. It is one of the first Lilacs to bloom in the spring here. This plant has not been found growing wild and there is no record that it has been seen by anyone in China since Fortune's time. It is probably a garden form of S. affinis, a white-flowered form commonly planted and probably the only Lilac in the gardens of Peking. Another form probably of S. oblata is the purple-flowered plant from northern China usually called the variety Giraldii of S. affi-The Korean S. dilatata is probably also only a wild form of S. The two best known of the Lilacs of eastern Asia, S. pubescens and S. villosa, were raised here from seeds sent in 1883 by Dr. Bretschneider of the Russian Legation in Peking. S. pubescens is a tall shrub with erect stems, small leaves and broad clusters of small pale mauve flowers with a long slender corolla tube. For the fragrance of the flowers, which is more pungent and delightful than that of any other Lilac, S. pubescens should find a place in every northern garden. The plants in the United States have failed to produce seeds, and as this species is unusually difficult to increase by cuttings it has remained one of the rarest Lilacs in American gardens. S. villosa is a large, round-topped bush, from ten to twelve feet tall and wide, with large, broad, elliptic to oblong leaves bright green and dull on the upper surface and pale below, and broad or narrow clusters of fleshcolored or nearly white flowers which have a rather disagreeable odor like those of the Privet. In spite of this drawback S. villosa is a valuable plant; its habit is excellent, it flowers freely every year, and the flowers do not open until most of those of the other Lilacs have faded. Of the new Lilacs from western China and Korea raised from seeds collected by Wilson and other travelers the most promising are S. Meyeri, S. microphylla, S. Julianae, S. tomentella, S. reflexa, S. Komarowii, S. Sweginzowii and S. dilatata.

Hybrid Lilacs. The first hybrid Lilac appeared in the Botanic Garden in Rouen in 1810 and was the result of crossing S. vulgaris and S. persica. It is one of the most delightful of all Lilacs and grows into a bush ten or twelve feet high and broad and of rather open habit. It is very hardy and blooms freely every year, and should be in every garden where Lilacs are grown. Its flowers resemble those of the Persian Lilac and are produced in massive clusters sometimes two feet in length, and are so heavy that the slender branches can hardly support them. There are forms with darker red flowers and with nearly white flowers. Through a misunderstanding of its origin this plant unfortunately must be called S. chinensis.

The next hybrid Lilac to appear was S. hyacinthiflora which is the result of crossing S. oblata with S. vulgaris coerulea plena. It is a large, round-topped shrub of excellent habit, with leaves resembling in shape those of S. oblata, and small clusters of semi-double, extremely

fragrant flowers. It blooms earlier than any of the forms of S. vulgaris but has little to recommend it as a garden plant.

The general name of Syringa Henryi has been given to a group of hybrid Lilacs between S. Josikaea and S. villosa which was obtained in Paris. The handsomest of this breed, S. Lutêce, is a valuable addition to the late-flowing Lilacs. Another interesting hybrid was obtained by Lemoine by crossing S. vulgaris with the variety of S. affinis with fragrant violet-colored flowers (var. Giraldii). The plants grow rapidly and are tall narrow shrubs. Like their Chinese parent they bloom early and the flowers are fragrant. Forms of this hybrid, Berryer, Claude Bernard, Lamartine, Mirabeau, Pascale and Vauban are in the Arboretum Collection.

A lover of Lilacs living in Manitoba has recently obtained a hybrid of S. pubescens and S. villosa which promises to be interesting, and there is still much work to be done in raising new hybrids between the species of western China.

Rhododendron (Azalea) obtusum var. Kaempferi is the only red-flowered Azalea which is hardy in this climate. It has been largely used in the Arboretum and is now flowering at least two weeks earlier than usual. Its flowers furnish the most surprising and spectacular display of the year. They are delicate, however, and when fully exposed to the sun lose their color; and this Azalea gives most satisfaction when it is planted in the shade of trees or on the northern border of a wood of conifers. It is planted in masses at the lower end of Azalea Path and in a large group under the shade of the Hemlocks on Hemlock Hill and on the northern edge of Hemlock Hill in a long narrow band between the Hemlocks and the Laurels. The tallest plants in the Arboretum are now more than ten feet high and rarely fail to flower profusely even when growing in complete shade. This is one of the best shrubs which has been introduced by the Arboretum.

Rhododendron (Azalea) Vaseyi from the southern Appalachian Mountains is flowering profusely this year. The pure pink flowers appear on the leafless branchlets and in delicacy and purity of color are not surpassed by the flowers of any other plant. It is only in comparatively recent years that this Azalea has been known to botanists or has found its way into gardens. It is perfectly hardy, the flower-buds are not injured by severe cold, and in time it will grow into a tall, usually rather narrow shrub. This beautiful Azalea has been planted on both sides of the Meadow Road, the largest group being at the northern end of the first pond.

Rhododendron (Azalea) luteum, a native of the Caucasus, has bloomed in the Arboretum several times and, although the buds are often injured, is in good condition this year. It is growing on the right hand side of Azalea Path below the plant of Rhododendron (Azalea) reticulatum. If the flower-buds of this Caucasian plant were hardier this would be one of the most delightful of all Azaleas as the flowers are charming in color and more fragrant than those of any other Azalea.



1925. "Hybrid Lilacs." *Bulletin of popular information - Arnold Arboretum, Harvard University* 11(6), 23–24. https://doi.org/10.5962/p.321570.

View This Item Online: https://www.biodiversitylibrary.org/item/216996

DOI: https://doi.org/10.5962/p.321570

Permalink: https://www.biodiversitylibrary.org/partpdf/321570

Holding Institution

Harvard University Botany Libraries

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Arnold Arboretum of Harvard University

License: http://creativecommons.org/licenses/by-nc-sa/4.0/

Rights: https://biodiversitylibrary.org/permissions

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.