

SCALOPS LATIMANUS.—BACH.

TEXAN SHREW MOLE.

SYNONYME.

SCALOPS LATIMANUS.—Bach, Boston Jour. Nat. History, vol. i., p. 41.

DESCRIPTION.

Larger than the common shrew-mole, intermediate in size between *S. Townsendi* and *S. Breweri*. Hair longer and thinner than in either of the other species, and slightly curled. Palms larger than in any other known species. Tail naked.

COLOUR.

Colour nearly black.

DIMENSIONS.

	Inches.	Lines.
Length to root of the tail, - - - - -	7	7
“ of the tail, - - - - -		10
Breadth of the palm, - - - - -		10
“ of the tarsus, - - - - -		7

GEOGRAPHICAL DISTRIBUTION.

Mexico and Texas.



Audubon, John James and Bachman, John. 1854. "Scalops latimanus, Texan Shrew Mole." *The quadrupeds of North America* 3, 323–323.

<https://doi.org/10.5962/p.322536>.

View This Item Online: <https://www.biodiversitylibrary.org/item/108515>

DOI: <https://doi.org/10.5962/p.322536>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322536>

Holding Institution

Duke University Libraries

Sponsored by

Duke University Libraries

Copyright & Reuse

Copyright Status: Not provided. Contact Holding Institution to verify copyright status.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.