RECTES JOBIENSIS, Meyer.

RECTES JOBIENSIS, *Meyer*.

Jobi-Island Wood-Shrike.

Rectes jobiensis, Meyer, Sitz. k. Akad. der Wiss. zu Wien, lxi. p. 205 (1874).—Sclater, Ibis 1874, p. 417.—Salvad. Ann. Mus. Civ. Genova, vii. p. 173 (1875), viii. p. 40 (1876).—Sharpe, Cat. B. Brit. Mus. iii. p. 287 (1877).—Gieb. Thes. Orn. iii. p. 412 (1877).
Rhectes jobiensis, Salvad. Ann. Mus. Civ. Genova, xv. p. 43 no. 12 (1879).—Id. Orn. Papuasias, &c. ii. p. 201 (1881).

THE Shrike-like genus *Rectes* consists of a small number of birds found in New Guinea and in the neighbouring islands. It is divided into two sections, consisting of those species which have a distinct cap of darker colour on the head, and those species which have no cap whatever; and the present bird will be seen at once to belong to the second section. Its nearest allies therefore are *Rectes cervineiventris* from Waigiou, and *R. cristatus* of New Guinea; but it is very distinct from both of these by reason of its foxy red colour both above and below.

The present bird was discovered by Dr. A. B. Meyer in the island of Jobi, which is situated in the Bay of Geelvink, in North-western New Guinea. He preserved several specimens of it; and it has also been met with there by Dr. Beccari and in the neighbouring little islet of Krudu.

Dr. Meyer says that the note of this bird is a "chrrr;" and its habits are doubtless the same as those of the rest of the genus, of which at present we unfortunately know but little.

The following description of the species I take from Mr. Bowdler Sharpe's Catalogue of Birds:—

Adult male. General colour above bright foxy red, paler on the head; wing-coverts like the back; quills dark brown, externally rufous, the inner secondaries entirely rufous; tail-feathers chestnut, the shafts rufous; lores, sides of face, and under surface of body deep rich tawny buff, darker on the throat and fore neck, and inclining to chestnut on the lower flanks and under tail-coverts; under wing-coverts bright tawny, like the breast; quills dusky brown below, rufous on the outer web and along the inner web; bill light yellowish horn-colour; feet blackish, claws pale. Total length 9·2 inches, culmen 11·5, wing 4·65, tail 4·35, tarsus 1·35.

Adult female. Similar in colouring to the male.

Total length 10·5 inches, wing 4·6, tail 4·3, tarsus 1·25.

Young male. Similar in colour to the adult, but rather duller; the bill shorter and duller brown.

The above descriptions were taken from the typical specimens lent to Mr. Sharpe by Dr. Meyer; and from one of these I have drawn the accompanying figure, which is of about the natural size.

Gould, John and Sharpe, Richard Bowdler. 1881. "Rectes jobiensis, Jobi-Island Wood-Shrike [Pl. 55]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 1(XII), –.

<https://doi.org/10.5962/p.322654>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229855>

DOI: <https://doi.org/10.5962/p.322654>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322654>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.