GRAUCALUS PUSILLUS, *Ramsay*.

Ramsay's Cuckoo-Shrike.

Graucalus, sp. nov., Ramsay, Nature, 1879, p. 125.

Graucalus pusillus, Ramsay, Proc. Linn. Soc. N. S. Wales, iv. p. 71 (1879).—Salvad. Ann. Mus. Civic. Genov. xv. p. 35 (1879).—Id. Ibis, 1880, p. 128.—Id. Orn. Papuasia, etc. ii. p. 140 (1881).

Graucalus dussumieri (?), Ramsay, Proc. Linn. Soc. N. S. W. iv. p. 71 (1879).

Graucalus salamonis, Ramsay, Proc. Linn. Soc. N. S. W. iv. p. 314 (1879).

THIS species is a small island race of *Graucalus axillaris*, the males of the two species being almost identical in colour. In the present bird, which comes from the Solomon Islands, the colour is a clearer slaty blue in both sexes and the under wing-coverts and axillaries are barred with black and white, and the bars are narrower.

In the females of *G. pusillus* the thighs are uniform blue-grey, whereas in *G. axillaris* they are barred with black and white like the rest of the under surface.

As far as is known at present, this Cuckoo-Shrike is found only in Guadalcanar, where it was discovered by Mr. Cockerell.

The following descriptions have been drawn up from the typical specimens:—

Adult male. General colour above clear blue-grey; wing-coverts like the back; large outer feather of bastard wing black; primary-coverts and quills black, externally blue-grey, more hoary on the outer webs of the primaries; tail-feathers black, the centre feathers ashy for two thirds of their length on the inner web, not quite so extended on the outer web, the outermost feather dull ashy at the tip; head clear blue-grey like the back; a narrow frontal line and lores black, occupying a distinct space in front of the eye and extending below the latter, as well as the upper and under edge of the eyelid; sides of face, ear-coverts, cheeks, and entire under surface of body clear blue-grey, with a tiny spot of black at the base of the chin and base of cheeks; under tail-coverts darker grey, with blackish shaft-stripes; axillaries barred, ashy grey on the inner web and on the base of the outer, the latter otherwise barred with black and ashy whitish; under wing-coverts slaty blue, with less distinct bars of black and ashy whitish; quills ashy blackish below, greyer on the inner web especially near the base, the inner web edged with white. Total length 9 inches, culmen 0·7, wing 4·95, tail 4·2, tarsus 0·8.

Adult female. Differs from the male in having the base of the forehead greyish black instead of deep black, the loreal patch being also ashy blackish; the throat and chest are uniform blue-grey, but the rest of the under surface is barred with black and ashy white on a blue-grey ground; on each side of the lower back a conspicuous patch of silky white; thighs uniform blue-grey. Total length 8·5 inches, culmen 0·65, wing 4·95, tail 3·1, tarsus 0·8.

The Plate represents the male and female birds of the natural size, the figures being drawn from the type specimens which Mr. Ramsay has kindly lent to us.

[R. B. S.]

Gould, John and Sharpe, Richard Bowdler. 1884. "Graucalus pusillus, Ramsay's Cuckoo-Shrike [Pl. 4]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 2(XVII), –.
<https://doi.org/10.5962/p.322659>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229877>

DOI: <https://doi.org/10.5962/p.322659>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322659>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.