PECILODRYAS BIMACULATA.

W. Hart del. et lith.

Walter Imp.

PŒCILODRYAS BIMACULATA.

Black-and-White Flycatcher.

Myiolestes? *bimaculata*, Salvad. Ann. Mus. Civic. Genov. vi. p. 84.

Pachycephala? *bimaculata*, Salvad. Ann. Mus. Civic. Genov. vii. p. 935 (1875), x. p. 142 (1877).

Pœcilodryas bimaculata, Sharpe, Notes Leyden Mus. i. p. 25 (1878).—Id. Cat. Birds Brit. Mus. iv. p. 244 (1879).

—Salvad. Ann. Mus. Civic. Genov. xiv. p. 502 (1879).—Id. Orn. Papuasias, etc. ii. p. 85 (1882).

Pœcilodryas sylvia, Ramsay, Trans. Linn. Soc. N. S. Wales, viii. p. 5 (1883).

THE genus *Pœcilodryas* consists of two groups, or sections, characterized by the colour of the abdomen; in one section the abdomen is white, in the other yellow. The white-bellied group may be further subdivided into those which have the throat white and those which have a black throat. The present species belongs to the latter section, which now contains three species—*P. bimaculata*, *P. æthiops*, and *P. albinotata*, the latter being distinguished by its bluish-grey upper surface.

P. bimaculata and *P. æthiops* have the upper surface black, with the rump and upper and under tail-coverts white; but they may easily be distinguished from each other, *P. bimaculata* having the abdomen white, a long white patch on the sides of the fore neck and chest, and the inner wing-coverts black; whereas in *P. æthiops* the abdomen is black, the inner wing-coverts are white, forming a shoulder-patch, and there is no white on the sides of the fore neck.

The present species was discovered in north-western New Guinea by Signor D'Albertis, and has been met with in the same locality by Dr. Beccari and M. Laglaize. It has been more recently obtained in the Astrolabe Mountains by Mr. Goldie, and a specimen from that range of mountains has been lent to me by Mr. Ramsay.

The following description is copied from the 'British Museum Catalogue of Birds':—

"*Adult male*. Above velvety black; upper tail-coverts white, forming a band across; wings and tail black; sides of face, sides of neck, throat, and breast black, as well as the flanks and thighs; abdomen, vent, and under tail-coverts white; on each side of the chest a broad line of pure white feathers running from the sides of the lower throat to the sides of the upper breast; under wing-coverts black; quills ashy black below. Total length 5.1 inches, culmen 0.6, wing 3.3, tail 2, tarsus 0.85."

The figures in the Plate, which represent an adult bird of the natural size, are drawn from the Astrolabe-Mountain specimen lent to us by Mr. Ramsay, the type of his *P. sylvia*. [R. B. S.]

Gould, John and Sharpe, Richard Bowdler. 1884. "Pœcilodryas bimaculata, Black-and-White Flycatcher [Pl. 17]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 2(XVI), -. <https://doi.org/10.5962/p.322672>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229877>

DOI: <https://doi.org/10.5962/p.322672>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322672>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.