TODOPSIS GRAYI, Wall.

TODOPSIS GRAYI, *Wall.*

Gray's Todopsis.

Todopsis grayi, Wallace, P. Z. S. 1862, p. 166.—Finsch, Neu-Guinea, p. 168.—Beccari, Ann. Mus. Civic. Genov. vii. p. 709.—Meyer, Sitz. k. Akad. Wien, lxix. pp. 81, 212.
Tchitrea grayi, Gray, Hand-list of Birds, i. p. 334, no. 5034.

THIS is one of the most distinct species of *Todopsis* yet discovered, being totally unlike any of the other members of the genus, its silvery grey or glaucous coloration being quite sufficient to distinguish it at a glance. It was discovered by Mr. Wallace in the mountains of Sorong in North-western New Guinea; and it still remains one of the rarest species of the genus, as I am only aware of one other instance of its capture, viz. the specimen mentioned by Beccari as having been obtained by Mr. Bruijn's hunters in the Arfak mountains. It is probable, however, that the *Todopsis sericyanea* of Rosenberg, and the *Myiagra glauca* of Schlegel may also belong to the present species; but at present I have not been able to determine this for certain.

It is much to be regretted that Dr. Beccari never met with this bird himself, as from his pen we should have undoubtedly received some information as to its habits, of which I can at present record absolutely nothing. The following description of the type is from Mr. Sharpe's 'Catalogue of Birds':—

Adult. General colour above dull blue, brighter and more cobalt on the lower back and rump; wing-coverts brown, slightly washed externally with olive, and tipped with dull verditer blue; quills brown, broadly washed externally with olivaceous brown; tail-feathers brown, externally bluish, slightly washed with olivaceous on the margins and tipped with white; head dull greyish blue, inclining to ashy verditer above the eyes and on the sides of the crown: feathers in front of the eye blackish, as also a broad streak above the ear-coverts; round the eye a circlet of verditer feathers; ear-coverts verditer blue, the feathers rather elongated, and streaked with brighter cobalt; cheeks and throat silvery whitish, with a bluish gloss, the rest of the under surface pale ashy blue, washed with a cobalt shade; under wing-coverts like the breast; thighs olivaceous brown.

Total length 5·7 inches, culmen 0·7, wing 2·55, tail 2·5, tarsus 1·0.

The figures in the Plate, which represent the species in two different attitudes, are taken from the type specimen in the British Museum.

Gould, John and Sharpe, Richard Bowdler. 1878. "Todopsis grayi, Gray's Todopsis [Pl. 24]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 2(VIII), –.

<https://doi.org/10.5962/p.322679>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229877>

DOI: <https://doi.org/10.5962/p.322679>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322679>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.