RHIPIDURA LEUCOTHORAX, Salvad.

RHIPIDURA LEUCOTHORAX, *Salvad.*

White-breasted Fantail Flycatcher.

Rhipidura leucothorax, *Salvad. Ann. Mus. Civic. Genov.* vi. p. 311 (1874).—*Id.* & *D'Albert. op. cit.* vii. p. 820 (1875).—*Salvad. op. cit.* ix. p. 25 (1876), x. p. 134 (1876).—*Oustalet, Bull. Soc. Philom.* 1877, p. 6.—*Sharpe, Cat. B. Brit. Mus.* iv. p. 327 (1879).—*D'Albert. & Salvad. Ann. Mus. Civic. Genov.* xiv. p. 60 (1879).—*Salvad. t. c.* p. 498 (1879).—*Id. Orn. Papuasias, etc.* ii. p. 58 (1881).

Rhipidura episcopalis, *Ramsay, Proc. Linn. Soc. N. S. W.* ii. p. 371 (1878).—*Salvad. Ibis*, 1879, p. 323.

THIS species belongs to the section of the genus *Rhipidura* with blackish under tail-coverts and rounded white spots on the chest. The large white breast-patch distinguishes it from all its near allies.

In North-western New Guinea the present bird has been found in Hatam and at Mariati by Bruijn, and *D'Albertis* procured specimens in Southern New Guinea on the Fly River, near Hall Bay and at Naiabui. *Mr. Ramsay* has also described a bird as *R. episcopalis* from Goldie River, in the interior of South-eastern New Guinea, which is certainly the same as *R. leucothorax*. It should be noticed, however, that the single example in the British Museum, determined by Count *Salvadori* himself to be *R. leucothorax*, differs from his description in having the chin white, and may belong to another species, unless it is the young bird.

The following is a description of the type specimen of *Rhipidura episcopalis*, which has been lent to us by *Mr. Ramsay* :—

Adult male. General colour above earthy brown, the head a little more dusky than the back and blacker on the forehead; scapulars like the back; wing-coverts black, spotted with white at their ends; primary-coverts blackish; quills dark sooty brown; upper tail-coverts black; tail-feathers black, tipped with white, increasing in extent toward the outermost; lores and base of forehead, feathers above and below the eye, and ear-coverts black, extending on to the sides of the neck; above the eye a band of white separated from the crown by a narrow line of black; hinder part of cheeks white, widening out on the sides of the throat and neck, and forming a large white patch; fore part of cheeks and throat black, widening out on the fore neck and chest, and having rounded white spots on the latter; remainder of breast white, the adjoining chest-feathers black, tipped with white, or white with black edges; sides of body and flanks pale ashy brown; vent and under tail-coverts black, the latter with white tips; thighs black; axillaries white with black bases; under wing-coverts black, tipped with white; quills dusky below, pale brown along the edge of the inner web. Total length 6·8 inches, culmen 0·65, wing 3·05, tail 3·5, tarsus 0·85.

The Plate has been drawn from the specimen above described, which is represented in two positions, of the size of life.

[R. B. S.]

Gould, John and Sharpe, Richard Bowdler. 1884. "Rhipidura leucothorax, White-breasted Fantail Flycatcher [Pl. 27]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 2(XVIII), -. <https://doi.org/10.5962/p.322682>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229877>

DOI: <https://doi.org/10.5962/p.322682>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322682>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.