RHIPIDURA FUSCORUFA, *Sclater*.

RHIPIDURA FUSCO-RUFA, *Sclater*.

Timor-Laut Fantail Flycatcher.

Rhipidura fusco-rufa, Sclater, Proc. Zool. Soc. 1893, p. 197, pl. xxvii.

THIS Flycatcher appears to us, after a comparison with the "Keys to the species" of *Rhipidura* given in the British-Museum 'Catalogue of Birds,' and in Count Salvadori's 'Ornitologia della Papuasie,' to belong to a totally new section of the genus. Its rufous outer tail-feathers and a peculiar distribution of rufous and brown coloration about the bird render it quite distinct from any other known species. In the character of the tail it approaches somewhat the female of *R. brachyrhyncha*, Schlegel, from New Guinea; but there the resemblance ends, as the coloration of the rest of the plumage is totally different. Mr. Forbes sent a large series of this species from Larat, Moloe, and Loetoe; and it would appear to be common in the Tenimber Islands which he visited.

The following is a description of the type specimen:—

Adult male. General colour above dull chocolate-brown, the head and hind neck duller and more sooty brown; wing-coverts dusky brown, the least and median series tipped and the greater series edged with pale ferruginous; bastard wing, primary-coverts, and quills dusky black; the secondaries edged with pale ferruginous, more broadly on the inner ones; the four centre tail-feathers entirely dusky blackish, the next pair blackish, excepting for a shade of pale ferruginous towards the end of the feather, the three outermost entirely pale ferruginous with yellowish-white shafts; sides of face, lores, and ear-coverts dusky blackish; the fore part of the cheeks white, and a half-concealed white spot above the eye; throat and fore neck white, the sides of the latter ashy streaked with white; remainder of the under surface very pale ferruginous, deepening in tint on the thighs, vent, and under tail-coverts; under wing-coverts and axillaries richer ferruginous; quills dusky below, ashy rufous along the edge of the inner webs; "bill, legs, and feet black; iris dark brown" (*H. O. Forbes*). Total length 6·75 inches, culmen 0·75, wing 3·6, tail 3·5, tarsus 0·75.

The Plate represents an adult male of this species, in two positions, of the natural size.

[R. B. S.]

Gould, John and Sharpe, Richard Bowdler. 1883. "Rhipidura fuscorufa, Timor-Laut Fantail Flycatcher [Pl. 31]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 2(XV), –. <https://doi.org/10.5962/p.322686>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229877>

DOI: <https://doi.org/10.5962/p.322686>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322686>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.