GRALLINA BRUIJNII, *Salvad.*

GRALLINA BRUIJNI, *Salvad.*

Bruijn's Grallina.

Grallina bruijni, Salvad. Ann. Mus. Civ. Genova, vii. p. 929 (1875).—Sharpe, Cat. B. Brit. Mus. iii. p. 273 (1877).—Salvad. Ann. Mus. Civ. Genova, xv. p. 42, n. 4 (1879).—Salvad. Orn. Papuasias, ii. p. 191 (1881).—Sharpe, Journ. Linn. Soc., Zool. vol. xvi. p. 435 (1882).
Pomareopsis semiatra, Oustalet, Bull. Ass. Sci. de France, 1880, p. 173.

THE discovery of a species of *Grallina* in New Guinea is of great interest, as the genus was until recently considered to be entirely peculiar to Australia. In general appearance the New-Guinea bird differs considerably from its Australian relative, and is on the whole the handsomer bird of the two; but the same difference in sexes which characterizes the Pied *Grallina* is seen in the species from New Guinea.

The present species was discovered in the Arfak Mountains by the hunters employed by Mr. Bruijn, after whom it is named. More recently it has been sent from the mountains of South-eastern New Guinea by Mr. Goldie, who procured it from the Morocco district at the back of the Astrolabe range. Mr. Goldie states that the native name is *Tada*, and gives the following note:—"These birds are found flying about creeks and hopping about stones; they seem to feed on insects obtained there."

The following descriptions are taken from a pair of Mr. Goldie's birds in the British Museum.

Adult male. General colour above glossy blue-black, including the scapulars; all the wing-coverts pure white, forming a large shoulder-patch; bastard wing, primary-coverts, and quills black with glossy blue-black margins; lower back white; rump, upper tail-coverts, and basal half of the tail creamy buff, the terminal half of the latter black; feathers round the eye and ear-coverts blue-black, with a streak of white above the eye; the lower ear-coverts creamy white, as also the sides of the neck, forming a large patch; throat and entire under surface blue-black; lower flanks and thighs white; under tail-coverts creamy buff; on the sides of the lower back a patch of white feathers; under wing-coverts and axillaries blue-black; quills below black, the inner edges rather browner.

Total length 6·9 inches, culmen 0·9, wing 4·05, tail 3·05, tarsus 1·1.

Adult female. Resembles the male above, but differs as follows. Lores and broad eyebrow white; eyelid, feathers below the eye, cheeks, ear-coverts, and throat blue-black; remainder of under surface creamy white, deeper creamy buff on the flanks; abdomen and under tail-coverts, under wing-coverts, and axillaries white, and base of quills also, broader on the secondaries. Culmen 0·85 inch, wing 4·05, tail 2·9, tarsus 1·05.

The Plate represents an adult male and female of the natural size; and for the loan of the birds I am indebted to Mr. Edward Gerrard, junr. [R.B.S.]

Gould, John and Sharpe, Richard Bowdler. 1883. "Grallina bruijnii, Bruijn's Grallina [Pl. 11]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 3(XIV), –.

<https://doi.org/10.5962/p.322724>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229963>

DOI: <https://doi.org/10.5962/p.322724>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322724>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.