PACHYCEPHALA CHRISTOPHORI, *Tristram*.

PACHYCEPHALA CHRISTOPHORI, *Tristr.*

San-Christoval Thickhead.

Pachycephalus christophori, Tristram, Ibis, 1879, p. 441.

Pachycephala christophori, Salvad. Ibis, 1880, p. 131.—Id. Orn. Papuasie e delle Molucche, ii. p. 216 (1881).—

Ramsay, Proc. Linn. Soc. N. S. W. vi. p. 178 (1882), vii. p. 25 (1883).—Salvad. Ibis, 1884, p. 323.—

Tristr. t. c. p. 398.

Pachycephala astrolabi, juv., Gadow, Cat. Birds in Brit. Mus. ix. p. 200 (1883).

THIS species was discovered by Lieut. Richards in the island of San Christoval in the Solomon group, and was described by Canon Tristram. It is undoubtedly a distinct species, without the yellow collar round the hind neck or the black head of *P. astrolabi*, which is its nearest ally; it is also a much smaller bird than the last named, which is an inhabitant of Guadalcanar in the Solomon Archipelago. According to Mr. Ramsay, *P. christophori* has also been found in the island of Ugi by Messrs. Morton and Stephens.

Dr. Gadow considered that the present species was merely the young of *P. astrolabi*; but in this he was undoubtedly wrong, as has been shown by Canon Tristram and Count Salvadori.

The following is a description of the typical specimens in Canon Tristram's collection:—

Adult male. General colour dull yellowish green, a little lighter on the lower rump and upper tail-coverts; lesser wing-coverts a little yellower than the back; median and greater coverts dusky blackish, washed externally with olive-yellow; bastard-wing, primary-coverts, and quills also dusky blackish, edged with olive-yellow; the primaries margined with ashy grey; tail-feathers dusky blackish, the centre ones dull olive-yellowish towards the base, the remainder washed externally and tipped with dull olive-yellow; crown of head a little more dusky yellowish green than the back; the lores black, as also a frontal band; fore part of eyelid and feathers below the eye blackish; ear-coverts dull olive-brown with whitish shaft-lines; cheeks and throat bright yellow, followed by a broad black band across the lower throat, fore neck, and chest; breast and abdomen bright yellow, inclining to orange on the upper breast; sides of body and flanks also yellow, very slightly tinged with greenish; thighs and under tail-coverts paler yellow; under wing-coverts and axillaries white, washed with yellow on the edges; quills below dusky, white along the inner edge: "feet ash-colour, iris grey" (*G. E. Richards*). Total length 6·4 inches, culmen 0·8, wing 3·4, tail 2·3, tarsus 1·05.

Adult female. Brighter yellowish green than the male, the head like the back, but with no black forehead; ear-coverts like the crown, tinged with reddish brown, of which colour there is also a shade on the quills; upper tail-coverts and tail, cheeks and entire under surface of body bright yellow, paler on the throat, slightly washed with greenish on the fore neck, chest, sides of body, and flanks, and tinged with orange on the breast; no black pectoral collar; under wing-coverts and axillaries yellow; quills dusky below, reddish along the edge of the inner web. Total length 5·65 inches, culmen 0·8, wing 3·25, tail 2·2, tarsus 0·95.

The young male resembles the old female, but has a few black feathers on the breast, indicating the appearance of the pectoral band.

The Plate illustrates the male and female of the natural size, the figures having been drawn from the type specimens kindly lent to us by Canon Tristram.

[R. B. S.]

Gould, John and Sharpe, Richard Bowdler. 1887. "Pachycephala christophori, San-Christoval Thickhead [Pl. 23]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 3(XXIII), -. <https://doi.org/10.5962/p.322736>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229963>

DOI: <https://doi.org/10.5962/p.322736>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322736>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.