MELIRRHOPHETES LEUCOSTEPHES, Meyer.

MELIRRHOPHETES LEUCOSTEPHES, *Meyer.*

White-faced Honey-eater.

Melirrhophetes leucostephes, A. B. Meyer, Sitzungsab. der k.-k. Akad. d. Wissenschaften zu Wien, lxx. Juni 1874, p. 111.—T. Salvadori, Ann. del Mus. Civ. di Genova, vol. vii. 1875, p. 776.

THIS large and attractive Honey-eater must form a conspicuous feature among the flowering trees of New Guinea, much as do the large species of this family so numerous spread over Australia; at least we may surmise this, as not even a few words have been placed on record by its discoverer Dr. Meyer, or by any other person. These meagre descriptions, however, testify to the rarity of the subject, and point out to future travellers over New Guinea how very desirable it would be for them to jot down any such particulars before they leave the forests, or at least as soon after as an opportunity may occur. All I can now say is that my artist, Mr. Hart, has made as faithful a portrait of this new bird as possible, to which I add Dr. Meyer's description very kindly forwarded to me in a letter.

"Feathers of forehead and those which encircle the naked skin surrounding the eye white; throat, crown, and hinder part of head, nape, ears, and a patch which borders the front of the eye black; naked eye-skin and lengthened skin-fold of the mouth yellowish; caruncles of the throat orange; back brownish, feathers of the upper part edged with white and brownish white; under surface black, with some white markings on the breast and abdomen; upper part of wings bright olivaceous; under surface greyish, with light-brown edges on the bases of the inner webs; under wing-coverts blackish, with light brown intermixed; upper part of tail brown, each feather having a lighter edge on the outer web; bill bluish grey, lighter at the tip; feet and tarsi blackish grey, soles of feet lighter."

Total length 11 inches, bill $1\frac{5}{8}$, wings 5, tail 5, tarsi $1\frac{1}{4}$.

According to Salvadori the sexes are alike in colour.

Hab. Arfak Mountains in New Guinea.

The figure in the Plate is of the natural size.

Gould, John and Sharpe, Richard Bowdler. 1877. "Melirrhophetes leucostephes, White-faced Honey-eater [Pl. 33]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 3(IV), -. <https://doi.org/10.5962/p.322746>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229963>

DOI: <https://doi.org/10.5962/p.322746>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322746>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.