MELIARCHUS SCLATERI.

MELIARCHUS SCLATERI.

Sclater's Honey-eater.

Philemon vulturinus (nec Reichenb.), Sclater, Proc. Zool. Soc. 1869, pp. 120, 124.

Philemon sclateri, Gray, Ann. and Mag. Nat. Hist. 1870, v. p. 327.—Id. Cruise of the Curaçoa, Birds, p. 362, pl. v. (1873).—Tristr. Ibis, 1879, p. 439.—Gadow, Cat. Birds in Brit. Mus. ix. p. 279 (1884).

Meliarchus sclateri, Salvad. Ann. Mus. Civic. Genov. xvi. p. 75 (1880).—Id. Orn. Papuasie e delle Molucche, ii. p. 322 (1881).

THIS fine species of Honey-eater appears to be confined to the Solomon Islands, and is probably found only on the island of San Christoval. Here it was procured by the late Mr. Brenchley, and Lieut. Richards also met with it at Makira Harbour, in the same island. Very few specimens exist in European collections, and nothing has been recorded of its habits.

We follow Count Salvadori in referring the present species to a distinct genus, as it appears to us to be by no means a true *Philemon*, excepting under the very elastic definition which Dr. Gadow gives to the latter genus.

The following description is taken from a specimen lent to us by Mr. E. P. Ramsay:—

Adult male. General colour above dull olive on the back and mantle, dull reddish brown on the body, rump, and upper tail-coverts; wing-coverts dark brown, edged with olive; bastard-wing blackish brown; primary-coverts and quills dusky brown, edged with lighter olive-yellow, except on the inner secondaries; tail-feathers reddish brown with olive margins; crown of head pale yellow, streaked with black, the hind neck with dusky; feathers round the eye and eyebrow and the upper part of the ear-coverts yellowish white, with a black spot on the lores; cheeks and lower part of ear-coverts black, streaked with pale yellow edges to the feathers; throat pale ashy grey; breast pale olive-yellow, with dusky centres to the feathers; lower breast and abdomen dull ashy; sides of body and flanks reddish brown washed with olive; thighs and under tail-coverts reddish brown, edged with olive-yellow; axillaries and under wing-coverts dark ashy, washed with olive; quills dusky below, ashy along the inner web: "bill yellowish; feet ash-colour; iris brown" (*G. E. Richards*). Total length 9·5 inches, culmen 1·5, wing 4·5, tail 4·25, tarsus 1·35.

The Plate represents an adult bird of the natural size, and is drawn from the above-mentioned specimen lent by Mr. Ramsay. It was procured at Makira Harbour by Lieut. Richards.

[R. B. S.]

Gould, John and Sharpe, Richard Bowdler. 1886. "Meliaichus sclateri, Sclater's Honey-eater [Pl. 52]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 3(XXI), –.
<https://doi.org/10.5962/p.322765>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229963>

DOI: <https://doi.org/10.5962/p.322765>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322765>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.