



MYZOMELA WAKOLOENSIS, Forbes.

W. Hart del. et lith.

Mintern. Bros. imp.

MYZOMELA WAKOLOENSIS, *H. O. Forbes.*

Lake Wakolo Honey-eater.

Myzomela wakoloensis, Forbes, Proc. Zool. Soc. 1883, p. 116.

THIS species of Honey-sucker was discovered by Mr. H. O. Forbes during his expedition to the Malay Archipelago, in the island of Bourou. Even in that island he met with it in one place only, Lake Wakolo, a locality to which few Europeans have ever penetrated before.

In general appearance it resembles *M. cruentata* of Meyer (figured in this work) more than any other species of *Myzomela*, but it is quite distinct specifically. The tint of the red colouring is quite different, and the black wings with their olive margins to the quills, as well as the absence of red on the tail, are good specific characters.

Adult male. General colour above scarlet vermilion; the feathers of the back and scapulars black at base, with broad ends of scarlet vermilion; rump and upper tail-coverts like the back; lesser and median wing-coverts black, the innermost of the latter tipped with scarlet; greater coverts and quills blackish, edged narrowly with olive-yellowish; bastard wing and primary-coverts blackish; tail-feathers blue-black; head like the back; lores black; sides of face, ear-coverts, cheeks, throat, and under surface of body bright scarlet, the feathers of the latter part with dusky bases; vent and under tail-coverts pale olive-yellowish, with dusky centres and somewhat tinged with red; under wing-coverts and axillaries white, with a yellowish tinge along the edge of the wing; quills dusky below, ashy white along the edge of the inner web, with dusky centres; axillaries and under wing-coverts white; the quills dusky below, whitish along the edge of the inner web; "bill black, the lower mandible yellow at base; legs and feet sooty grey; iris dark brown" (*H. O. Forbes*). Total length 3.6 inches, culmen 0.5, wing 2.05, tail 1.3, tarsus 0.65.

Mr. Forbes unfortunately did not procure an adult female, but he managed to get the immature male, which will doubtless give us some idea of the plumage of the old hen bird.

The young male is brown above, with pale streaks on the forehead; wing-coverts dusky brown, edged with paler or white brown; quills dusky, edged with an olive fringe, the inner secondaries margined with pale brown, like the coverts; tail-feathers dusky brown, with olive-brown edges; lores brown, tinged with red; ear-coverts and cheeks ochreous brown, with a reddish tinge; throat pale yellowish, with a distinct wash of red on the chin; lower throat and fore neck more ashy; abdomen yellowish white, the breast and sides of the body browner; under tail-coverts ruddy brown. Total length 3.7 inches, culmen 0.6, wing 2.1, tail 1.2, tarsus 0.6.

[R. B. S.]



BHL

Biodiversity Heritage Library

Gould, John and Sharpe, Richard Bowdler. 1884. "Myzomela wakoloensis, Lake Wakolo Honey-eater [Pl. 64]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 3(XVIII), -. <https://doi.org/10.5962/p.322777>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229963>

DOI: <https://doi.org/10.5962/p.322777>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322777>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.