



ÆDISTOMA PYGMÆUM, Salvad.

## ÆDISTOMA PYGMÆUM, *Salvad.*

### Pigmy Honey-eater.

*Ædistoma pygmæum*, *Salvad.* Ann. Mus. Civic. Genov. vii. p. 952 (1875).—Id. op. cit. xvi. p. 74 (1880).—Id. Orn. Papuasie e delle Molucche, ii. p. 312 (1881).—Gadow, Cat. Birds in Brit. Mus. ix. p. 293 (1884).

---

For an opportunity of figuring this curious little bird we are indebted to the kindness of the Marquis Doria, who was so good as to send us one of the typical specimens from the Genoa Museum. Its small size, slender bill, and sober colouring recall the species of *Melilestes*, and, as in the latter genus, *Ædistoma* has very pronounced silky tufts on the flanks. Count Salvadori considers *Ædistoma* to be very nearly allied to *Melilestes*, but to be distinguished principally by the form of the bill; and this will be easily discernible on a comparison of the Plates given in the present work. We agree with him that the genera *Melilestes* and *Ædistoma*, along with *Glycichæra*, form a natural section of the *Meliphagidæ*, though they show a certain connection with the Sun-birds of the genera *Arachnothera* and *Anthothreptes*.

This species, as far as is known, is confined to the Arfak Mountains in North-western New Guinea.

The following description is taken from the typical specimen belonging to the Genoa Museum:—

*Adult female.* General colour above dark olive-green, slightly greyer on the head and more distinctly yellowish towards the rump, the feathers of which are long and silky and completely hide the upper tail-coverts; wing-coverts below the back, bastard-wing, primary-coverts, and quills ashy brown, edged with ashy, the secondaries with olive-yellow, lighter than the back; centre tail-feathers olive-green, the remainder dusky edged with olive-greenish; in front of the eyes a dusky spot; eyelid whitish; sides of face and ear-coverts pale ashy, whiter below the eye; throat whitish, with a slight greenish tinge; remainder of under surface pale sulphur-yellow, olive on the fore neck and chest, yellowish white on the centre of the breast and abdomen; axillaries pale sulphur-yellow like the sides of the body; quills dusky below, ashy along the inner web: "bill dusky, the base of the lower mandible dusky below; feet lead-colour" (*Beccari*). Total length 2·8 inches, culmen 0·5, wing 1·9, tail 0·8, tarsus 0·5.

The male, according to Count Salvadori, does not differ from the female in colour.

The figures in the Plate represent an adult bird in two positions, drawn from the typical specimen lent to us by the Marquis Doria.

[R. B. S.]



Gould, John and Sharpe, Richard Bowdler. 1885. "Ædistoma pygmæum, Pigmy Honey-eater [Pl. 8]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 4(XX), –.  
<https://doi.org/10.5962/p.322793>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/230755>

**DOI:** <https://doi.org/10.5962/p.322793>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/322793>

#### **Holding Institution**

Smithsonian Libraries and Archives

#### **Sponsored by**

Biodiversity Heritage Library

#### **Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.