



ÆGOTHELES WALLACHI, Gray.

W. Hart del et lith.

Mintern Bros. imp.



## ÆGOTHELES WALLACII, *Gray.*

### Wallace's Goatsucker.

- Ægotheles wallacii*, Gray, Proc. Zool. Soc. 1859, p. 154.—Id. Proc. Zool. Soc. 1861, p. 433.—Finsch, Neu-Guinea, p. 162 (1865).—Schlegel, Nederl. Tijdschr. Dierk. iii. p. 340 (1866).—Gray, Hand-list Birds, i. p. 55, no. 603 (1869).—Sclater, Proc. Zool. Soc. 1873, p. 696.—Meyer, Sitz. k. Akad. Wien, lxix. p. 75 (1874).—Salvad. Ann. Mus. Civic. Genov. x. p. 310 (1877).—Id. Proc. Zool. Soc. 1878, p. 94.—Id. Orn. Papuasie e delle Molucche, ii. p. 526 (1880).—Id. Report Voy. H.M.S. 'Challenger,' p. 77 (1882).—Meyer in Madarász, Zeitschr. ges. Orn. i. p. 278, pl. xvii. fig. 4 (1884).—Guillemard, Proc. Zool. Soc. 1885, p. 630.
- ? *Caprimulgus brachyurus*, Schl. Nederl. Tijdschr. Dierk. iii. p. 340 (1866, ex Rosenb. MSS.).—Rosenb. Nat. Tijdschr. Nederl. Ind. xxix. p. 143 (1867).—Id. Reis. naar Zuidoostereil. p. 37 (1867).

---

THIS interesting Goatsucker was discovered by Mr. A. R. Wallace during his travels in the east, at Dorei in New Guinea, and the type specimen is in the British Museum. It has since been met with in the same locality by Mr. Bruijn's hunters, as well as on the Arfak Mountains by Dr. A. B. Meyer, and in Atam by Signor D'Albertis. It appears to represent in North-western New Guinea the Australian *Ægotheles novæ-hollandiæ*, and it is replaced in South-eastern New Guinea by *Æ. bennetti*. It differs from the latter, and consequently also from *Æ. novæ-hollandiæ*, in its dark coloration, and in having the fore part of the crown varied with rufous, and in exhibiting some whitish-red spots on the scapular feathers.

A Goatsucker from the Aru Islands has been described by Baron von Rosenberg as a distinct species under the name of *Caprimulgus brachyurus*. Count Salvadori, who has examined the type in the Leyden Museum, is of opinion that it is a young bird of the genus *Ægotheles*, with an imperfectly developed tail, probably referable to *Æ. wallacii*; but he has also seen a second specimen from the Aru Islands collected during the 'Challenger' expedition, and he believes that the Aru bird is probably distinct, by reason of its smaller dimensions and more minute vermiculations. We have examined the last-named specimen and find that it fully bears out Count Salvadori's opinion; but it will be better to wait for a larger series of specimens before venturing to separate the Aru bird specifically, as Goatsuckers vary so much in the intensity of their coloration.

Dr. Meyer has recently received a specimen from the same group of islands, where the bird is called by the natives "Tatar faffu." The iris was greyish brown and the feet dark flesh-colour. The egg, which is figured by Dr. Meyer, is cream-coloured, with scribblings of dusky greyish.

We do not give a detailed description of this species, the characters having been well pointed out above.

The Plate represents two adult birds of the natural size, the figures having been drawn from a specimen lent to us by Dr. Guillemard.

[R. B. S.]



Gould, John and Sharpe, Richard Bowdler. 1886. "Ægotheles wallacii, Wallace's Goatsucker [Pl. 39]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 4(XXI), –.

<https://doi.org/10.5962/p.322824>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/230755>

**DOI:** <https://doi.org/10.5962/p.322824>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/322824>

#### **Holding Institution**

Smithsonian Libraries and Archives

#### **Sponsored by**

Biodiversity Heritage Library

#### **Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.