CEYX SOLITARIA,

CEYX SOLITARIA.

Solitary Kingfisher.

- Ceyx meninting*, Less. Voy. Coquille, Zool. i. p. 691 (1826).—Cuv. Règne An. i. p. 444 (1829).—Less. Traité, p. 241 (1831).—Pucher. Rev. et Mag. 1861, p. 345.
- Ceyx solitaria*, Temm. Pl. Col. 595. fig. 2 (1836).—Gray, P. Z. S. 1858, p. 172; 1859, p. 155; 1861, p. 433.—Sharpe, P. Z. S. 1868, p. 271.—Id. Monogr. Alced. pl. 38 (1868).—Salvad. Atti R. Accad. Torino, iv. p. 474 (1869).—Gray, Handl. B. i. p. 95 (1869).—Salvad. Ann. Mus. Civ. Genoa, vii. p. 765 (1875).
- Alcyone solitaria*, Gray, Gen. B. i. p. 82 (1847).—Bp. Consp. i. p. 158 (1850).—Reich. Handb. Alced. p. 7, taf. ccxcviii. fig. 3067 (1851).—Bp. Consp. Vol. Anis. p. 10, sp. 358 (1854).—Rosenb. J. f. O. 1864, p. 118.
- Alcedo solitaria*, Schl. M. P. B. Alcedines, p. 17 (1863).—Id. Voy. Nederl. Ind. Alced. pp. 12, 48, pl. 3. fig. 5 (1864).—Id. Revue Alced. p. 9 (1874).
-

THIS little three-toed Kingfisher is an inhabitant of New Guinea and the neighbouring islands. The original type came from the Bay of Lobo; and the Leiden Museum has specimens from Sorong, Dorey, and Andai, collected by the late Dr. Bernstein and by Von Rosenberg. In the island of Salvatty, Bernstein collected examples at Kalwal and Sailolo; and it has been sent from the island of Batanta by Von Rosenberg: in the last-named locality Dr. Beccari also met with it. Mr. Hoedt is our authority for the occurrence of the bird in Mysol, five individuals being in the Leiden Museum from Waaigama and Kasim. In his original 'Catalogue of the Kingfishers of the Leiden Museum,' Professor Schlegel recorded a specimen from Ceram; but Mr. Wallace at the time doubted the occurrence of the species in that island, and it now seems that the specimen in question was a Mysol one. Lastly, it occurs in the Aru Islands, whence we saw a large series in the collection recently formed there by Mr. Cockerell. Von Rosenberg's localities are Wokam and Wonoumbai. Mr. Cockerell is likewise our authority for the occurrence of the present species in Australia, as specimens were in his last collection from Cape York, and he assures me that Mr. Jardine had obtained them in the neighbourhood of Somerset. Beyond this, I have never in all my experience heard of the species nor seen a single example from the Australian continent.

All that is at present written concerning its economy is contained in the following brief note, communicated by Mr. Wallace to Mr. Sharpe's 'Monograph.' He observes that "the stomachs of those he killed contained the remains of water-beetles and other insects."

The present species is distinguished first of all by having only three toes, a peculiarity of the genera *Ceyx* and *Alcyone*; and from all the members of the latter it differs by its black bill and yellow belly.

In the Plate the bird is represented of the natural size, drawn from an Aru-Island specimen in my collection.

Gould, John and Sharpe, Richard Bowdler. 1876. "Ceyx solitaria, Solitary Kingfisher [Pl. 45]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 4(III), –.
<https://doi.org/10.5962/p.322830>.

View This Item Online: <https://www.biodiversitylibrary.org/item/230755>

DOI: <https://doi.org/10.5962/p.322830>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322830>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.