MONTIFRINGILLA RUFICOLLIS. W.T. Blanford

MONTIFRINGILLA RUFICOLLIS, *Blanford.*

Red-necked Mountain-Finch.

Montifringilla ruficollis, Blanf., J. A. S. B., xli. p. 67 (1872).

THE birds composing the genus *Montifringilla* are truly, as their name implies, inhabitants of the mountains, being found in the elevated ranges from the Alps of Southern Europe to the Himalayas. The one which I now figure seems to be the smallest of the genus; but it is at the same time the prettiest and most conspicuously marked; and I consider Mr. Blanford very fortunate in crowning the difficult journey which he made in company with Captain Elwes to the Thibetan frontier, by the discovery of such an interesting species.

"This bird," writes Mr. Blanford, "differs widely from every described form. In structure it is closely allied to *M. nivalis*, and differs from *Leucosticte* in its shorter wings and tail. The plumage may become whiter in winter.

"I met with one flock of this new Finch near Kangra Lama pass, and with another at Prálung, both places inhabited by purely Thibetan forms, and at elevations of 15000 to 16000 feet. It is probably common in Thibet. Dr. Stoliczka is of opinion that it is the same as the undescribed *Montifringilla* found by him in Ladak (J. A. S. B. 1868, pt. ii. p. 62), but of which his specimen is not now in Calcutta."

Subjoined is Mr. Blanford's original description.

"*Male*.—Forehead whitish, passing into the rather pale umber-brown of the head; supercilia white; lores and a line from them passing under the eye black, this line is continued posteriorly over the ear-coverts, and its colour changes to dark ferruginous; back umber-brown, with broad central dusky streaks to the feathers; rump more ferruginous; wings brown, the first primary with the outer web white, the others with fulvous outer margins, all with white internal edges and a broad white wing-band, only visible on the expanded wing, formed by a large spot on the inner webs of all the primaries except the first four, and the whole basal portion of the inner web of the remaining quills except the last three, which have broad fulvous borders; smaller wing-coverts mostly white, angle of wing greyish; upper tail-coverts very long, pale umber with a fulvous tinge; central tail-feathers and the tips of the remainder for about half an inch dark brown, with fulvous margins; basal portion of all the tail-feathers except the central pair pale ashy, with some white, on both inner and outer webs, between the grey portion and the brown tips, increasing in quantity on the outer feathers and running up the external web, which is entirely white in the outermost pair; sides of the head below the black eye-streak, chin, and throat white, with two black lines, one from each side of the base of the lower mandible, running backwards and diverging; ear-coverts bright ferruginous; sides of the neck the same, but a little paler, the rufous tint forming a semi-collar, only interrupted for a very narrow space in front; remainder of the lower parts white, with an isabelline tinge. Iris reddish brown, bill and legs black.

"The female wants the whitish forehead and the rufescent tinge on the rump; the demi-collar is brown posteriorly, and, in the only specimen obtained, it is continuous round the front of the neck.

	Wing.	Tail.	Tarsus.	Hind claw.	Bill from forehead.	Bill from gape.
"Male	3.63	3.23	0.85	0.36	0.42	0.53.
Female	3.56	2.1	0.87	0.32	0.43	0.54."

The figures in the accompanying Plate were drawn from the types, for the loan of which I have to thank Mr. Blanford. They are of the size of life.

Gould, John. 1875. "Red-necked Mountain-Finch, *Montifringilla ruficollis*,
Blanford. [Pl. 5]." *The Birds of Asia* 5(XXVII), –.
<https://doi.org/10.5962/p.322977>.

View This Item Online: <https://www.biodiversitylibrary.org/item/121124>

DOI: <https://doi.org/10.5962/p.322977>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322977>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.