LIOPTILA ANNECTANS, Blyth.

W. Hart del. et lith.

Walter imp.

LIOPTILA ANNECTENS, *Blyth.*

Slender-billed Chat-Thrush.

Leioptila annectans, Blyth, Journ. As. Soc. Beng. xvi. p. 450 (1847).—Id. Cat. B. Mus. As. Soc. p. 337 (1849).—Jerd. B. India, ii. p. 248 (1863).—Godwin-Austen, J. As. Soc. Beng. xxxix. p. 109 (1870).
Cutia annectans, Gray, Hand-l. B. i. p. 314, no. 4690 (1869).
Lioptila annectans, Hume, Str. F. 1877, p. 110.—Id. Str. F. 1879, p. 104.

THE present species has generally been considered a connecting link between the genera *Cutia* and *Liothrix*; and its name "*annectans*" shows that Mr. Blyth regarded it as joining these two groups together. Although its exact relations were believed by him to be with *Sibia* and *Yuhina*, I myself think that its affinities are more with the Chat-Thrushes or *Thamnobiæ*, and that it is an eastern representative of the African genus *Cossypha*.

The range of the present species extends from Sikkim to the Khasia hills, and in the Karen and Tenasserim hills it is replaced by a slightly darker race which was called *L. saturata* by the late Lord Tweeddale. Of the latter bird Mr. Davison gives the following account:—"I only found this bird at Mooleyit, quite near the top, usually in pairs, sometimes singly. I found it generally about the large trees surrounding the 'sakans' or camping-grounds—strange to say, climbing about the trunk and branches much after the manner of a Nuthatch. I have also seen it hunting about the leaves and smaller branches of the tree-tops. Those I killed had eaten only insects. I never remember to have heard it utter any note. It was very rare even where it did occur, and I only saw some six or seven."

The following description is of a specimen kindly lent to me by Captain Wardlaw Ramsay.

Adult. General colour above rich orange-chestnut from the lower mantle downwards; upper scapulars black, the lower ones chestnut or bordered with black; head, nape, and upper mantle black, streaked conspicuously with white, the feathers of the hind neck and mantle white on the inner web; lesser and median wing-coverts black, the feathers edged with ashy-grey; greater series black tipped with orange-chestnut, forming a wing-bar; bastard wing and primary-coverts black; quills black, margined with pale lavender-grey, the inner secondaries tipped with white, the innermost orange-chestnut near the base of the outer web; tail-feathers black, the centre ones fringed with ashy whitish, the ends pure white on the other feathers, increasing in extent towards the outermost; lores and sides of face and ear-coverts black; cheeks and under surface of body white, the flanks and under tail-coverts orange-buff; thighs ashy with whitish edgings; under wing-coverts and axillaries pure white; quills blackish below, white along the inner margin; "bill black, the base of the lower mandible yellow; legs pale fleshy brown; iris brown" (JERDON). Total length 6·5 inches, culmen 0·65, wing 3·1, tail 3·1, tarsus 0·95.

The figures in the Plate represent two specimens, of the size of life, from the above-named skins lent me by Captain Wardlaw Ramsay.

[R. B. S.]

Gould, John. 1883. "Slender-billed Chat-Thrush, *Leioptila annectens*, Blyth. [Pl. 30]." *The Birds of Asia* 3(XXXIV), -. <https://doi.org/10.5962/p.322983>.

View This Item Online: <https://www.biodiversitylibrary.org/item/118635>

DOI: <https://doi.org/10.5962/p.322983>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322983>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.