ALCEDO BENGALENSIS, *Cmel.*

Stoddard and McRicker, del. et lith.

Walter & Cohen, Imp.

ALCEDO BENGALENSIS, Gmel.

Indian Kingfisher.

- Alcedo bengalensis*, Gmel. Edit. Linn. Syst. Nat., tom. i. p. 450.—Bonn. et Vieill. Ency. Méth., Orn., part i. p. 291.
—Shaw, Gen. Zool., vol. viii. part i. p. 102.—Id. Steph. Cont., vol. xiii. part ii. p. 103.—Sykes, Proc. of Comm. of Sci. and Corr. of Zool. Soc., part ii. p. 84.—McClell. Proc. Zool. Soc., part vii. p. 156.
—Jerd. Madras Journ. of Lit. and Sci., vol. xi. p. 231.—Pears. Journ. Asiat. Soc. Beng., vol. x. p. 636.—Gray and Mitch. Gen. of Birds, vol. i. p. 81.—Gray, Cat. of Spec. of Birds in Brit. Mus., part ii. sec. i. p. 63.—Id. Cat. of Spec. and Draw. of Mamm. and Birds pres. to Brit. Mus. by B. H. Hodgson, Esq., p. 57.—Blyth, Journ. of Asiat. Soc. Beng., vol. xv. p. 11.—Id. Cat. of Birds in Mus. Asiat. Soc. Calcutta, p. 49.—Bonap. Consp. Gen. Av., tom. i. p. 158.—Dr. F. B. Hamilton's MS., vol. i. p. 27.—Cass. Cat. of Halc. in Mus. Acad. Sci. Philad., p. 3.—Horsf. and Moore, Cat. of Birds in Mus. East Ind. Comp., vol. i. p. 129.—Kittl. Kupf. Vög., t. 29. fig. 2.—Cab. et Hein. Mus. Hein., Theil ii. p. 144.—Irby in Ibis, vol. iii. p. 228.—Swinh. in Ibis, vol. iii. pp. 31 & 328.—Bonap. Consp. Vol. Anisod. in Ateneo Italiano, no. 11, Agosto 1854.—Sieb., Temm. and Schleg. Faun. Jap., tab. xxxviii.
Alcedo hispidioides, Less. Compl. Buff., tom. ix. p. 345.
— *hispidula*, Cat. of Rivoli Coll., pp. 2, 3.
— *Bengalensis*, var. *Indica*, Reichenb. Handb., tom. i. p. 3, 3a, tab. 393. fig. 3047.
Ispida Bengalensis, Klein, Av., p. 34, no. 2.—Briss. Orn., tom. iv. p. 475.—Buff. Hist. Nat. des Ois., tom. xiii. p. 297.
— *minor*, Briss. Orn., tom. iv. p. 477.
Alcedo ispida Bengalensis, Sieb., Temm. et Schleg. Faun. Jap., p. 76.
— *japonica*, Bonap. Consp. Vol. Anisod. in Ateneo Italiano, no. 11, Agosto 1854.
The Little Indian Kingfisher, Edw. Nat. Hist. of Birds, vol. i. pl. 11.
Chota Kilkila of the Hindoos (Jerdon).
Chota match-ranga of the Musselmans (Dr. F. B. Hamilton).
Seeta koona, Malwa (Boys).
Match-ranga of the Bengalese (Blyth and Hamilton).
Rajah whodan, Malay (Blyth).
Mal-pilli hudua (*piuchi* or "small"), in Ceylon (Blyth).
Tow-yü-long of the Cantonese (Swinhoe).

If I were writing the history of the British Kingfisher (*Alcedo ispida*), I should state that the middle portion of Europe was probably the centre of its area of range, and that from thence it radiated westward to the British Islands, northwards to Norway and Sweden, southward to Morocco, and eastward to Asia Minor, and that, in the latter country, it inosculates with another species, the *Alcedo Bengalensis*, the subject of the present memoir. Here, then, the two species meet; but this appears to be the western limit of the Indian bird, its true habitat being Scinde, Bombay, and the whole of the Peninsula of India, wherever situations suited to its habits occur; thence it advances eastward to China and Japan, and southward through the Indian Islands to Timor, according to the authors of the 'Fauna Japonica.' Specimens from these widely extended localities present certain trifling differences both in size and colour, but not to the extent observable between the Indian and European species,—the *Alcedo Bengalensis* differing from the *A. ispida* not only in being considerably smaller in size, but more brilliantly coloured, and in the more decided character of the markings of its head and wing-coverts, these features being so apparent that the experienced ornithologist would have no difficulty in deciding at a glance to which country a specimen belonged, were his opinion required. I have said this much on the subject in order to prove the right of Asia to claim a true and distinct species of this form among her avi-fauna; but as the subject has been most carefully investigated by my friend Schlegel in the 'Fauna Japonica,' it is only justice to him to transcribe his remarks:—

"*The common Kingfisher of Bengal.*—This bird, which takes the place of the ordinary Kingfisher throughout a great part of southern and eastern Asia, does not appear to differ from that species except in its smaller size, and occasionally by the depth of the tints of its plumage. Some modern authors, regarding these differences of too little importance to serve as a means of specific distinction, have considered these birds as belonging to the same species, and have not even admitted the existence of several races of this bird. Putting on one side this question of nomenclature, we shall limit ourselves to record the observations we

have made upon these different birds, which offer to one another a striking resemblance with respect to the distribution of their tints. The ordinary Kingfisher (*Alcedo ispida* of authors—a species known throughout the whole world) is of the following proportions:—Length of wings, 2 inches 10 lines; length of beak, from the front, 1 inch 4 lines; breadth of beak, near the front, $3\frac{1}{4}$ lines; length of tail, 1 inch 4 lines; tarsi, $3\frac{3}{4}$ lines.

“The ordinary Kingfisher of Bengal (*Alcedo Bengalensis*, Gmelin, figured by Edwards, pl. 11, and by Kittlitz, Kupfertafeln, pl. 29, fig. 1) may be constantly distinguished from the common species by its less robust size and less squat form, although its beak may be a little longer than in the former. This little race has been observed at Bengal; our travellers have obtained a great number of individuals from Japan and Timor; and Kittlitz says that it is common in the Isle of Luçon, and that its habits are identically the same as those of the European race. We have before us a complete series of individuals received from Bengal, from Japan, and from Timor; these all exhibit slight but perceptible differences.

“Those from Bengal present the same gradation of tints as those of the common Kingfisher of Europe; but their wings do not exceed 2 inches 7 lines in length, the tail is 1 inch 4 lines, and the beak 1 inch 5 to 6 lines long.

“The individuals from Japan still more closely resemble those from Bengal with regard to the gradation of their tints; but they have the wings constantly a little longer, in that they measure 2 inches $8\frac{1}{2}$ lines long.

“Lastly, the individuals from Timor resemble those from Japan in the length of their wings; but they differ from them, as well as from those of Bengal, in the tints of the upper part of the bird, which are much more lively, of an almost perfect blue, and very little inclining to green, as seen in individuals from Japan and Bengal, as well as in the common European species.”

I may here remark that the late Prince Charles Bonaparte regarded the Japanese bird as distinct, and that he therefore designated it *Alcedo Japonica* (see his paper in the ‘Ateneo Italiano’ for August 1854).

Colonel Sykes says, “This species affects brooks; it is never seen in gardens.”

Mr. Pearson informs us that “It is frequently seen seated on a stick standing upright in a paddy-field, watching for small fishes in the shallow water.”

From Dr. Francis Buchanan Hamilton’s MS. we learn that “It builds its nest in holes, which it excavates in abrupt banks and mud walls.”

The following brief notice of this bird occurs in the notes of the late Captain Boys:—“Frequents the mountain-streams and shady spots; is more frequently seen on the branch of a tree overhanging the water than on any projecting stone or stump in it. It flies with great rapidity and ease, generally very near the water. It is abundant near the cantonments at Nibon, each rivulet having a pair in about every two miles. It constantly remains at the same spot, unless forced to remove by necessity and the want of food. The sexes are alike in plumage; but that of the female is of a somewhat duller hue.”

Mr. Swinhoe states that it is sometimes seen, but is not common, between Takoo and Peking, in North China.

The male has the crown and nape and a stripe from the angle of the lower mandible deep-bluish green, transversely barred with bright verditer-green; plumage of the upper surface and wings bluish green; each of the wing-coverts with a spot of verditer-green at the tip; back and upper tail-coverts beautiful verditer-green; lores, space below the eye, ear-coverts, under wing-coverts, and under surface of the body reddish chestnut; behind the ear-coverts a streak of silky white; primaries and secondaries brown, margined externally with green; tail deep blue; irides brown; feet orange; bill black, lighter at the base of the under mandible.

The female is very similar; but the markings of the head and wings are not so fine, the base of the under mandible is lighter, and her general hue is duller than that of her mate.

In the immature bird the general colouring is less brilliant, and the upper surface and breast are clouded with olive-brown.

The figures represent two fully adults, of the natural size.

Gould, John. 1862. "Indian Kingfisher, *Alcedo bengalensis*, Gmel. [Pl. 53]." *The Birds of Asia* 1(XIV), -. <https://doi.org/10.5962/p.323034>.

View This Item Online: <https://www.biodiversitylibrary.org/item/115342>

DOI: <https://doi.org/10.5962/p.323034>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323034>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.