PITTA CUCULLATA, Hartlaub.

J. Gould & W. Hart, del. et lith.

Walter Lloyd.

PITTA CUCULLATA, *Hartl.*

Hooded Pitta.

- Pitta cucullata*, Hartl. Rev. et Mag. de Zool. 1833, p. 65.—Gray, Gen. B. i. p. 213 (1846).—Blyth, Cat. B. Mus. A. S. B. p. 157 (1849).—Moore, P. Z. S. 1854, p. 274.—Horsf. & Moore, Cat. B. Mus. E.I. Co. ii. p. 399 (1856).—Cass. U.S. Expl. Exp. Birds, p. 437 (1858).—Jerd. B. Ind. i. p. 504 (1862).—Gray, Hand-l. B. p. 295 (1869).
— *nigricollis*, Blyth, J. A. S. B. xii. p. 960 (1843).
— *rhodogaster*, Hodgs. J. A. S. B. xii. p. 961 (1843).
— *malaccensis*, Müll. & Schleg. Verh. Natuurl. Gesch. Zool. p. 19 (1844).
— *coronata*, Peale, U.S. Expl. Exp. Birds, p. 891 (1848).
Brachyurus cucullatus, Bp. Consp. i. p. 255 (1850).—Elliot, Monogr. Pittidæ, pl. xxviii. (1863).—Hume, Str. F. 1875, p. 109.
Melanopitta cucullata, Bp. Consp. Volucr. Anis. p. 7 (1854).—Hume, S. F. 1874, p. 475.—Id. Nests & Eggs Ind. B. p. 225 (1875).

THE close connexion which exists between the avifauna of the Eastern Himalayas and that of the Malayan peninsula is illustrated by the occurrence of several peculiar forms; but by none is the fact better exemplified than by the subject of the present article. Its range extends from Nepal and Sikhim, through Assam, Burmah, and Tenasserim to the Malayan peninsula; and specimens from all these localities are identical.

The following observation is extracted from Mr. Hume's 'Nests and Eggs of Indian Birds':—

"According to Mr. Hodgson's notes and drawings, the Hooded Pitta breeds in the central regions of Nepal and about Darjeeling in April and May. They build a large globular nest, one of which measured nearly 6·75 inches externally, and had a circular opening, fully 3 inches in diameter, on one side. They place their nests, very generally on the ground, in clumps of bamboos; and they construct them of dry bamboo leaves and twigs, and stems of plants, firmly and compactly interwoven. The exterior is rough and strong; the interior lined with soft vegetable fibres. They lay four eggs, very broad oval, glossy, with a pinky white ground, pretty thickly spotted all over with reddish and brownish purple; an egg figured measures 0·96 by 0·79 inch."

Dr. Jerdon says:—"I only procured one specimen, which was killed by a Lepcha when seated on her nest on the banks of the great Rungit river, about 1200 feet above the sea. The nest was composed chiefly of roots and other fibrous matter, with a few hairs, and contained three eggs of a faint greenish white, with a few reddish and some fawn-coloured spots."

Mr. Oates has found the species in Upper Pegu; and Mr. Hume writes as follows with respect to it:—

"Pegu specimens are absolutely identical with many others that I have from Sikhim, where the bird is very common. Blyth apparently considers (*Ibis*, 1866, p. 74) that our Indian bird is distinct from the Malaccan one, and should stand under his name *nigricollis*; as far south as Tavoy, at any rate, all are of one and the same species. Mr. Oates correctly points out (as I have previously noticed) that, in describing this species, both Dr. Jerdon (*B. of Ind. i.* p. 505) and Mr. Elliot (*Ibis*, 1870, p. 420) omit the conspicuous black patch, which on the centre of the lower abdomen surmounts the rich vermilion of the lower ventral region."

Mr. Oates remarks:—"I met with this bird in one ravine only in the evergreen forests, where I procured several specimens. I searched many precisely similar localities, but never again met with it. Two pairs that I measured varied as follows:—

"Length 7·3 to 7·55, expanse 13·5 to 14·5, tail from vent 1·55 to 1·65, wing 4·25 to 4·5, bill from gape 1·05 to 1·08, tarsus 1·6 to 1·7.

"The bill was black; the inside of the mouth dusky fleshy; irides dark coffee-brown; eyelids pale plumbeous-fleshy; legs fleshy pink; claws pinkish horny."

Mr. Davison has also met with the species in Tenasserim, at Amherst; and two examples procured by Dr. Cantor in Malacca are recorded by Messrs. Horsfield and Moore in their 'Catalogue.'

Body green above; head light chestnut-brown, the lores, eyebrows, sides of face, sides of neck, as well as the hind part of the latter and throat, black; upper tail-coverts rich cobalt; least wing-coverts rich cobalt; primaries with a broad band of white; under surface of body green; on the upper abdomen a black patch; the lower abdomen and under tail-coverts scarlet; thighs brown.

The figures in the Plate are of the natural size, drawn from specimens in my own collection.

Gould, John. 1877. "Hooded Pitta, *Pitta cucullata*, Hartl. [Pl. 82]." *The Birds of Asia* 5(XXX), -. <https://doi.org/10.5962/p.323107>.

View This Item Online: <https://www.biodiversitylibrary.org/item/121124>

DOI: <https://doi.org/10.5962/p.323107>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323107>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.