MYZORNIS PYRRHOPTERA, *Hodgs.*

J. Gould and H.C. Richter, del. & lith.

Hallward & Patten, imp.

MYZORNIS PYRRHOURA, *Hodgs.*

Red-tailed Myzornis.

- Myzornis pyrrhoura*, Hodgs. Blyth, Journ. Asiat. Soc. Beng. vol. xii. p. 984, and vol. xiv. p. 561.—Gray, Cat. of Spec. and Draw. of Mamm. and Birds presented to the Brit. Mus. by B. H. Hodgson, Esq., p. 74.—Blyth, Cat. of Birds in Mus. Asiat. Soc. Calcutta, p. 101.—Horsf. Cat. of Birds in Mus. East Ind. Comp. vol. i. p. 263.
- Yuhina pyrrhoura*, Gray and Mitch. Gen. of Birds, vol. i. p. 199. pl. 53, *Yuhina*, sp. 5.—Bonap. Consp. Gen. Av. 397, *Yuhina*, sp. 3.

THAT so beautiful a bird as the one here represented will become a general favourite with collectors there can be but little doubt, and one would fain hope that the time may arrive when some native Nepaulese, inspired with a love for ornithology, may give to the world a description of the habits and economy of this and the many other rare species which adorn his luxuriant country, or that some European may be favourably located for obtaining this desirable information: at present all that has been recorded respecting it is comprised in a line by Mr. Hodgson, who states that "this bird inhabits the northern and central hills of Nepaul."

I believe that the accompanying Plate represents both sexes; if so, it will be seen that they very nearly resemble each other; the only difference being in the smaller amount of red on the throat of the female and the lesser brilliancy of the red of her outer tail-feathers.

In the male the feathers of the crown are black, broadly margined with green; all the upper surface and wing-coverts dark green, becoming paler on the rump and upper tail-coverts; spurious wing orange-yellow; quills black, the apical half of the primaries margined with grey and tipped with white; the basal half of the secondaries margined with red, passing into greyish white on the apical portion and largely tipped with white; two central tail-feathers dark green tipped with black, the remainder fine crimson-red tipped with black; lores black; under surface green with a patch of dull red on the breast; vent and under tail-coverts paler red; bill black; feet fleshy.

The female is similar in colour, but the hues are not so bright, and she has merely an indication of the red on the breast and under tail-coverts.

The figures are of the natural size, and the plant is the *Cirrhopetalum cornutum*.

Gould, John. 1856. "Red-tailed Myzornis, *Myzornis pyrrhoura*, Hodgs. [Pl. 20]." *The Birds of Asia* 4(VIII), -. <https://doi.org/10.5962/p.323121>.

View This Item Online: <https://www.biodiversitylibrary.org/item/120503>

DOI: <https://doi.org/10.5962/p.323121>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323121>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.