CORACIAS AFFINIS, *McClell.*

Burmese Roller.

- Coracias affinis*, McClell. in Proc. of Zool. Soc., 1839, p. 164.—Gray, Gen. of Birds, vol. i. p. 62, pl. xxi. *Coracias*, sp. 8.—Blyth, Journ. Asiat. Soc. Beng., vol. xiv. p. 190, and vol. xix. p. 228.—Id. Cat. of Birds in Mus. Asiat. Soc. Calcutta, p. 51.—Horsf. and Moore, Cat. of Birds in Mus. East-Ind. Comp., vol. ii. p. 574.—Jerd. Birds of India, vol. i. p. 217.—Cab. et Heine, Mus. Hein., Theil ii. p. 118.—Gould in Proc. of Zool. Soc., 1859, pp. 150, 151.—Schomb. Ibis, 1864, p. 246.
- *katnas*, F. (B.) Hamilton, MS., vol. ii. p. 123.
- Katnas*, Gualpara, F. (B.) Hamilton.
- Nghet-kha*, Arracan, Capt. Abbott.

BOTH Mr. Blyth and Mr. Jerdon believe that, where the *Coracias Indica* and *C. affinis* inosculate, an interbreeding frequently takes place, since specimens partaking of the colouring of both are often met with. We know that such an intermixture occasionally occurs with some few other species; but this must be regarded as quite exceptional, and not the rule. When it does, it is mostly among the Gallinaceæ, whose hotter blood renders them more salacious than insessorial birds. On the interbreeding of the *C. Indica* and *C. affinis* Mr. Blyth has published some remarks which I take the liberty of transcribing, as I also do the accounts given of the latter species by Mr. Jerdon, Dr. Francis Buchanan Hamilton, and Sir Robert Schomburgk.

"The numerous specimens of Rollers from Assam, Arracan, and Tenasserim which I have seen," says Mr. Blyth, "all pertain strictly to this species, having the upper parts greener than in *C. Indica*, the neck and breast, devoid of the reddish-brown colour proper to the latter species, being purplish-dusky, varied with bright purple on the fore neck; and the entire under surface of the wing, except near the tips of the primaries, is deep purple; but I have obtained several specimens in the vicinity of Calcutta, and some from Tipperah, which present every gradation of plumage from one to the other of these species, and also some in the pure *C. affinis* plumage; from which I infer that, where found together in the same locality, they not unfrequently interbreed, and tend to merge into a single blended race. It may be further remarked that I have never seen an example of the true *C. affinis* with the broad purple terminal band to the tail, which distinguishes the adult of *C. Indica*; but I have seen this imperfectly developed in the mixed race, which latter has also commonly the fore part of the under surface of the wing intermingled purple and verditer."

Dr. F. Buchanan Hamilton mentions, in his MS., that "this bird is very common at Gualpara throughout the year, the natives considering it as distinct from the *Nil-khant*."

"This species," says Mr. Jerdon, "chiefly differs from the Indian one in the upper parts being greener, the neck and breast without any reddish-brown, being dusky purplish varied with bright purple, and in the wing being deep purple; it also wants the broad terminal purple band to the tail."

"The Burmese Roller is the only one found to the east of the Bay of Bengal, and in Assam, but it spreads into the Sunderbunds, Tipperah, and even the neighbourhood of Calcutta itself, where it mingles and interbreeds with the Indian Roller. Blyth has procured many specimens, which present every gradation of plumage from one to the other. The same intermingling of affined species takes place in the Green Pigeons of the south and north of India, and in several other birds. It does not differ in its habits or voice from the Indian Roller; those which I saw in Burmah were decidedly more wary and less familiar than the Indian bird."

Sir Robert Schomburgk states that "this handsome bird, the *Nook tackah* of the Siamese, is dressed in blues of all shades, azure and smaragdinous included, shading off into green and grey. His greatest enemies are the Crows. The bird is by no means uncommon in the gardens attached to the houses in Bangkok; but scarcely have the Crows in the vicinity discovered one before they make a simultaneous attack upon him. The Roller endeavours to avoid them, now by making gyrations in the air, and then by alighting upon the branch of a tree, uttering all the time pitiful cries; nor are his tormentors satisfied until he leaves the grounds. The beautiful blue wing-feathers of this bird, principally the primaries and scapularies, are exported to China for tiny fans; into which they are also manufactured in Siam, but they are not so well made as those that come from China."

The bird figured on the accompanying Plate, from Assam specimens, will be readily recognized as a distinct species by every ornithologist. It will be seen that there is not the slightest indication of the terminal band on the tail which is so conspicuous in *Coracias Indica*; the throat also is more blue, and in certain positions that colour is also visible on the abdomen, while in others it is inclined to verditer green.

Crown of the head and nape bluish green, bordered over the upper mandible and along the lores with buff, and above the eyes with bright verditer; back of the neck, back, and scapularies olive green; lower part of the back deep blue, each feather tipped with verditer blue; upper tail-coverts verditer blue; lesser wing-coverts deep blue; greater coverts of the primaries and spurious wing bright verditer blue; coverts of the secondaries dull greenish blue, becoming of a purer blue at the base; primaries deep rich blue, crossed towards the tip with a broad band of verditer blue, which is brightest on the external web, beyond this the tip is blue, shading into dull black at the extremity, shafts black; secondaries deep blue, with a band of verditer blue on the middle of the outer web only; tertiaries olive green; lores and throat purplish brown, passing into fawn-colour on the breast; the darker feathers of the sides and base of throat have a stripe of buff, and those in the centre of the throat a line of bright blue down the shaft; under wing-coverts deep blue; lower part of the abdomen greenish blue, passing into the dull verditer blue of the vent and under tail-coverts; two middle tail-feathers dull green; the remainder deep blue, largely tipped with pale verditer blue; shafts black; bill black; legs and feet dusky yellow; irides reddish-brown.

The sexes present no appreciable difference in their colouring

The figure of this fine species is of the size of life.

Gould, John. 1869. "Burmese Roller, *Coracias affinis*, McClell. [Pl. 55]." *The Birds of Asia* 1(XXI), -. <https://doi.org/10.5962/p.323171>.

View This Item Online: <https://www.biodiversitylibrary.org/item/115342>

DOI: <https://doi.org/10.5962/p.323171>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323171>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.