PHYLLORNIS CYANOPOGON, Temm.

PHYLLORNIS CYANOPOGON, Temm.

All Green Phyllornis.

- Phyllornis cyanopogon*, Temm. Pl. Col. 512. fig. 1.—Less. Traité d'Orn., p. 301.—Blyth, Cat. of Birds in Mus. Asiat. Soc. Calcutta, p. 213.—Gray and Mitch. Gen. of Birds, vol. i. p. 124, *Phyllornis*, sp. 6.—Moore in Proc. of Zool. Soc., part xxii. p. 280.
- *Malabaricus*, Blyth, Journ. Asiat. Soc. Beng., vol. xii. p. 957.
- Burong Daun* of the Malays in Sumatra, Raffles.
- Phyllornis mystacalis*, Moore in Proc. of Zool. Soc., part xxii. p. 280.
- *cyanopogon*, Blyth, Journ. Asiat. Soc. Beng., vol. xiv. p. 564.
- Chloropsis mystacalis*, Swains. in Lard. Cycl., Anim. in Menag., part iii. p. 96.—Gray and Mitch. Gen. of Birds, vol. i. p. 124, *Phyllornis*, sp. 7.
- *cyanopogon*, Jard. and Selby, syn. spec. *Chloropsis* in Ill. Orn., vol. ii. text to pl. 100. sp. 6.

No confusion need ever exist as to the identity of the bird here represented, since it differs from all its congeners in the absence of any blue or greenish blue on the shoulder, and in the almost uniform green colouring of its entire plumage, which is only relieved in the male by the black colouring of the throat and the usual stripe of blue springing from the base of the lower mandible.

The *Phyllornis cyanopogon*, which must rank as one of the smaller members of the genus, is undoubtedly the bird described under the name of *Phyllornis mystacalis* by the late Mr. Swainson, who appears not to have been aware of the localities in which it is found, but which we now know to be Malacca, and the Malay countries generally. Most of the specimens received in Europe are sent from Singapore. Temminck states that it is also found in Sumatra, but not in abundance.

The sexes of this species, like those of all the other members of the genus, may be at once distinguished by the absence of any black colouring on the throat of the female.

The male has the general plumage grass-green above and paler beneath; lores, chin, and throat deep velvety black, within which a narrow stripe of blue descends from the angle of the lower mandible; inner webs of the primaries and secondaries brownish black; tail green above, beneath slaty brown; bill black; irides brown; feet olive-grey.

The female differs in having the throat green instead of black, with, in some specimens, a slight blue moustache.

The Plate represents the two sexes of the natural size, on the *Ceropegia Cumingiana*.

Gould, John. 1861. "All-green Phyllornis, Phyllornis cyanopogon, Temm. [Pl. 18]." *The Birds of Asia* 3(XIII), -. <https://doi.org/10.5962/p.323233>.

View This Item Online: <https://www.biodiversitylibrary.org/item/118635>

DOI: <https://doi.org/10.5962/p.323233>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323233>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.