PARUS MINOR, Temm & Schleg

J. Gould and H. C. Fischer, del. et lith.

Heilmann & Witten, Imp.

PARUS MINOR, *Temm. et Schleg.*

Chinese Tit.

Parus minor, Temm. and Schleg. Faun. Japonica, Aves, p. 70. pl. xxxiii.—Bonap. Consp. Gen. Av., p. 229, *Parus*, sp. 9.

OVER what extent of habitat this well-defined species may range, will be for future ornithologists to determine, when the great country of China has become more open to scientific investigation ; at present we only know that the bird is a native of the districts in the neighbourhood of Shanghai and the Island of Japan, from both of which countries I possess examples. Although somewhat smaller in size, it is evidently the representative in China of the *Parus major* of Europe, from which it is distinguished by the total absence of yellow on the flanks, and the greater amount of white on the outer tail-feathers, as it is from the *Parus cinereus* of India by its yellow nape. MM. Temminck and Schlegel have given figures of this bird in the volume "Aves" of Siebold's interesting work on the Fauna of Japan, but afford us no information as to its habits and economy ; we may infer, however, that they resemble those of its near allies, which, as is well known, obtain their insect food among the branches of thickly-foliaged trees ; that they are pert, restless, and active in their manners, and stationary or non-migratory.

The sexes appear to assimilate most closely in their colouring, which may be thus described :—

Crown of the head, nape, throat, band on the sides of the neck, chest, and an irregular mark down the centre of the abdomen glossy bluish black ; cheeks and ear-coverts white ; centre of the back yellowish olive, separated from the black of the nape by a mark of white ; scapularies, lower part of the back, and upper tail-coverts blue-grey ; wing-coverts black, tipped with white, and margined with blue-grey ; primaries and secondaries slaty black, the former narrowly margined at the base with blue-grey, and towards the apex with white, the latter more broadly with dull white ; sides of the chest and flanks creamy ; tail slaty black, margined with blue-grey, the outer feather white, margined internally with brown, and the next on each side with a wedge-shaped mark of white at the tip ; bill bluish black ; irides dark brown ; feet blackish blue.

The figures are the size of life. The plant is a species of *Uvularia*, copied from one of the fine collections of botanical drawings at the East India House.

Gould, John. 1858. "Chinese Tit, *Parus minor*, Temm. et Schleg. [Pl. 56]." *The Birds of Asia* 2(X), -. <https://doi.org/10.5962/p.323242>.

View This Item Online: <https://www.biodiversitylibrary.org/item/118636>

DOI: <https://doi.org/10.5962/p.323242>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323242>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.