EUPLOCAMUS LINEATUS.

EUPLOCAMUS LINEATUS, Vigors.

Lineated Pheasant.

Lophophorus Cuvieri, Temm. Pl. Col., v. pl. 1 (1820, hybrid).

Lineated Pheasant, Lath. Gen. Hist. B., viii. p. 201.

Phasianus lineatus, Vigors, P. Z. S., 1831, p. 24 (ex Lath. MS.).—Jerdon, B. Ind., iii. p. 531.—Beavan, Ibis, 1868, p. 381.

Gennæus lineatus, Wagler, Isis, 1832, p. 1228.

Phasianus Reynaudi, Lesson, in Bélanger, Voy. Ind. Orient. Zool., p. 276, pls. 8, 9.

Phasianus fasciatus, M'Clell. Calcutta Journ. N. H., ii. p. 146, pl. 3.

Lophophorus leucomelas, Gray, List of Genera, 1840, p. 60.

Alectrophasis leucomelanos, Gray, List of Genera, 1841, p. 78.

Gallophasis fasciatus, Gray, Gen. B., iii. p. 498.

Euplocamus lineatus, Gray, List of Genera, p. 78.—Blyth, Cat. B. Mus. A. S. B., p. 244.—Sclater, P. Z. S., 1863, p. 120.—Id. text to Wolf's Zool. Sketches, 2nd ser. pl. 38.—Blyth, Ibis, 1867, p. 153.—Gray, List of Gallinæ, p. 34.—Gray, Hand-l. B., ii. p. 260.—Elliot, Monogr. Phasian., ii. pl. xxxiii.

Grammatoptilus lineatus, Reichenb. Syst. Av., p. 30.

If there is one species more than another which unites the two sections of the genus *Euplocamus* (that is to say the Kaleeges) with the Silver Pheasant, it is the present bird. The largely developed facial wattle, the finely pencilled plumage extending over the whole upper surface, and the form of the tail incline it to the Silver Pheasant (*Gallophasis nycthemerus*); but then the latter species and its immediate allies are distinguished by their bright red legs, while in the present bird they are bluish grey. This may seem to many a character of slight importance; but to my mind it is not so, for I find the colour of the legs in these Pheasants of great differential value. Those who know our Silver Pheasant, the beautiful *Euplocamus Swinhoei*, or its elegant congener *E. praelatus*, and the still more curious *Lobiophasis Bulweri*, will remember that all these birds have delicate legs of a bright red colour, whereas the present bird and all the Kaleeges have them dark-coloured. I lay the greater stress on this point because Mr. Elliot, in his noble monograph of the family, has given bright red legs to *E. albocristatus*; but this must be an oversight, for I find that Dr. Jerdon gives them as "dark horny," which accords with my own experience. Again, in the description of the present species, he gives the legs as red, whereas they are figured brown in the plate; and that the latter is correct will be found on a reference to Wolf's plate in the 'Zoological Sketches,' taken from the living bird.

As is the case with most of the members of the genus *Euplocamus*, of which allied species interbreed on the confines of their respective ranges, the Lineated Pheasant is often found crossed with *E. Horsfieldi*; and were hybrids to be noticed, it would stand as *Euplocamus Cuvieri*, as our first notice of the species was in 1820, when Temminck figured a hybrid specimen. Concerning this I may with advantage quote Mr. Blyth's remarks:—

"The *Lophophorus Cuvieri* represents one of the hybrid races referred to, between *Gallophasis lineatus* and *G. Horsfieldi*. These completely pass one into the other in the province of Arakan, whence some living specimens have been received by the Zoological Society. In like manner *G. albocristatus* and *G. melanonotus* interbreed in the intermediate province of Nipâl, *G. melanonotus* being the species inhabiting Sikkim and Butan, where most assuredly *G. lineatus* is unknown, the latter inhabiting southward of the range of *G. Horsfieldi*, i. e. in Pegu and the Tenasserim provinces, where I have personally observed it in the forests."

Captain Beavan writes as follows:—"Dr. Jerdon mentions the singular drumming noise made by the male. With respect to *Euplocamus lineatus*, which is, in my opinion, a true Kaleege, I may mention that the Burmese take advantage of this habit of the birds, and by imitating it with a rough kind of machine catch numbers of the latter species. It is like the crow of a cock, a kind of challenge uttered by one male to excite another to a battle—'in defiance of his fellows.'"

Mr. Elliot further observes:—"Mr. Blyth informs me that the natives snare a cock bird of this species, and secure him near the jungles, when his cries serve to bring the males who may be within hearing into the open, looking for their antagonist, whom they immediately attack, thus giving the concealed gunner an opportunity to shoot them; and in this way many birds are obtained. Of course it is only the cocks which are procured, the hens never paying any attention to the crowing, but remaining quietly in the forest."

The following descriptions are transcribed from Mr. Elliot's 'Monograph':—

"*Male*.—Head with a long occipital black crest. Rest of head, neck, and entire upper parts black, finely vermiculated with white. Inner web of middle tail-feathers buff; outer black, crossed with white zigzag lines; rest of tail-feathers black, with diagonal white lines. Primaries brown, crossed with fine white lines. Breast and underparts bluish black. Centre of flank-feathers white. Bare skin of face red.

"*Female*.—Head and occipital crest dark brown, mottled with light brown; neck and upper part of back light brown, with V-shaped white marks near the end of the feathers, divided from the brown edges by a line of black; rest of back, wings, and upper tail-coverts greyish brown, finely mottled with blackish brown. Chin whitish brown. Breast and flanks rufous brown; shafts and centre of feathers white, with black lines around their edges. Abdomen and rest of parts black, the feathers tipped with chestnut. Central tail-feathers buff, mottled with black on inner webs; remainder black, irregularly crossed with white lines. Bill: upper mandible black, lower horn-colour. Feet lead-colour."

The Plate represents the male and female of this Pheasant, of about three quarters of the natural size.

Gould, John. 1875. "Lineated Pheasant, *Euplocamus lineatus*, Vigors. [Pl. 14]." *The Birds of Asia* 7(XXVII), -. <https://doi.org/10.5962/p.323259>.

View This Item Online: <https://www.biodiversitylibrary.org/item/122491>

DOI: <https://doi.org/10.5962/p.323259>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323259>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.