CHRYSOCOLAPTES XANTHOCEPHALUS, *Walden.*

CHRYSOCOLAPTES XANTHOCEPHALUS, *Walden.*

Negros Yellow-faced Woodpecker.

Chrysocolaptes xanthocephalus, Walden & Layard, Ibis, 1872, p. 99, pl. iv.—Walden, Trans. Zool. Soc. ix. p. 147.—Sharpe, Trans. Linn. Soc., 2nd series, Zoology, vol. i. part 5.

THE Philippine Islands seem to possess several peculiar species of Woodpeckers, all belonging to the same type. Thus *C. hæmatribon* inhabits Luzon, *C. xanthocephalus* is found in Negros, while *C. maculiceps* is the representative form in Basilan, and *C. erythrocephalus* in Palawan; both these latter species were discovered by Dr. Steere. There is also a fifth species, *C. lucidus*, from Luzon, with which, Lord Tweeddale thinks, Mr. Sharpe's *C. maculiceps* may be identical.

The subject of the present article was discovered in the island of Negros by Mr. L. C. Layard, who, however, only succeeded in capturing a female bird. This, however, was so evidently distinct that Lord Tweeddale had no difficulty in separating it specifically. He remarks:—"The carmine dorsal colouring of this species closely resembles that of *Ch. carlotta* (Malh.), *Ch. hæmatribon* (Wagler), and *Brachypternus erythronotus* (V.) apud Malherbe. The male bird may prove to possess a red head, as in the rest of the genus." That this latter supposition was correct has now been proved by Dr. Steere's specimen, which was also procured in the island of Negros. He says:—"I saw more of this Woodpecker, which was not rare; but I had the greatest difficulty in getting about the forests in Negros, owing to the obstruction caused by the trees blown down in a recent typhoon."

The following is the description of the male bird given by Mr. Sharpe:—

"General colour above ruddy crimson, all the feathers olive-brown at the base, the upper tail-coverts entirely of the latter colour, with a crimson wash on their margins; scapulars and wing coverts exactly resembling the back, the bastard wing, primary-coverts, and quills brown, externally yellowish olive, the secondaries brown, inclining to olive on the outer webs, which are externally crimson, like the back; all the quills spotted with white on the inner web, much more largely on the secondaries; tail-feathers dark brown, with a slight wash of olive on the outer webs; crown of head bright crimson, with yellow bases to the feathers; lores, eyebrow, and entire sides of face bright yellow, as also the throat, which is bordered on each side with a narrow black moustachial line, and has also another down the centre of the throat; sides of neck yellow, with three lines of black running down from behind the ear-coverts; neck all round and fore neck scaly in appearance, the feathers being yellowish buff, fringed with black; remainder of under surface yellowish buff, the sides of the body somewhat streaked with olive-brown; under wing-coverts dull fulvous, with olive-brown margins to the feathers, which are also slightly washed with red; bill stone-brown (in skin); legs yellowish, claws black; 'iris carmine' (*Steere*). Total length 10·5 inches, culmen 1·6, wing 5·6, tail 3·8, tarsus 1·05."

The figure of the male bird in the Plate is taken from the skin procured by Dr. Steere at San Bernardino, Negros, and kindly lent to me by him. For the loan of the female I have to thank the Marquis of Tweeddale, who has been so good as to forward it to me for the purpose of the present work. The two figures are about the natural size.

Gould, John. 1877. "Negros Yellow-faced Woodpecker, *Chrysocolaptes xanthocephalus* [Pl. 37]." *The Birds of Asia* 6(XXX), –.

<https://doi.org/10.5962/p.323296>.

View This Item Online: <https://www.biodiversitylibrary.org/item/122488>

DOI: <https://doi.org/10.5962/p.323296>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323296>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.