CUCULUS HYPERYTHRUS, Gould.

CUCULUS HYPERYTHRUS, *Gould.*

Rufous-bellied Cuckoo.

Cuculus hyperythrus, Gould in Proc. of Zool. Soc. April 22, 1856.

CHINA has hitherto been, as it were, a sealed country to the scientific explorer, consequently all the natural productions of that part of the world have been regarded with especial interest. That its feathered inhabitants partake of a triple character is certain; in the first place, it possesses several species of birds which are precisely identical with some of those of the British Islands; in the next, it is tenanted by several additional species of the same forms; and in the third, by forms which are peculiarly its own: the present species, which is a true *Cuculus*, pertains to the second of these divisions. I have never seen examples from any other country; neither, so far as I am aware, has it ever before received a specific appellation; for I have looked in vain for a description of it in every work to which I could obtain access, particularly in the valuable monograph of the genus lately published by Mr. Blyth. In size it is rather less than the Common Cuckoo of Europe, and it is altogether less elegant in its general contour. The rufous colouring of the breast and under surface, and the black marks on the throat and cheeks, are characters seldom seen among the *Cuculidæ*, and by which it may at once be distinguished.

A fine specimen of this bird, now at the British Museum, but which was formerly in my possession, was shot at Shanghai, and this, I regret to say, is all I know respecting it.

Crown of the head, all the upper surface and wings dark slate-grey; spurious wings white; lores, ear-coverts, moustache and spot on the chin black; throat white, with a fine line of brown down the shaft of each feather; under surface dull rusty red; tail grey, crossed by two narrow irregular bands of black bordered with brown, and by a very broad band of black near the extremity, the tip being reddish brown; upper mandible black; lower mandible and feet yellow.

Total length, $11\frac{1}{2}$ inches; bill, $1\frac{1}{8}$; wing, 8; tail, $6\frac{1}{2}$.

The figures are of the natural size.

Gould, John. 1856. "Rufous-bellied Cuckoo, *Cuculus hyperythrus* [Pl. 43]." *The Birds of Asia* 6(VIII), -. <https://doi.org/10.5962/p.323307>.

View This Item Online: <https://www.biodiversitylibrary.org/item/122488>

DOI: <https://doi.org/10.5962/p.323307>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323307>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.